

Due to the relative simplicity of the technique, many anesthesiologists consider intravenous regional anesthesia to be both virtually foolproof and, barring tourniquet failure, a guarantee of adequate analgesia. However, many studies have shown that failure to provide good analgesia is not uncommon and may be related to the particular technique used. Inadequate analgesia (defined differently in each study) occurs^{7,13,14,16,17} at rates varying from 7% to 15%. Of particular interest is the work of Sorbie and Chacha,⁷ which noted a much higher percentage of unsuccessful blocks following proximal cubital injections (22.7%) as compared with distal hand injections (4.1%). Hollingworth *et al.*¹⁶ revealed the apparent importance of individual technique on the success of the block. Aside from specifying uniform drugs and dosages, their study¹⁶ does not report the precise techniques used in providing intravenous regional anesthesia. Inadequate analgesia occurred with widely different frequencies among eight physicians—from 0% to 31%. These physicians might have used different injection rates, tourniquet pressures, *etc.*, and thereby affected the number of successful blocks.

In conclusion, our data indicate that injections for intravenous regional anesthesia should be made over at least 90 s, into a distal vein, with a tourniquet pressure of at least 300 mmHg, and after exsanguination with an Esmarch's bandage. By avoiding MVP and thus keeping all the anesthetic isolated in the limb, a decrease in the incidence and severity of toxic symptoms and an increased percentage of successful blocks should result.

REFERENCES

1. Heath ML: Bupivacaine toxicity and Bier blocks. *ANESTHESIOLOGY* 59:481, 1983
2. Rosenberg PH, Kalso EA, Tuominen MK, Linden HB: Acute bupivacaine toxicity as a result of venous leakage under the

tourniquet cuff during a Bier block. *ANESTHESIOLOGY* 58:95-98, 1983

3. El-Hassan KM, Hutton P, Black AMS: Venous pressure and arm volume changes during simulated Bier's block. *Anaesthesia* 39: 229-235, 1984
4. Davies JAH, Hall ID, Wilkey AD, Smith JE, Walford AJ, Kale VR: Intravenous regional analgesia—The danger of the congested arm and the value of occlusion pressure. *Anaesthesia* 39:416-421, 1983
5. Emmett C: More trouble with Bier's block. *Anaesthesia* 38:390, 1983
6. Davies JAH, Gill SS, Weber JCP: Intravenous regional analgesia using bupivacaine. *Anaesthesia* 36:331, 1981
7. Sorbie C, Chacha P: Regional anaesthesia by the intravenous route. *Br Med J* 1:957-960, 1965
8. Fleming SA, Veiga-Pires JA, McCutcheon RM, Emanuel CI: A demonstration of the site of action of intravenous lignocaine. *Can Anaesth Soc J* 13:21-27, 1966
9. Raj PP, Garcia CE, Borleson JW, Jenkins MT: The site of action of intravenous regional anesthesia. *Anesth Analg* 51:776-786, 1972
10. Kalso E, Tuominen M, Rosenberg PH, Alila A: Bupivacaine blood levels after intravenous regional anesthesia of the arm. *Reg Anaesth* 5:81-84, 1982
11. American Heart Association, Kirkendall WM, Burton AC, Epstein FH, Freis ED: Recommendations for human blood pressure determination by sphygmomanometers. *Circulation* 36:980-988, 1967
12. Finegan BA, Bukht MD: Venous pressures in the isolated upper limb during saline injection. *Can Anaesth Soc J* 31:364-367, 1984
13. Ware RJ: Intravenous regional analgesia using bupivacaine. *Anaesthesia* 30:817-822, 1975
14. Holmes CM: Intravenous regional analgesia—A useful method of producing analgesia of the limbs. *Lancet* 1:245-247, 1963
15. Bell HM, Slater EM, Harris WH: Regional anesthesia with intravenous lidocaine. *JAMA* 186:544-549, 1963
16. Hollingworth A, Wallace WA, Dabir R, Ellis SJ, Smith AFM: Comparison of bupivacaine and prilocaine used in Bier block—A double blind trial. *Injury* 13:331-336, 1982
17. Kennedy BR, Duthie AM, Parbrook GD, Carr TL: Intravenous regional analgesia: An appraisal. *Br Med J* 1:954-957, 1965

Anesthesiology
65:320-322, 1986

Bradycardia Delays the Onset of Action of Intravenous Atropine in Infants

G. ZIMMERMAN, M.D.,* DAVID J. STEWARD, M.B., F.R.C.P.(C)

Atropine, a vagolytic drug, is commonly used in pediatric patients to prevent or reverse bradycardia.¹ Rou-

tine preoperative administration of an anticholinergic is decreasing in adults,² and some have recommended that this practice be discontinued in pediatric patients.³ Atropine is then administered only if indicated (*e.g.*, to treat bradycardia or precede succinylcholine). To achieve a rapid effect in treating bradycardia, iv atropine is often used. Sometimes a long delay in restoring normal heart rate has been observed, especially if severe bradycardia is present. This delay may be caused by the low cardiac output and prolonged circulation time associated with severe bradycardia.

To study this further, we collected data on pediatric patients to examine the relationship between the heart

* Resident in Anaesthesia, University of British Columbia.


† Professor of Anaesthesia, University of British Columbia, and Anaesthetist-in-Chief, British Columbia Children's Hospital.

Received from the Department of Anaesthesia, University of British Columbia and British Columbia Children's Hospital, Vancouver, British Columbia. Accepted for publication April 7, 1986.

Address reprint requests to Dr. Steward: Department of Anaesthesia, British Columbia Children's Hospital, 4480 Oak Street, Vancouver, British Columbia V6H 3V4, Canada.

Key Words: Anesthesia: pediatric. Parasympathetic nervous system: atropine. Heart: arrhythmia, bradycardia.

FIG. 1. Relationship between initial heart rate and delay in onset of action of intravenous atropine in 14 infants less than 2 yr of age (Group 2).



rate at the time of iv injection of atropine and the delay in the onset of action on the heart.

METHODS

This study was approved by the Human Investigation Review Committee. It is a common practice with pediatric patients to omit im premedication and to commence an inhalation induction with halothane. Once the patient is anesthetized, a venipuncture is performed and iv atropine administered.⁴ Following this procedure we also recorded a strip of the electrocardiogram tracing. This tracing was marked with the time at which atropine (0.02 mg/kg) was injected rapidly into a vein on the dorsum of the hand. The tracing was later examined to determine the heart rate at the time of atropine administration and the delay in seconds until a sustained increase in heart rate occurred. This point was defined at the R wave after which four successive R-R intervals were shorter than the preceding series of R-R intervals.

RESULTS

Data were obtained from a total of 29 patients. After preliminary analysis, when it became apparent that the results in infants were different from those in older children, the patients were divided into two groups: Group 1 comprised 15 children aged 2 to 10 yr (mean age 4.5 yr \pm 2.0 yr SD). Group 2 comprised 14 infants aged under 2 yr (mean age 8.8 months \pm 5.2 months SD). The children in Group 1 demonstrated no relationship between initial heart rate and the delay in drug action. The infants

in Group 2 showed an inverse relationship between initial heart rate and the delay in drug action. The slower the initial heart rate, the longer was the delay in onset of drug action (fig. 1).

DISCUSSION

Our results show that the time of onset of cardiac vagolytic action of iv atropine in infants is significantly influenced by the heart rate at the time of drug administration. In older children no such relationship was demonstrated. These results might be anticipated from a knowledge of the factors that influence the distribution of a drug from its iv injection site to its site of action.

The speed of onset of the effect of iv atropine will be dependent on the circulation time from the site of injection to the postganglionic parasympathetic nerve endings in the cardiac plexi.⁵ The circulation time is related inversely to the cardiac output, and a decrease in cardiac output is followed by an increased circulation time.⁶ Cardiac output is a function of heart rate and stroke volume. Small infants have a relatively fixed stroke volume due to the presence of less compliant ventricles.⁷ The cardiac output of the infant is therefore closely related to the heart rate. In children, as in adults, the cardiac output is less dependent on the heart rate, as increased diastolic filling of the ventricle and consequent increased stroke volume can partially compensate for a reduced heart rate. Thus, beyond infancy, circulation time depends less on heart rate. Our findings are consistent with this.

We conclude that in the infant, bradycardia should be treated promptly, before a serious decrease in cardiac

output results. As the common causes of intraoperative bradycardia are hypoxia or vagal activity, such therapy should consist of ventilation with oxygen and iv atropine. To obtain a rapid response, iv atropine should be administered before marked bradycardia is present.

REFERENCES

1. Smith RM: Preparing children for operation, *Anesthesia for Infants and Children*, 4th edition. St. Louis, Mosby, 1980, p 98
2. Holt AT: Premedication with atropine should not be routine. *Lancet* 2:984-985, 1962
3. Hendriks PH, Govaerts MGM: Pediatric premedication: Atropine or not. *Acta Anaesthesiol Belg [Suppl]* 31:195-196, 1980
4. Gregory GA: Induction of anesthesia, *Pediatric anesthesia*. New York, Churchill-Livingstone, 1983, p 441
5. Morton HJ, Thomas ET: Effect of atropine on the heart rate. *Lancet* 2:1313, 1958
6. Mountcastle VB: The heart as a pump, *Medical Physiology*, 14th edition. St. Louis, Mosby, 1980, pp 1000-1001
7. Friedman WF: The intrinsic properties of the developing heart. *Prog Cardiovasc Dis* 15:87-111, 1972

Anesthesiology
65:322-325, 1986

Effect of Droperidol Pretreatment on Postanesthetic Vomiting in Children Undergoing Strabismus Surgery

J. LERMAN, M.D., F.R.C.P.C.,* S. EUSTIS, M.D.,† D. R. SMITH, M.D., F.R.C.S.C.‡

Vomiting is a frequent and often disturbing complication following strabismus surgery, particularly in children.¹⁻⁵ Droperidol, a potent antiemetic, reduces the incidence of vomiting after many types of surgery, including strabismus surgery.⁶⁻¹⁴ Abramowitz *et al.* reported that intravenous droperidol 0.075 mg · kg⁻¹ reduces the incidence of vomiting in children undergoing strabismus repair from 85% (control group) to 43% when administered after manipulation of the eye.¹ However, Meyers and Tomeldan observed that intramuscular droperidol 0.10 mg · kg⁻¹ reduces the incidence of vomiting after eye surgery in adults to 10% when administered preoperatively and before manipulation of the eye.¹⁵ The lower incidence of vomiting in the latter study may be attributed, in part, to the more effective antiemetic action of droperidol when administered before manipulation of the eye rather than afterward. We speculated that if intravenous droperidol were administered before manipulation of the eye, the incidence of vomiting after strabismus repair in children might be reduced to a similar extent as in adults.¹⁵ Therefore, we determined the incidence of

vomiting after strabismus repair in children who received intravenous droperidol pretreatment, or one of the routine postanesthetic analgesics used in our institution: rectal acetaminophen or intramuscular codeine.

METHODS

With approval from the Committee on Human Research, a prospective, randomized study was undertaken. Informed written consent was obtained from the parents of 100 children scheduled for elective strabismus repair. The children were ASA Physical Status 1 or 2, fasting, unpremedicated, and older than 2 yr of age. Children with a history of motion sickness or vomiting after previous strabismus surgery were not excluded from the study.

The children were randomly assigned to one of three treatment groups: intravenous droperidol (0.075 mg · kg⁻¹) (n = 31); rectal acetaminophen (10 mg · kg⁻¹) (n = 35); or intramuscular codeine (1.5 mg · kg⁻¹) (n = 34). Droperidol was given at induction of anesthesia immediately after succinylcholine. Acetaminophen and codeine were given in the postanesthetic room (PAR) when the children were arousable. Acetaminophen was administered instead of a placebo to alleviate any possible discomfort after surgery. Intramuscular codeine was included as the third treatment after it was incriminated as a possible cause of vomiting after strabismus repair by one of our ophthalmologists. §

After a precordial stethoscope, electrocardiogram, blood pressure cuff, and Doppler probe were applied, general anesthesia was induced with intravenous thiopental (5 mg · kg⁻¹), atropine (0.02 mg · kg⁻¹), and succinyl-

* Assistant Professor, Director of Anaesthesia Research.

† Fellow in Ophthalmology.

‡ Assistant Professor of Ophthalmology.

Received from the Departments of Anaesthesia and Ophthalmology, and the Research Institute, The Hospital for Sick Children, University of Toronto, Toronto, Ontario. Presented in part, at the Annual Meeting of the American Society of Anesthesiologists, San Francisco, California, October 1985. Accepted for publication April 21, 1986.

Address reprint requests to Dr. Lerman: Department of Anaesthesia, The Hospital for Sick Children, 555 University Avenue, Toronto, Ontario M5G 1X8.

Key words: Anesthesia: pediatric; ophthalmologic. Narcotics: codeine. Vomiting: antiemetics; droperidol.

§ Smith DR: Personal communication.