

can be judged. When the blood volume is reduced by less than 10 per cent, the arterial pressure is maintained primarily by the non-neural factors. With a volume reduction between 10 and 25 per cent, the sympathetic activity, the inhibition of vagi, and the non-neural factors are of about equal importance. When the reduction of blood volume is greater than 25 per cent, the sympathetic activity becomes the most significant compensatory factor. (Chien, S., and Billig, S.: *Effect of Hemorrhage on Cardiac Output of Sympathectomized Dogs*, *Amer. J. Physiol.* 201: 475 (Sept.) 1961.)

**AURICULAR PRESSURE** In five dogs and one human being, right auricular, intrathoracic (esophageal), intratracheal, and arterial pressures were measured during spontaneous respiration, manually controlled respiration, IPBB, and positive-negative phase respiration. Right auricular pressure is low during spontaneous inspiration, and elevated during expiration. The pressures in the right atrium show a direct relationship to those in the respiratory tract. With high insufflation pressures, the pressure in the right atrium is markedly increased and venous filling is diminished. The addition of a negative phase during expiration causes restitution to almost physiological conditions. The right auricular pressure curves resemble those obtained during spontaneous respiration. (Hanquet, M., and Lefebvre, L.: *Respiration Contrôlée et Pression Auriculaire Droite*, *Acta Anaesth. Belg.* 11: 381 (Dec.) 1960.)

**VENTRICULAR TACHYCARDIA** Quinidine and procaine amide are the most effective drugs for treatment of ventricular tachycardia, and the majority of episodes respond to these agents. However, in large doses, these drugs significantly depress myocardial contractility with resultant heart failure and shock. The object of electric countershock, whether the underlying arrhythmia is ventricular tachycardia or fibrillation, is to achieve simultaneous, uniform depolarization of all parts of the ventricle, thereby promptly extinguishing ectopic foci and permitting the sinus node to resume as pacemaker. When countershock is applied directly to the exposed heart, rela-

tively small amounts of current (50 to 150 volts) are employed; greater current, usually around 350 to 440 volts, is necessary when countershock is applied externally. The obvious advantages of electric countershock are that it is easily administered, the results are immediate, there is no lasting depression of cardiac function, and there are no serious after-effects. Combined with the safety factors of effective external pacemakers and external cardiac massage, this new technique may be the treatment of choice for ventricular tachycardia when anti-arrhythmic drugs have proven ineffective. (Alexander, S., and others: *Use of External Electric Countershock in the Treatment of Ventricular Tachycardia*, *J. M. A.* 177: 916 (Sept. 30) 1961.)

**STARLING'S LAW** Changes in effective left ventricular end-diastolic pressure were determined by measuring left ventricular pressure with a catheter introduced through the atrial septum, and intraesophageal pressure with a balloon. The activity of the autonomic nervous system was reduced with an infusion of trimethaphan administered at a constant rate. The reduction in circulatory reactivity was indicated by the absence of arterial pressure response to the cold pressor test. Measurements of cardiac output, stroke volume, left ventricular work, power, tension-time index, and the duration and mean rate of left ventricular ejection were carried out before, during and upon completion of a transfusion of 1,500 ml. of the subject's own blood. Transfusion resulted in a significant elevation of left ventricular end-diastolic pressure in each subject. As this occurred, left ventricular performance also became augmented. These data are consistent with the hypothesis that the end-diastolic pressure is an important determinant of the characteristics of ventricular contraction and that Starling's Law of the heart is applicable to man. (Braunwald, E., and others: *Studies on Starling's Law of the Heart. V. Left Ventricular Function in Man*, *J. Clin. Invest.* 40: 1882 (Oct.) 1961.)

**BARORECEPTORS** The relationship between pulmonary arterial pressure and the impulse activity recorded from pulmonary arterial baroreceptor fibers has been studied in