

22. Paulet, G.: Activité cholinérasique et fonctionnement des centres respiratoires, *J. Physiol. (Paris)* 48: 915, 1956.
23. Randall, L. O., and Lehmann, G.: Pharmacological properties of some neostigmine analogs, *J. Pharmacol. Exp. Ther.* 99: 16, 1950.
24. Randall, L. O.: Anticurare action of phenolic quaternary ammonium salts, *J. Pharmacol. Exp. Ther.* 100: 83, 1950.
25. Wescoe, W. C., Riker, W. F., Jr., and Beach, V. L.: Studies on the inter-relationships of certain cholinergic compounds. III. The reactions between 3-acetoxy phenyltrimethylammonium methylsulfate, 3-hydroxy phenyltrimethylammonium bromide and cholinesterase, *J. Pharmacol. Exp. Ther.* 99: 265, 1950.
26. Wescoe, W. C., and Riker, W. F., Jr.: Pharmacology of anticurare agents, *Ann. N. Y. Acad. Sci.* 54: 438, 1951.
27. Randall, L. O., and Jampolsky, L. M.: Pharmacology of drugs affecting skeletal muscle, *Amer. J. Phys. Med.* 32: 102, 1953.
28. Riker, W. F., Jr., and Wescoe, W. C.: The direct action of prostigmine on skeletal muscle; its relationship to the choline esters, *J. Pharmacol. Exp. Ther.* 88: 58, 1946.
29. Smith, C. M., Cohen, H. L., Palikan, E. W., and Unna, K. R.: Mode of action of antagonists to curare, *J. Pharmacol. Exp. Ther.* 105: 391, 1952.
30. Nastuk, W. L., and Alexander, J. T.: The action of 3-hydroxyphenylidimethylammonium (Tensilon) on neuromuscular transmission in the frog, *J. Pharmacol. Exp. Ther.* 111: 302, 1954.
31. Riker, W. F., Jr.: Excitatory and anti-curare properties of acetylcholine and related quaternary ammonium compounds at the neuromuscular junction, *Pharmacol. Rev.* 5: 1, 1953.
32. Riker, W. F., Jr., Roberts, J., Standaert, F. G., and Fujimori, H.: The motor nerve terminal as the primary focus for drug-induced facilitation of neuromuscular transmission, *J. Pharmacol. Exp. Ther.* 121: 286, 1957.
33. Riker, W. F., Jr., Werner, G., Roberts, J., and Kuperman, A.: Pharmacologic evidence for the existence of a presynaptic event in neuromuscular transmission, *J. Pharmacol. Exp. Ther.* 125: 150, 1959.
34. Blaber, L. C., and Bowman, W. C.: A comparison between the effects of edrophonium and choline in the skeletal muscles of the cat, *Brit. J. Pharmacol.* 14: 456, 1959.
35. Pelikan, E. W.: In discussion of: Synthetic curare-like agents and their antagonists, *Ann. N. Y. Acad. Sci.* 54: 474, 1951.
36. Waser, P. G.: Autoradiographie von Endplatten mit Radiocurarin nach Denervierung, Prostigmin oder SKF 525A, *Helv. Physiol. Acta.* 16: 171, 1958.

---

### Anesthesia

**SPINAL ANESTHESIA** There is a small but definite increase in femoral vein oxygen tension and a small but definite decrease in arteriovenous oxygen difference after spinal block in patients with peripheral vascular disease. This is presumably due to an increase in flow to the extremity, primarily to muscle. The mechanism is presumably that of a preganglionic block causing an arterial and arteriolar dilatation. (Cerrilli, G. J., and Engell, H. C.: *The Effect of Spinal Anesthesia on Femoral Vein Oxygen Tension, Surgery* 60: 668 (Sept.) 1966.)

**OPIATES IN OBSTETRICS** In a double blind study of 471 patients using four treatment groups nearly similar in all respects to insure random sampling, it was found that anileridine and meperidine produced exactly comparable effects, with 30 mg. anileridine being equivalent to 75 mg. meperidine. Apgar ratings were essentially the same in all four treatment groups. No significant side effects were noted with either drug and no real advantage was obtained when either were combined with a phenothiazine. (Cavanagh, D., and others: *Comparison of Anileridine and Meperidine as Obstetric Analgesia, Amer. J. Obstet. Gynec.* 96: 213 (Sept.) 1966.)