

- ard for the estimation of methoxyflurane levels in blood, *Brit. J. Anaesth.* 38: 29, 1966.
9. Courtin, R. F., Bickford, R. G., and Faulconer, A., Jr.: The classification and significance of electro-encephalographic patterns produced by nitrous oxide-ether anesthesia during surgical operations, *Proc. Mayo Clin.* 25: 197, 1950.
  10. Faulconer, A., Jr.: Correlation of concentrations of ether in arterial blood with electro-encephalographic patterns occurring during ether-oxygen and during nitrous oxide, oxygen and ether anesthesia of human surgical patients, *ANESTHESIOLOGY* 13: 361, 1952.
  11. Courtin, R. F.: Electroencephalographic and clinical observations with trichlorethylene and nitrous oxide anesthesia, *Dallas Med. J.* 41: 613, 1955.
  12. Pearce, W. C., Knott, J. R., and Bjurstrom, R. O.: Studies on nitrous oxide, meperidine and levallorphan with unipolar electroencephalography, *ANESTHESIOLOGY* 18: 310, 1957.
  13. Possati, S., Faulconer, A., Jr., Bickford, R. G., and Hunter, R. C.: Electroencephalographic patterns during anesthesia with cyclopropane: correlation with concentration of cyclopropane in arterial blood, *Anesth. Analg.* 32: 130, 1953.
  14. Galla, S. J., Rocco, A. G., and Vandam, L. D.: Evaluation of the traditional signs and stages of anesthesia: an electroencephalographic and clinical study, *ANESTHESIOLOGY* 19: 328, 1958.
  15. Clowes, G. H. A., Jr., Kretschmer, H. E., McBurney, R. W., and Simeone, F. A.: The electro-encephalogram in the evaluation of the effects of anesthetic agents and carbon dioxide accumulation during surgery, *Ann. Surg.* 138: 558, 1953.
  16. Brazier, M. A. B.: The physiological effects of carbon dioxide on the activity of the central nervous system in man: with special reference to the problem of high altitude flying, *Medicine* 22: 205, 1943.

---

### Nervous System

**MONAMINE OXIDASE INHIBITORS** The duration of action of norepinephrine released from cytoplasmic storage granules of neurones upon neuronal stimulation is largely limited by uptake again into nerve cytoplasm and storage granules and to a much lesser extent by diffusion away from the nerve ending and enzymatic destruction by catechol-O-methyl transferase. The chief function of monamine oxidase present in the mitochondria of adrenergic nerve endings may be to metabolize cytoplasmic norepinephrine that leaks from storage granules and escapes re-uptake. Monamine oxidase inhibitors cause an increase in neuronal stores of norepinephrine, but the mechanisms for disposing of norepinephrine released as a neurotransmitter or injected into the circulation are not disturbed other than by the limitation on tissue uptake imposed by the greater tissue stores. In experiments with human volunteers taking monamine oxidase inhibitors it was found, as expected, that injected norepinephrine is not potentiated. Ephedrine acts by releasing stored norepinephrine and since these stores are increased by monamine oxidase inhibitors the action of ephedrine is substantially potentiated. Phenylephrine acts chiefly directly on receptors and only to a small extent by release of stored norepinephrine and little potentiation occurs after intravenous administration, as expected. Substantial amounts of monamine oxidase are found in the intestine and liver and since phenylephrine acts as a substrate for monamine oxidase, administration of an inhibitor must be expected to increase absorption of phenylephrine. In 3 subjects, amounts of phenylephrine which barely caused blood pressure rise when given orally before monamine oxidase inhibitors, caused such dramatic blood pressure rise after inhibitors that the administration of the alpha blocking agent, phentolamine, was necessary. The inhibitors used in this study were phelzine and tranlycypromine. (*Elis, J., and others: Modification by Monoamine Oxidase Inhibitors of the Effect of Some Sympathomimetics of Blood Pressure, Brit. Med. J. I: 75 (April) 1967.*)