

depolarization block in 10 of 24 subjects. Hexafluorenum and *d*-tubocurarine always increased the neuromuscular effects of one another. The interaction of hexafluorenum and gallamine was not consistent. The hexafluorenum-induced neuromuscular block was increased by the subsequent administration of gallamine. When hexafluorenum was administered after gallamine the intensity of the block was increased in 5, unchanged in 2 and antagonized in 10 of the 17 subjects tested. There was a consistent mutual antagonism between the neuromuscular effects of hexafluorenum and decamethonium. The prior administration of hexafluorenum markedly potentiated and prolonged the neuromuscular blocking action of succinylcholine.

### References

1. Folds, F. F., Molloy, R. E., Zsigmond, E. K., and Zwart, J. A.: Hexafluorenum: Its anticholinesterase and neuromuscular activity, *J. Pharmacol. Exp. Ther.* 129: 400, 1960.
2. Macri, F. J.: Curare-like activity of some bisfluorenyl-bis-quaternary ammonium compounds, *Proc. Soc. Exp. Biol. Med.* 85: 603, 1954.
3. Folds, F. F., Hillmer, N. R., Molloy, R. E., and Monte, A. P.: Potentiation of the neuromuscular effect of succinylcholine by hexafluorenum, *ANESTHESIOLOGY* 21: 50, 1960.
4. Folds, F. F., Monte, A. P., Brunn, H. M., Jr., and Wolfson, B.: Studies with muscle relaxants in unanesthetized subjects, *ANESTHESIOLOGY* 22: 230, 1961.
5. Duncalf, D., Folds, F. F., Jacobson, E., and Vargas, P. G.: Investigation of the cardiovascular effects of hexafluorenum, *Brit. J. Anaesth.* 37: 591, 1965.
6. Rizzi, R.: Su una nuova associazione curarica, *Acta anaesth.* 7: 59, 1956.

7. Arrowood, J. G., and Kaplan, M. S.: Mylaxen: Intensification of neuromuscular blocking effect by combination with succinylcholine, *Anesth. Analg.* 35: 412, 1956.
8. Nastuck, W. L., and Karis, J. H.: The blocking action of hexafluorenum on neuromuscular transmission and its interaction with succinylcholine, *J. Pharmacol. Exp. Ther.* 144: 236, 1964.
9. Katz, R. L., Gissen, A. J., and Karis, J. H.: The effects of hexafluorenum and edrophonium on the neuromuscular blocking actions of succinylcholine, decamethonium, imbretil and *d*-tubocurarine, *ANESTHESIOLOGY* 26: 154, 1965.
10. Churchill-Davidson, H. C., and Christie, T. H.: The diagnosis of neuromuscular block in man, *Brit. J. Anaesth.* 31: 290, 1959.
11. Folds, F. F., Monte, A. P., Brunn, H. M., Jr., and Wolfson, B.: The influence of exercise on the neuromuscular activity of relaxant drugs, *Canad. Anaesth. Soc. J.* 8: 118, 1961.
12. Jenden, D. J., Kamijo, K., and Taylor, D. B.: The action of decamethonium (C10) on the isolated rabbit lumbrical muscle, *J. Pharmacol. Ther.* 111: 229, 1954.
13. Folds, F. F., Wnuck, A. L., Hodges, R. J. H., Thesleff, S., and deBeer, E. J.: The mode of action of depolarizing relaxants, *Anesth. Analg.* 36: 23, 1957.
14. Katz, R. L.: Neuromuscular effects of *d*-tubocurarine, edrophonium and neostigmine in man, *ANESTHESIOLOGY* 28: 327, 1967.
15. Waser, P. G., and Luthi, U.: Autoradiographische Lokalisation von <sup>14</sup>C-calebasencurarin. I. und <sup>14</sup>C-Decamethonium in der Motorischen Endplatte, *Arch. Int. Pharmacodyn.* 112: 273, 1957.
16. Thesleff, S.: Investigation of muscle-relaxing action of succinylcholine-iodide in man, *Acta physiol. Scand.* 25: 348, 1952.
17. McCaul, K., and Robinson, C. D.: Suxamethonium "extension" by tetrahydroaminacrine, *Brit. J. Anaesth.* 34: 536, 1962.

### Nervous System

**NORADRENALINE RELEASE** The view has been held for many years that the impulse which passes down the sympathetic postganglionic fiber releases noradrenaline directly. Current evidence indicates that the release is a more complicated process and that it closely resembles the mechanism of release of adrenaline from the adrenal medulla. The impulse passing down the fiber first releases acetylcholine, which, after leaving the fibers, makes the membrane of the fiber permeable to calcium ions. These ions enter the fiber and release noradrenaline from the granules in which they are held. (Burn, J. H.: *Release of Noradrenaline from the Sympathetic Postganglionic Fibre*, *Brit. Med. J.* 1: 197 (April) 1967.)