

3. Greenfield, A. D. M.: The circulation through the skin. In Hamilton, W. F., and Dow, P., eds.: *Handbook of Physiology—Circulation*. Baltimore, Williams and Wilkins Co., 1963.
4. Severinghaus, J. W.: The rate of uptake of nitrous oxide in man, *J. Clin. Invest.* 33: 1183, 1954.
5. Engstrom, C. G., Herzog, P., and Norlander, O.: A method for continuous measurement of oxygen consumption in the presence of inert gases during controlled ventilation, *Acta Anesth. Scand.* 5: 115, 1961.
6. Smith, T. C.: Nitrous oxide and low inflow circle systems, *ANESTHESIOLOGY* 27: 266, 1966.
7. Sechzer, P. H.: Cyclopropane. In Papper, E. M., and Kitz, R. J. (eds.): *Uptake and Distribution of Anesthetic Agents*. New York, 1962.
8. Eger, E. I., II, Johnson, E. A., Larson, C. P., and Severinghaus, J. W.: The uptake and distribution of intravenous ether, *ANESTHESIOLOGY* 23: 647, 1962.
9. Eger, E. I., II, and Guadagni, N. P.: Halothane uptake in man at a constant alveolar concentration, *ANESTHESIOLOGY* 24: 299, 1963.
10. Eger, E. I., and Larson, C. P., Jr.: Anesthetic solubility in blood and tissues: Values and significance, *Brit. J. Anaesth.* 36: 140, 1963.
11. Perl, W., Rackow, H., Salanitro, E., Wolf, G. L., and Epstein, R. M.: Intertissue diffusion effect for inert fat-soluble gases, *J. Appl. Physiol.* 20: 621, 1965.
12. Kety, S. K.: Exchange of inert gases at lung and tissues, *Pharmacol. Rev.* 3: 1, 1951.
13. Klocke, R. A., Gurtner, G. H., and Farhe, E.: Gas transfer across the skin in man, *Appl. Physiol.* 18: 311, 1963.

---

### Drugs

**NARCOTIC ANTAGONIST** A study was performed in volunteers to determine the relative degrees of respiratory depression produced by nalorphine and morphine and their interaction when administered together. The relative potency of nalorphine when compared with morphine was found to be 0.74; that is, 13.5 mg nalorphine produced depression in respiration approximately equivalent to 10 mg morphine. The data also suggest that antagonism of the respiratory effects of morphine by nalorphine is related primarily to differences in intrinsic activity of the drugs and not solely to receptor affinity. Their interaction could thus be classified as competitive dualism. (Bellville, J., and Fleischli, G.: *The Interaction of Morphine and Nalorphine on Respiration, Clin. Pharmacol. Therap.* 9: 152 (March) 1968.)

**RESERPINE** Hypertensive patients were studied before and after 30 days of treatment with reserpine. Cardiac output, measured both at rest and after induced atrial tachycardia, was found to be reduced approximately 25 per cent after reserpine treatment, while resting pulse was little affected. Atrioventricular conduction time increased significantly after reserpine. These results indicate that reserpine may lower cardiac output at rest significantly with no further decrease during atrial tachycardia. Reserpine may enhance second-degree heart block during induced atrial tachycardia. (Cohen, S. I., and others: *Effects of Reserpine Therapy on Cardiac Output and Atrioventricular Conduction During Rest and Controlled Heart Rates in Patients with Essential Hypertension, Circulation* 37: 738 (May) 1968.)