

results suggest that Forane neither depresses the human heart nor predisposes to arrhythmias. In miniature swine the metabolism of Forane is less than that of halothane or methoxyflurane. Muscle relaxation is good, and compared with halothane, Forane markedly potentiates the effect of *d*-tubocurarine. On the other hand, Forane is a profound respiratory depressant.

Does Forane possess other defects? Does it produce hepatic or renal toxic effects in man? Although 500 human exposures have failed to reveal any such toxicity, thousands of exposures may be necessary to uncover a "sensitization" phenomenon or a subtle toxic effect. Is Forane metabolized in man? What are its uptake, distribution and excretion characteristics? How soluble is it in body tissues? Does it enhance uterine bleeding? Does it have adverse effects on the fetus? Are there unfavorable or favorable interactions with other drugs patients might receive concomitantly? The list of unanswered questions is long. If initial results are confirmed by further studies, Forane may constitute a major advance in the search for the perfect anesthetic.

JAMES F. VITCHA
Manager, Chemical Product
Development
Ohio Medical Products, A Division
of Air Reduction Company, Inc.
Central Research Laboratories
Murray Hill, New Jersey

References

1. Lucas GHW, Henderson VE: New anesthetic gas cyclopropane. *Canad Med Ass J* 21: 173, 1929
2. Stiles JA, Neff WB, Rovenstine EA, et al.: Cyclopropane as an anesthetic agent. *Anesth Analg* 13:56, 1934
3. Davis HE, Krantz JC Jr: Studies of isopropenyl vinyl ether in man (propethylene ether). *ANESTHESIOLOGY* 5:159-162, 1944
4. Fisher AJ, Whitacre RJ: Clinical use of metopryl as an anesthetic agent. *ANESTHESIOLOGY* 8:156-158, 1947
5. Sadove MS, Wyant GM, Cletcher JO Jr: Ethyl vinyl ether: Pharmacological and clinical evaluation. *Anesth Analg* 34:235-240, 1955
6. Downing RC: History of organic fluorine industry, Kirk-Othmer Encyclopedia of Chemical Technology. Second edition. New York, Interscience, 1966, vol. 9, pp 704-707
7. Raventos J: Action of fluothane: A new volatile anesthetic. *Brit J Pharmacol* 11:394, 1956
8. Fabian LW, Carnes MA, DeWitt H: Laboratory and clinical investigation of some newly synthesized fluorocarbon anesthetics. *Anesth Analg* 39:456, 1960
9. Artusio JF Jr: Clinical investigation of telflurane, 2-bromo-1,1,1,2-tetrafluoroethane. *DA* 708, *Fed Proc* 22:186, 1963
10. Sulman M, Sadove MS: 1,1,1,2-Tetrafluoroethane: An inhalation anesthetic agent of intermediate potency. *Anesth Analg* 46: 629-633, 1967
11. Artusio, JF Jr, Van Poznak A, Hunt RE, et al.: Clinical evaluation of methoxyflurane in man. *ANESTHESIOLOGY* 21:512, 1960

Drugs

HALOTHANE HEPATITIS Halothane-induced stimulation of lymphocytes was observed in ten of 15 patients for whom diagnoses of drug-induced hepatitis were made following halothane (nine patients) or methoxyflurane (one patient) anesthesia. Stimulation of lymphocytes was measured serologically by the incorporation of ³H-thymidine into the DNA lymphocytes. Three patients exposed to halothane who did not develop jaundice, nine patients with hepatic disease and six healthy persons served as controls. None of the control patients showed any evidence of stimulation of lymphocytes in the presence of halothane. These data support the hypothesis that sensitization to halothane may be involved in the pathogenesis of hepatic damage following halothane and methoxyflurane. (Paronetto, F., and Popper, H.: *Lymphocyte Stimulation Induced by Halothane in Patients with Hepatitis Following Exposure to Halothane*, *New Engl. J. Med.* 283: 277 (Aug.) 1970.)