

- anesthetics. ANESTHESIOLOGY 28:432-437, 1967
6. Reynolds F, Taylor G: Maternal and neonatal blood concentrations of bupivacaine. A comparison with lignocaine during continuous extradural analgesia. Anaesthesia 25:14-23, 1970
 7. Tucker GT, Boyes RN, Bridenbaugh PO, et al: Binding of amilide-type local anesthetic in human plasma: II. Implications *in vivo*, with special reference to transplacental distribution. ANESTHESIOLOGY 33:304-314, 1970
 8. Finster M, Morishima HO, Boyes RN, et al: The placental transfer of lidocaine and its uptake by fetal tissues. ANESTHESIOLOGY 36:159-163, 1972
 9. Tucker GT, Morishima HO, Finster M, et al: Binding of amilide-type local anesthetics in human plasma. I. Relationships between binding, physicochemical properties and anesthetic activity. ANESTHESIOLOGY 33: 287-303, 1970
 10. Lund PC, Cwik JC, Cannon RT: Etidocaine (Duranest): A clinical and laboratory evaluation. Acta Anaesthesiol Scand 18:176-188, 1974
 11. Bridenbaugh PO, Tucker GT, Moore DC, et al: Preliminary clinical evaluation of etidocaine (Duranest): A new long-acting local anesthetic agent. Acta Anaesthesiol Scand 18: 165-171, 1974
 12. Editorial: Obstetric analgesia and the newborn baby. Lancet I:1090, 1974
 13. Desmond MM: Obstetric medication and infant behavior. ANESTHESIOLOGY 40:111-113, 1974
 14. Dubowitz V: Neurological fragility in the newborn: Influence of medication in labour. Br J Anaesth 47:1005-1010, 1975
 15. Tronick E, Wise S, Als H, et al: Regional obstetric anesthesia and newborn behavior: Effect over the first ten days of life. Pediatrics 58:94-100, 1976
 16. Levinson G, Shnider SM: Placental transfer of local anesthetics: Clinical implications. Parturition and Perinatology. Clinical Anesthesia. Philadelphia, F.A. Davis, 1973, pp 173-185

Thoracic Anesthesia

FIBEROPTIC BRONCHOSCOPY The fiberoptic bronchoscope is especially valuable for evaluation of peripheral lung densities, in elucidating hemorrhage of obscure cause, and in examination of patients on respirators or those who have cervical or cranial disease. The rigid bronchoscope retains certain advantages: it is preferred for removal of a foreign body or broncholith, for pediatric bronchoscopy, for aspiration of massive hemorrhage or retained thick secretions, and

for use in patients with narrowed tracheal diameters. The fiberoptic scope permits visualization of the larynx and nasopharynx. (Krook CJ: *Fiberoptic bronchoscopy*. *Carle Selected Papers* 28: 26, 1975, Urbana, Ill.)
ABSTRACTER'S COMMENT: The fiberoptic bronchoscope can serve as a guide over which an endotracheal tube can be passed in a patient whose anatomy prevents visualization of the larynx with a conventional laryngoscope.