

Effects of Propranolol on the Cardiovascular and Renin-Angiotensin Systems during Hypotension Produced by Sodium Nitroprusside in Humans

Wayne K. Marshall, M.D.,* Robert F. Bedford, M.D.,† William P. Arnold, M.D.,‡ Edward D. Miller, M.D.,§ David E. Longnecker, M.D.,¶ Michael D. Sussman, M.D.,** Michael W. Hakala, M.D.††

The authors examined the effects of controlled hypotension induced with sodium nitroprusside (SNP) with and without propranolol on the cardiovascular, pulmonary, and renin-angiotensin systems in 10 consecutive anesthetized patients with kyphoscoliosis undergoing posterior spinal fusion. SNP infusion ($4.1 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) alone decreased mean systemic arterial pressure (SAP) by $25 \text{ torr} \pm 3 \text{ SE}$ ($P < 0.001$), systemic vascular resistance index (SVRI) by $1113 \text{ dyne} \cdot \text{sec} \cdot \text{cm}^{-5} \cdot \text{m}^2 \pm 125 \text{ SE}$ ($P < 0.001$), mean pulmonary artery pressure (PAP) by $6 \text{ torr} \pm 2 \text{ SE}$ ($P < 0.02$), pulmonary capillary wedge pressure (PCWP) by $4 \text{ torr} \pm 1 \text{ SE}$ ($P < 0.01$), pulmonary vascular resistance (PVR) by $50 \text{ dyne} \cdot \text{sec} \cdot \text{cm}^{-5} \pm 18$ ($P < 0.05$), and Pa_{O_2} by $16 \text{ torr} \pm 7 \text{ SE}$ ($P < 0.05$), whereas cardiac index increased by $1.08 \text{ l} \cdot \text{min}^{-1} \cdot \text{m}^2 \pm 0.24 \text{ SE}$ ($P < 0.01$) and heart rate increased $16 \text{ beats/min} \pm 5 \text{ SE}$ ($P < 0.02$). After 40 min of hypotension, 0.03 mg/kg propranolol was injected intravenously while the SNP infusion rate was held constant. Ten min later there was a significant decrease in the heart rate ($10 \text{ beats/min} \pm 4 \text{ SE}$, $P < 0.02$) and cardiac index ($0.65 \text{ l} \cdot \text{min}^{-1} \cdot \text{m}^2 \pm 0.21$, $P < 0.02$). Plasma renin activity (PRA) increased from $2.37 \text{ ng} \cdot \text{ml}^{-1} \cdot \text{h}^{-1} \pm 0.7 \text{ SE}$ before anesthesia to $6.50 \text{ ng} \cdot \text{ml}^{-1} \cdot \text{h}^{-1} \pm 1.45 \text{ SE}$ ($P < 0.05$) after 40 min of nitroprusside infusion. Forty min after propranolol there was a significant reduction in PRA to $4.07 \text{ ng} \cdot \text{ml}^{-1} \cdot \text{h}^{-1} \pm 0.73 \text{ SE}$ ($P < 0.05$). Thus propranolol, when given during SNP hypotension, exhibits an early cardiovascular response manifested as a decrease in cardiac output and heart rate and a delayed action on the kidney resulting in an inhibition of renin release. (Key words: Anesthetic technique: hypotension, induced, nitroprusside. Polypeptides: renin-angiotensin. Surgery: orthopedic. Sympathetic nervous system: sympatholytic agents, propranolol.)

ALTHOUGH the cardiovascular and pulmonary consequences of kyphoscoliosis in awake unanesthetized individuals have been well-documented,¹⁻⁹ data pertaining to the hemodynamic and respiratory effects of

anesthesia and/or induced hypotension in these patients are limited. Controlled hypotension with sodium nitroprusside (SNP) has been shown to be effective in reducing blood loss associated with mandibular surgery¹⁰ and total hip arthroplasty.¹¹ SNP dose requirement can be reduced by giving propranolol in a dose large enough to slow heart rate during controlled hypotension,¹² thus decreasing the likelihood of developing cyanide toxicity. The precise mechanism of action of propranolol in this circumstance has not yet been defined, although a possible effect on the renin-angiotensin system has been suggested.¹³

The present study examined the following areas in patients with kyphoscoliosis: 1) the effects of anesthesia and hypotension induced with SNP on the pulmonary, cardiovascular, and renin-angiotensin systems; and 2) the effects of propranolol on the pulmonary, cardiovascular, and renin-angiotensin systems during hypotension induced with SNP.

Methods

The subjects were ten consecutive patients with moderately severe kyphoscoliosis, ages 13-36 yr, undergoing posterior spinal fusion and Harrington rod insertion. The study protocol was reviewed by the Institutional Human Studies Committee and detailed informed consent was obtained from the subjects and/or their legal guardians on the day prior to operation.

Premedication consisted of morphine (0.1 mg/kg) and scopolamine (0.005 mg/kg), intramuscularly (IM), one hour prior to induction of anesthesia. After the patient arrived in the operating suite, a peripheral intravenous infusion was begun and a venous blood sample was obtained for plasma renin activity (PRA) measurement. Gallamine, $10-20 \text{ mg}$, was given intravenously (iv), and anesthesia was induced with sodium thiopental, $3-4 \text{ mg/kg}$, iv, and maintained with 70 per cent nitrous oxide in oxygen and morphine, 0.5 mg/kg , iv. Endotracheal intubation was facilitated with succinylcholine, 1 mg/kg , iv, after topical laryngotracheal anesthesia with 4 per cent lidocaine 160 mg . Curare, 0.3 mg/kg , iv, was then

* Research Fellow, Department of Anesthesiology.

† Assistant Professor of Anesthesiology and Neurological Surgery.

‡ Assistant Professor of Anesthesiology.

§ Associate Professor of Anesthesiology.

¶ Professor of Anesthesiology.

** Assistant Professor of Orthopedic Surgery and Pediatrics.

†† Associate Professor of Orthopedic Surgery and Pediatrics.

Received from the Departments of Anesthesiology and Orthopedic Surgery, University of Virginia Medical Center, Charlottesville, Virginia 22908. Accepted for publication February 6, 1981.

Address reprint requests to Dr. Marshall: Department of Anesthesiology, Pennsylvania State University College of Medicine, Milton S. Hershey Medical Center, 500 University Drive, Hershey, Pennsylvania 17033.

TABLE 1. Calculations and Formulae

$\overline{SAP} = \left(\frac{\text{Systolic} - \text{Diastolic}}{3} \right) + \text{Diastolic}$	(torr)
$\overline{PAP} = \left(\frac{\text{Systolic} - \text{Diastolic}}{3} \right) + \text{Diastolic}$	(torr)
$CI = \frac{CO}{BSA (m^2)}$	(l·min ⁻¹ ·m ⁻²)
$SVRI = \frac{(\overline{SAP} - CVP) \times 79.92}{CI}$	(dyne·sec·cm ⁻⁵ ·m ²)
$PVR = \frac{(\overline{PAP} - PCWP) \times 79.92}{CO}$	(dyne·sec·cm ⁻⁵)
$SVI = \frac{CI}{HR}$	(ml·m ²)

given and constant controlled mechanical ventilation was instituted to maintain the arterial CO₂ tension (P_{aCO₂}) near 35 torr throughout the operation.

After induction of anesthesia, a 20-gauge radial arterial catheter and a 7-French thermistor-tipped triple-lumen pulmonary artery catheter were introduced percutaneously. The latter was localized using pressure waveform control. The legs were wrapped with elastic bandages and patients were placed prone on a chest frame for operation.

Radial arterial, pulmonary arterial, and central venous pressures were continuously monitored (Bentley® Model 500 transducers) and recorded throughout the study period. The following variables were determined: heart rate (HR); mean systemic arterial pressure (SAP); mean pulmonary artery pres-

sure (\overline{PAP}); pulmonary capillary wedge pressure (PCWP); central venous pressure (CVP); and body temperature. Cardiac output (CO) was measured by thermal dilution technique in triplicate (Edwards® 9520-A computer). Body surface area (BSA) was calculated from standard nomograms using height and weight, and the following hemodynamic variables were calculated from appropriate formulae (table 1): cardiac index (CI); systemic vascular resistance index (SVRI); pulmonary vascular resistance (PVR); and stroke volume index (SVI). Arterial and mixed venous blood samples were obtained anaerobically for analysis of pH, oxygen, and CO₂ tensions (Corning® model 175 pH/blood-gas analyzer). Mixed venous blood samples were obtained for analysis of plasma renin activity (PRA; measured using radioimmunoassay, method of Haber; New England Nuclear Kit); and whole blood cyanide (by the method of Rodkey^{14,15}).

Hemodynamic measurements and blood samples were obtained: 1) prior to induction of anesthesia (blood samples only); 2) after anesthetic induction and placement in the prone position; 3) 10 min after incision (and infiltration with 1:500,000 epinephrine-saline mixture 25–50 ml); 4) after 10 min of constant SNP infusion sufficient to maintain \overline{SAP} at 71 torr ± 3 (SE); and 5) after 40 min of SNP while the SNP infusion rate was held constant (4.1 μg·kg⁻¹·min⁻¹ ± 0.53 SE). Propranolol, 0.03 mg/kg, iv, was then administered as a bolus. The SNP infusion rate continued at 4.1 μg·kg⁻¹·min⁻¹ ± 0.53 SE and data were again collected: 10 min and 40 min after propranolol. The study was then terminated. All PRA samples were drawn from the pulmonary artery ex-

TABLE 2. Hemodynamic and Laboratory Values before and during Nitroprusside Infusion

	Awake	Prone Anesthetized	Incision + 10 min	SNP + 10 min	SNP + 40 min	Propranolol + 10 min	Propranolol + 40 min
HR (beats/min)		64 ± 4	64 ± 6	78 ± 7	80 ± 7†	70 ± 6‡	69 ± 5‡
SAP (torr)		79 ± 2	99 ± 3*	71 ± 3†	74 ± 3*	76 ± 2	74 ± 3
\overline{PAP} (torr)		18 ± 2	22 ± 2*	14 ± 2†	16 ± 2†	17 ± 2	18 ± 2
PCWP (torr)		14 ± 2	16 ± 2*	8 ± 2†	12 ± 2†	13 ± 2	13 ± 2
CVP (torr)		9 ± 1	10 ± 1	6 ± 1†	8 ± 1*	9 ± 2	9 ± 1‡
CI (l·min ⁻¹ ·m ⁻²)		2.71 ± 0.09	2.91 ± 0.15	3.58 ± 0.41	3.99 ± 0.34†	3.34 ± 0.27‡	3.43 ± 0.20
PVR (dyne·sec·cm ⁻⁵)		91 ± 18	103 ± 18	74 ± 10	54 ± 6†	71 ± 9‡	71 ± 9‡
SVRI (dyne·sec·cm ⁻⁵ ·m ²)		2075 ± 87	2502 ± 163	1625 ± 145†	1389 ± 114†	1695 ± 137‡	1574 ± 129
SVI (ml/m ²)		43.47 ± 2.5	47.75 ± 3.9	48.92 ± 5.4	52.56 ± 5.6	49.90 ± 4.5	51.05 ± 3.7
P _{aO₂} (torr)		139 ± 12	126 ± 9	120 ± 7†	110 ± 7†	120 ± 6	123 ± 7
P _{aCO₂} (torr)		34 ± 2	34 ± 2	34 ± 2	35 ± 1	35 ± 1	35 ± 2
pH _a		7.46 ± 0.02	7.46 ± 0.02	7.43 ± 0.02	7.43 ± 0.02	7.44 ± 0.01	7.45 ± 0.02
P \overline{v} O ₂ (torr)		40 ± 1	38 ± 1	42 ± 2†	41 ± 1	39 ± 1	36 ± 1‡
Hct		36 ± 1	36 ± 1	35 ± 1	34 ± 1	34 ± 1	34 ± 1
PRA (ng·ml ⁻¹ ·h ⁻¹)	2.37 ± 0.7	5.35 ± 1.43*	5.42 ± 1.29	6.66 ± 1.69	6.50 ± 1.45	5.47 ± 1.01	4.07 ± 0.73‡
CN ⁻ (μmol/l)		1.02 ± 0.26			5.64 ± 0.85*		13.19 ± 2.26‡
Temperature (°C)		36.1 ± 0.1	35.6 ± 0.2	35.3 ± 0.2	35.1 ± 0.2	35.1 ± 0.3	35.0 ± 0.3

All values are means ± SE. * Indicates $P < 0.05$ vs. preceding value; † Indicates $P < 0.05$ vs. values 10 minutes after incision;

‡ Indicates $P < 0.05$ vs. values after 40 minutes of SNP (just before propranolol).

cept for those obtained before anesthetic induction which were from a peripheral vein.

All data were analyzed using Student's *t* test for paired data. *P* < 0.05 was regarded as significant.

Results

Incision resulted in significant increases in \overline{SAP} , \overline{PAP} , and PCWP as compared with control values (table 2). SNP infusion produced significant decreases in \overline{SAP} , SVRI, \overline{PAP} , PCWP, and PVR, accompanied by significant increases in HR and CI and a progressive reduction in Pa_{O_2} despite a constant inspired oxygen fraction of 0.3.

After propranolol administration there were significant decreases in HR and CI whereas SVRI, PVR, and CVP increased. The progressive reduction in arterial oxygen tension (Pa_{O_2}) observed during SNP infusion did not continue following propranolol.

PRA was $2.37 \text{ ng} \cdot \text{ml}^{-1} \cdot \text{h}^{-1} \pm 0.78 \text{ SE}$ prior to induction of anesthesia and increased significantly after induction of anesthesia, intubation, cannulation, and turning into the prone position. No further increase in PRA was observed after incision and SNP infusion. Propranolol administration significantly decreased PRA as compared with values measured just before propranolol was given. Control PRA values varied markedly between individuals, a finding also observed by others.¹⁶

Whole blood cyanide increased progressively during SNP infusion with a peak value of $13.19 \mu\text{mol/l} \pm 2.26 \text{ SE}$ being reached 40 min after propranolol was administered.

Discussion

The basic derangement in kyphoscoliosis is compression of lung tissue from chest wall distortion, with consequent loss of thoracic volume leading to ventilation-perfusion mis-match, hypoxia, and eventual pulmonary hypertension.⁵ Our data after anesthesia and intubation agree well with the observations of Bergofsky *et al.*² in five unanesthetized patients with moderately severe kyphoscoliosis who were clinically comparable to our patients. They observed a \overline{PAP} of 16 torr ± 2 (SE), a CI of $3.2 \text{ l} \cdot \text{min}^{-1} \cdot \text{m}^{-2} \pm 0.1$, and a normal PCWP.³

Infusion of the vasodilator SNP caused reductions in SVRI and PVR with resultant reductions in \overline{SAP} , \overline{PAP} , and PCWP. The concomitant increases in CI and HR are thought to be reflex responses to the decrease in \overline{SAP} ^{17,18} and may be baroreceptor-mediated.¹⁹ The hemodynamic response to propranolol was characterized by decreases in HR and CI within 10 min of the administration of the drug. These observa-

tions are in agreement with previously published data demonstrating that intravenous propranolol decreases HR and CI within minutes.²⁰

The significant increase in PRA after anesthesia and the absence of a further increase after epinephrine infiltration, surgical incision, and SNP infusion apparently reflect a near-maximal renin response to induction of anesthesia, endotracheal intubation, vascular cannulation, and turning prone.^{21,22} Propranolol significantly decreased PRA as compared with levels observed just before propranolol was given, but this was not seen until 40 min after the drug was administered. Davies and Slater also found a 15–30 min delay after propranolol before observing decreased PRA response to head-tilt maneuvers in healthy subjects.²³ In the rat PRA supports blood pressure during SNP-induced hypotension and this effect can be blocked by propranolol.²⁴ In addition, propranolol is known to block the renal release of renin in response to both generalized sympathetic stimulation²⁵ and direct stimulation of the renal nerves.²⁶ Thus, our data appear to indicate two distinct effects of propranolol during SNP hypotension: 1) an early, direct effect on the heart characterized by decreases in HR and CI; and 2) an effect on the kidney manifested by a decrease in PRA occurring later.

Arterial oxygenation decreased progressively during SNP infusion, despite a constant inspired oxygen fraction of 0.30. This may have been related to release of hypoxic pulmonary vasoconstriction by SNP, a significant phenomenon known to occur in dogs,^{27,28} but as yet unproven in humans.²⁹ PVR increased after propranolol, and it is possible that propranolol reversed the effect of SNP on hypoxic pulmonary vasoconstriction by restoring pulmonary vascular tone. This is highly speculative since propranolol has no apparent effect on arterial vasodilation caused by nitroglycerin, histamine, or acetylcholine.³⁰ In addition, the pulmonary vasculature is unresponsive to angiotensin-II³¹ which should have decreased along with the decrease in PRA values after propranolol. An alternative explanation for the effect of propranolol on oxygenation is that it reversed the progressive increase in cardiac output caused by SNP infusion. As recently pointed out by Cheney and Colley, the effect of cardiac output on oxygenation may be variable, but in regional atelectasis, such as is seen with kyphoscoliosis, shunt fraction has been found most frequently to vary inversely with cardiac output.³²

The continuous increase in CN^- levels during the study period was due to the constant rate of infusion of SNP. There was, however, no change in arterial *pH* or Pv_{O_2} suggestive of cyanide toxicity in any of the patients.

In summary, propranolol administration during SNP infusion decreases HR, CI, and PRA. These may be the mechanisms by which previously documented decreases in SNP dose requirement by propranolol¹² are achieved.

References

1. Kafer ER: Respiratory and cardiovascular functions in scoliosis and the principles of anesthetic management. *ANESTHESIOLOGY* 52:339-351, 1980
2. Bergofsky EG, Turino GM, Fishman AP: Cardiorespiratory failure in kyphoscoliosis. *Medicine* 38:263-317, 1959
3. Shneerson JM, Venco A, Prime FJ: A study of pulmonary artery pressure, electrocardiography, and mechanocardiography in thoracic scoliosis. *Thorax* 32:700-705, 1977
4. Shneerson JM: Pulmonary artery pressure in thoracic scoliosis during and after exercise while breathing air and pure oxygen. *Thorax* 33:747-754, 1978
5. Kafer ER: Idiopathic scoliosis: Gas exchange and the age dependence of arterial blood gases. *J Clin Invest* 58:825-833, 1976
6. Makley JT, Herndon CH, Inkley S, et al: Pulmonary function in paralytic and non-paralytic scoliosis before and after treatment. *J Bone Joint Surg (Am)* 50-A:1379-1380, 1968
7. Gazioglu K, Goldstein LA, Femi-Pearse D, et al: Pulmonary function in idiopathic scoliosis. *J Bone Joint Surg (Am)* 50-A: 1391-1399, 1968
8. Naeye RL: Kyphoscoliosis and Cor pulmonale. *Am J Pathol* 38: 561-573, 1961
9. Davies G, Reid L: Effect of scoliosis on growth of alveoli and pulmonary arteries and on right ventricle. *Arch Dis Child* 46:623-632, 1971
10. Sivarajan M, Amory DW, Everett GB, et al: Blood pressure, not cardiac output, determines blood loss during induced hypotension. *Anesth Analg (Cleve)* 59:203-206, 1980
11. Thompson GE, Miller RD, Stevens WC, et al: Hypotensive anesthesia for total hip arthroplasty: A study of blood loss and organ function (brain, heart, liver, and kidney). *ANESTHESIOLOGY* 48:91-96, 1978
12. Bedford RF, Berry FA, Longnecker DE: Impact of propranolol on hemodynamic responses and blood cyanide levels during nitroprusside infusion: A prospective study in anesthetized man. *Anesth Analg (Cleve)* 58:466-469, 1979
13. Khambatta HJ, Stone JG, Khan E: Propranolol abates nitroprusside-induced renin release. *ANESTHESIOLOGY* 51:S74, 1979
14. Rodkey FL, Collison HA: Determination of cyanide and nitroprusside in blood and plasma. *Clin Chem* 23:1969-1975, 1977
15. Rodkey FL, Robertson RF: Analytical precautions in measurement of blood cyanide. *Clin Chem* 24:2184-2185, 1978
16. Freeman RH, Davis JO: The control of renin secretion and metabolism. *Hypertension*. New York, McGraw-Hill, 1979, pp 210-240
17. Rowe GG, Henderson RH: Systemic and coronary hemodynamic effects of sodium nitroprusside. *Am Heart J* 87: 83-87, 1974
18. Tinker JH, Michenfelder JD: Sodium nitroprusside. *Pharmacology, toxicology, and therapeutics*. *ANESTHESIOLOGY* 45: 340-354, 1976
19. Guyton AC: *Textbook of Medical Physiology*, Philadelphia, W. B. Saunders, 1976, pp 268-269
20. Robin E, Cowan C, Puri P, et al: A comparative study of nitroglycerine and propranolol. *Circulation* 36:175-186, 1967
21. Jakubowski HD, Taube HD: Plasma renin activity in surgical patients. *Recent Progress in Anaesthesiology and Resuscitation*. Edited by Arias A, Llauro R, Nalda MA, et al. Amsterdam, Excerpta Medica (International Congress Series 347), 1975, pp 359-361
22. Bailey DR, Miller ED, Kaplan JA, et al: The renin-angiotensin-aldosterone system during cardiac surgery with morphine-nitrous oxide anesthesia. *ANESTHESIOLOGY* 42:538-544, 1975
23. Davies R, Slater JDH: Is the adrenergic control of renin release dominant in man? *Lancet* 2:594-596, 1976
24. Miller ED, Ackerly JA, Vaughan ED, et al: The renin-angiotensin system during controlled hypotension with sodium nitroprusside. *ANESTHESIOLOGY* 47:257-262, 1977
25. Zanchetti A, Stella A: Neural control of renin release. *Clin Sci Molec Med* 42(S):215-223, 1975
26. Imbs JL, Kraetz J, Schmidt M, et al: Beta-blocking drugs and renin secretions in the anesthetized dog. *Clin Sci Molec Med* 49(S):105-107, 1975
27. Colley PS, Cheney FW: Sodium nitroprusside increases Q_a/Q_t in dogs with regional atelectasis. *ANESTHESIOLOGY* 47: 338-341, 1977
28. Colley PS, Cheney FW, Hlastala MP: Ventilation-perfusion and gas exchange effects of sodium nitroprusside in dogs with normal and edematous lungs. *ANESTHESIOLOGY* 50: 489-495, 1979
29. Benumof JL: Hypoxic pulmonary vasoconstriction and infusion of sodium nitroprusside. *ANESTHESIOLOGY* 50:481-483, 1979
30. Goodman LS, Gillman A: *The Pharmacological Basis of Therapeutics*. 5th Edition. New York, MacMillan, 1975, pp 549
31. Page IH, Bumpass FM: Angiotensin. *Handbook of Experimental Pharmacology*, Vol 37. New York, Springer-Verlag, 1974
32. Cheney FW, Colley PS: The effect of cardiac output on arterial blood oxygenation. *ANESTHESIOLOGY* 52:496-503, 1980