

Title: EFFECTS OF ISOFLURANE ON FUNCTIONALLY SKINNED MYOCARDIAL FIBERS FROM RABBITS

Authors: J. Y. Su, Ph.D., and J. G. Bell, B.S.

Affiliation: Department of Anesthesiology, University of Washington, Seattle, Washington 98195

Introduction. Isoflurane(I)¹ like halothane (H) and enflurane(E),² at clinical concentrations depresses myocardial contractility in isolated intact cat papillary muscle. The mechanisms of the action of H^{3,4} and E⁵ have been shown mainly due to decreasing Ca²⁺ uptake by the sarcoplasmic reticulum(SR) in functionally skinned myocardial fibers from rabbits. The isomeric structure of I and E led us to hypothesize that I would have similar mechanisms of action as E.⁵ The present study was designed to test the hypothesis by studying the effects of I on functionally skinned myocardial fibers.

Methods. Right ventricular papillary muscles from rabbits were isolated. Pieces of the muscle were homogenized in relaxing solution(-log[Ca²⁺] = pCa > 9) to disrupt the sarcolemma. A fiber bundle was dissected from the homogenate and mounted between forceps, and attached to a photodiode tension transducer. **Study on the Contractile Proteins:** The fibers were activated with one of the submaximal [Ca²⁺](pCa 5.6-5.0) and the maximal [Ca²⁺](pCa 3.8) and were relaxed with a relaxing solution(pCa > 9) in between contractions. The steady-state tension development was measured. The submaximal Ca²⁺-activated tension development was expressed as percentage of the maximal Ca²⁺-activated tension as 100%. **Study on the SR:** The fibers were immersed sequentially in five different solutions to load Ca²⁺ into the SR and to release Ca²⁺ from the SR with caffeine producing a tension transient. The area of the tension transient was measured as an estimate of the amount of Ca²⁺ released. Three types of tests were performed: (i) uptake phase- the fibers immersed in loading solutions saturated with I, (ii) release phase- the fibers immersed in the releasing solution saturated with I, and (iii) uptake-and-release phase- the fibers immersed in (i) and (ii) solutions. Isoflurane was delivered through a Verni-Trol® vaporizing system. The partial pressure of I in the solutions was assayed by gas chromatograph. Each preparation was immersed in control solutions (saturated with pure N₂), followed by test solutions (saturated with I and N₂ mixture) and finally again the control solutions. The test results were compared with the mean of the two bracketing controls by paired t-test. The mean and the standard error of the mean were calculated from at least 6 preparations of at least 3 rabbits.

Results. Isoflurane (2,3,4%) slightly decreased(7-9%) the maximal Ca²⁺-activated tension. The submaximal Ca²⁺-activated tension was decreased by I(2-4%) so that higher[Ca²⁺] was required to generate the same amount of tension (Fig.1, left). Isoflurane(3,4%) decreased 16% and 30% respectively the caffeine-induced tension transient at the uptake phase, and did not change the tension transient at the release phase(Fig.1,

right). Isoflurane(4%) decreased 21% the tension transient at the uptake-and-release phase. However, I(4%) increased the submaximal caffeine (2 mM)-induced tension transient at the release phase(Fig.1, right).




Fig. 1. Effects of I(2,3,4%) on(left)[Ca²⁺]-tension relationship and on(right) the caffeine-induced tension transient at three experimental conditions(Mean ± SE(n)). * p < 0.05 compared with the control.

Discussion. The results prove that our hypothesis is correct: the decreased Ca²⁺ activation of the contractile proteins and decreased Ca²⁺ uptake by the SR by I are similar to that of H^{3,4} and E.⁵ At clinical concentrations of I(1-3%), the slight decrease in the Ca²⁺-activated tension development of the contractile proteins and Ca²⁺ uptake by the SR could only partly be responsible for the marked myocardial depression observed in isolated intact cat papillary muscle.¹ We conclude that I-induced myocardial depression is partly achieved by decreasing Ca²⁺ activation of the contractile proteins. Isoflurane has a much less depressive effect on the SR than H or E.

Supported by NIH, No. HL20754; isoflurane was supplied by Ohio Medical Anesthetics, Madison, WI.

References.

1. Kemmotsu O, Hashimoto Y, Shimosato S: Inotropic effects of isoflurane on mechanics of contraction in isolated cat papillary muscles from normal and failing hearts, *Anesthesiology* 39:470-477, 1973
2. Brown BR, Crout JR: A comparative study of the effects of five general anesthetics on myocardial contractility: I. Isometric conditions, *Anesthesiology* 34:236-245, 1971
3. Su JY, Kerrick WGL: Effects of halothane on Ca²⁺-activated tension development in mechanically disrupted rabbit myocardial fibers, *Pflügers Arch* 375:111-117, 1978
4. Su JY, Kerrick WGL: Effects of halothane on caffeine-induced tension transients in functionally skinned myocardial fibers, *Pflügers Arch* 380:29-34, 1979
5. Su JY, Kerrick WGL: Effects of enflurane on functionally skinned myocardial fibers from rabbits, *Anesthesiology* 95:753-764, 1980