

14. Hillyer KW, Johnston RR: Unsuspected dilution of anesthetic gases detected by an oxygen analyzer. *Anesth Analg (Cleve)* 57:491-492, 1978
15. Bageant RA, Hoyt JW, Epstein RM: Error in a pipeline gas concentration: An unanticipated consequence of a defective check valve. *ANESTHESIOLOGY* 54:166-169, 1981
16. Baraka A, Maullem M: Awareness during anesthesia due to a ventilator malfunction. *Anaesthesia* 34:678-679, 1979
17. Manawadu BR, Hartwig FE, Sherrill D, Swanson GD: Monitoring oxygen consumption utilizing low flow techniques, Low Flow and Closed System Anesthesia. Edited by Aldrete JA, Lowe HJ, Virtue RW. New York, Grune and Stratton, 1979, pp 147-150

(Accepted for publication January 21, 1983.)

Anesthesiology
59:162, 1983

Plasma-Sulfate-Conjugated Catecholamines during Anesthesia

To the Editor:—Recent reports in the Journal have examined the effects of anesthetics on adrenergic response during surgery.^{1,2} Responses were evaluated by measuring variations in concentrations of plasma norepinephrine and epinephrine. However, the methods used in these studies measured only free plasma catecholamines, whereas, in humans, 70–80% of total plasma catecholamines are sulfate conjugated.³ Thus, we performed a study in seven ASA class I or II patients (aged 25–69 years) to delineate variations in sulfate-conjugated catecholamines during anesthesia.

In all cases, induction was achieved with thiopental, 5 mg/kg, and succinylcholine, 1 mg/kg, followed by nitrous-oxide-halothane for maintenance. Radial artery blood samples were obtained simultaneously before induction of anesthesia (T1) (20 min after intraarterial insertion of catheter), 1 min after tracheal intubation (T2), and 1 min after skin incision (T3) for measurement of total and free norepinephrine (NE) and epinephrine (E) levels, using a radio-enzymatic assay method. Sulfate-conjugated catecholamines were calculated from these data by subtraction.³

As noted in other studies, free NE increased during nociceptive stimulations (table 1); free E levels decreased. Total NE and E levels did not change. The percentage of sulfate-conjugated NE decreased significantly, while the percentage of sulfate-conjugated E increased.

Sulfate-conjugated catecholamines are usually thought to be of minor importance in the regulation of circulation in humans except in some situations.⁴ The role they play in anesthetized patients currently is under study.

J. MARTY, M.D.
*Département d'Anesthésie
Hôpital Bichat*

J. COUTURE, M.D.
*Département d'Anesthésie
Hotel Dieu de Montréal*

J. DE CHAMPLAIN
*Département de Physiologie
Université de Montréal
Paris, France*

TABLE 1.

	T1	T2	T3
Total NE pg/ml	982 ± 174	1,031 ± 180	1,033 ± 190
Free NE pg/ml	185 ± 32	275 ± 48†	305* ± 74
Sulfate-conjugated NE in per cent	79.9 ± 3	71.2* ± 3.7	69.3* ± 2.3
Total E pg/ml	517 ± 188	441 ± 168	366 ± 171
Free E pg/ml	64 ± 15	29† ± 4	32† ± 6
Sulfate-conjugated E in per cent	82.4 ± 4.1	89.5† ± 2.1	89.6† ± 2.3

Values are mean ± SE.

* $P < 0.01$ versus T1 (analysis of variance).

† $P < 0.05$.

REFERENCES

1. Roizen MF, Horrigan RW, Frazer BM: Anesthetic doses blocking adrenergic (stress) and cardiovascular responses to incision MAC-BAR. *ANESTHESIOLOGY* 54:390-398, 1981
2. Pflug AE, Halter JB: Effects of spinal anesthesia on adrenergic tone and neuro-endocrine responses to surgical stress in man. *ANESTHESIOLOGY* 55:120-126, 1981
3. Johnson GA, Baker CA, Smith RT: Radioenzymatic assay of sulfate conjugates of catecholamines and dopa in plasma. *Life Sci* 26:1591-1598, 1980
4. Kuchel D, Burr NT, Hamet P, Larochelle P, Bourque M, Genest J: Essential hypertension with low conjugated catecholamines imitates pheochromocytoma. *Hypertension* 3:347-355, 1981

(Accepted for publication January 21, 1983.)