

patients receiving epidural morphine, patient satisfaction with the relief obtained was comparable. Factors responsible for the lack of correlation between apparent pain and increased pain tolerance and satisfaction may be related to a more uniform level of analgesia, a lack of troublesome side effects, independence, and an awareness that reliable analgesia will be provided quickly as required. Patients utilizing PCA morphine did not achieve total analgesia, but settled at a level in which comfort was balanced by troublesome side effects, *i.e.*, increasing sedation, nausea, and dizziness. It may well be that morphine is not the optimal narcotic for use in PCA. An agent with faster onset, less sedation, and fewer adverse effects might further increase patient satisfaction and increase levels of analgesia to levels achieved with epidural narcotics.

In summary, two newer modalities of providing postoperative pain relief offered either superior analgesia or higher patient satisfaction than traditional im dosing. Epidural morphine offers the best possible analgesia if one is willing to accept higher morbidity and risks of delayed respiratory depression. In settings where epidural narcotics cannot be administered, the high patient satisfaction and uniform analgesia offered by PCA provides an attractive alternative.

The authors wish to thank Nelsann Kadric, Leonda Fermo, and Peter Capello for their invaluable assistance during the study.

#### REFERENCES

1. Marks RM, Sachar EJ: Undertreatment of medical inpatients with narcotic analgesics. *Ann Intern Med* 78:173-181, 1973

Anesthesiology  
68:457-459, 1988

2. Kotelko DM, Dailey PA, Shnider SM, Rosen MA, Hughes SC, Brizgys RV: Epidural morphine analgesia after cesarean delivery. *Obstet Gynecol* 63:409-413, 1984
3. Graves DA, Foster TS, Batenhorst R, Bennett RL, Baumann TJ: Patient controlled analgesia. *Ann Intern Med* 99:360-365, 1983
4. Rosenberg PH, Heino A, Scheinin B: A comparison of intramuscular analgesia, intercostal block, epidural morphine and on demand IV fentanyl in the control of pain after upper abdominal surgery. *Acta Anesthesiol Scand* 28:603-607, 1984
5. Purves PG, Spelling SJ, Dykes V, Stanley TD: Improved postoperative analgesia: Epidural vs. patient-controlled analgesia (abstract). *Anesth Analg* 66:S143, 1987
6. Pybus DA, Torda TA: Dose-effect relationships of extradural morphine. *Br J Anaesth* 54:1259-1262, 1982
7. Nordberg G, Hedner T, Mellstrand T, Dahlstrom B: Pharmacokinetic aspects of epidural morphine analgesia. *ANESTHESIOLOGY* 58:545-551, 1983
8. Graves DA, Arrigo JM, Foster TS, Baumann TJ, Batenhorst RL: Relationship between plasma morphine concentrations and pharmacologic effects in postoperative patients using patient-controlled analgesia. *Clinical Pharm* 4:41-47, 1985
9. Cousins MJ, Mather LE: Intrathecal and epidural administration of opioids. *ANESTHESIOLOGY* 61:276-310, 1985
10. Martin R, Salbaing J, Blaise G, Tetrault JP, Tetrault L: Epidural morphine for postoperative pain relief: A dose-response curve. *ANESTHESIOLOGY* 56:423-426, 1982
11. Matthew EB, Henderson SK, Avram MJ, Glassenberg R, Cohen H: Epidural hydromorphone versus epidural morphine for postcesarean section analgesia. *Anesth Analg* 66:S112, 1987
12. Leicht CH, Rosen MA, Dailey PA, Hughes SC, Shnider SM, Baker BW, Cheek DB, O'Connor DE: Evaluation and comparison of epidural sufentanil citrate and morphine sulfate for analgesia after cesarean section. *ANESTHESIOLOGY* 65:A365, 1986
13. Leicht CH, Hughes SC, Dailey PA, Shnider SM, Rosen MA: Epidural morphine sulfate for analgesia after cesarean section: A prospective report of 1000 patients. *ANESTHESIOLOGY* 65:A366, 1986

## Venous Air Embolism during Removal of Tissue Expander in a Supine Child

ANDREW G. ROTH, M.D.,\* DIANE V. DADO, M.D.†

Venous air embolism (VAE) is a well-known complication of neurosurgical procedures, especially those

done in the sitting position.<sup>1,2</sup> Additionally, isolated case reports have also appeared in the literature describing VAE in adults and children having a variety of extra-cranial procedures performed.<sup>3-11</sup> We describe a case in which venous air embolism occurred in a child having a large scalp flap raised after it had been delayed and enlarged with a tissue expander.

#### CASE REPORT

A 14-month-old boy had a tissue expander placed in a sub-galeal pocket to delay and expand a scalp flap in preparation for removal of a giant nevus. Anesthesia consisted of halothane, N<sub>2</sub>O, and O<sub>2</sub>. A 4.5-mm ID oral endotracheal tube was inserted during paralysis from

\* Attending Anesthesiologist, Children's Memorial Hospital; Associate of Clinical Anesthesia, Northwestern University Medical School, Chicago, Illinois.

† Attending Physician, Children's Memorial Hospital; Assistant Professor of Surgery, Loyola University Medical School, Division of Plastic Surgery, Maywood, Illinois.

Received from the Department of Anesthesiology, Children's Memorial Hospital, 2300 Children's Plaza, Chicago, Illinois 60614. Accepted for publication October 12, 1987.

Address reprint requests to Dr. Roth.

Key words: Anesthesia; pediatric. Embolism; air. Surgery; plastic.

pancuronium, 0.1 mg/kg iv and ventilation was mechanically controlled. The anesthesia and operation lasted 70 min and was uneventful.

At age 17 months, this child returned to the operating room for removal of the tissue expander, excision of the scalp nevus, and closure of wound with flap advancement. The child weighed 12 kg and had an estimated blood volume of 900 cc. Monitoring included ECG, blood pressure cuff, chest stethoscope, rectal temperature, and inspired oxygen concentration. Anesthesia consisted of 1% halothane, N<sub>2</sub>O, and O<sub>2</sub> via mask, followed by iv pancuronium 0.1 mg/kg, orotracheal intubation, and mechanical ventilation. The child was placed in the supine position. His head was draped in a manner to allow side-to-side movement. The site of the incision was level with the heart at its lowest point and 7 cm above the heart at its highest. Surgery proceeded uneventfully for 1 h. At this time, the nevus had been excised, the tissue expanders removed, and flaps raised in the subgaleal plane. Moist laps were placed under the delayed flaps for hemostasis. The periosteum had not been incised or elevated. The estimated blood loss was 100 ml. Five per cent dextrose in lactated Ringer's solution had been infused. Suddenly arterial blood pressure (BP) decreased from 90 mmHg systolic to become unobtainable; the ECG showed a sinus rhythm increasing from 110 bpm to 125 bpm. No heart tones were heard and a pulse was not palpable. The presence of breath sounds was confirmed bilaterally. All anesthetics were discontinued immediately, 100% O<sub>2</sub> administered, chest compression begun, and epinephrine, 200 µg, was administered iv. BP returned to 80 mmHg systolic and later to 110 mmHg while the heart rate increased to 170 bpm.

An arterial line and precordial Doppler were placed. Arterial blood gas revealed a pH<sub>a</sub> of 7.66, PaCO<sub>2</sub> of 15 mmHg, a PaO<sub>2</sub> of 508 mmHg, BE of 1.6 mEq/l, and a Hct of 25%. Packed red blood cells, 60 ml, were administered iv. Anesthesia and surgery was then resumed with 1% halothane, air, and O<sub>2</sub> to allow the scalp wound to be closed. As the surgeon lifted the scalp flap, air sounds were immediately heard on the Doppler and the BP suddenly became unobtainable again. Oxygen, 100%, was administered, chest compressions begun, and 200 µg of epinephrine was administered iv. The head of the patient was lowered below the level of the heart while the surgeon flooded the field with saline. BP immediately returned to 100 mmHg while the heart rate increased to 170 bpm. The scalp was quickly closed while the patient was in the Trendelenburg position. The child was awakened and observed overnight in the Intensive Care Unit. A follow-up neurologic examination and CAT Scan of the brain were normal.

## DISCUSSION

This is the first reported case of VAE occurring with the removal of a tissue expander from under a delayed scalp flap. In general, air enters the vessel when venous pressure at the surgical site is lower than atmospheric pressure and the vessel is patent. Vessel patency is often maintained when vessels are imbedded in bony or fibrous structures that prevent retraction and collapse of the cut end of the vessel, thus explaining the association between VAE and craniotomies. Children are especially at risk for unexpected reductions in venous pressure, since acute blood loss is more difficult to estimate. Seemingly small amounts of blood loss may represent a significantly larger proportion of estimated blood volume, and may result in undetected low venous pressure.

Previously, a variety of operations have been associated with an increased risk for VAE, the most com-

mon being those in which the patient is undergoing a craniotomy in the sitting position.<sup>12</sup> VAE has occurred in extra-cranial procedures, such as mastectomy<sup>3</sup> and surgery for head and neck tumors,<sup>4</sup> diagnostic and therapeutic air injections,<sup>5,6</sup> venous catheter placement,<sup>7,8</sup> and chest injury,<sup>9,10</sup> and during total hip replacement.<sup>11</sup> Small children with their proportionately larger head size and abundant venous channels are especially prone to VAE during intracranial surgery, even while in the supine position.<sup>13,14</sup> Usually, the surgical procedures associated with an increased risk for VAE involve elevation of bony flaps, large patent blood vessels, and elevated extravascular pressure.

Our patient may have been at risk for VAE for multiple reasons. First, blood loss of approximately 100 ml (10–15% of estimated blood volume) occurred from the scalp incision. While blood replacement was felt to be adequate, venous pressure may have been lower than expected. Second, the placement of a tissue expander with rapid stretching and growth of the scalp may have promoted enlargement of previously present venous channels, as well as the development of new ones. An increased blood flow occurs in the microcirculation of flaps that have been delayed in pigs.<sup>15</sup> Indeed, examining human skin histologically has revealed capillaries and small vessels that are dilated in the dermis and an established vascular layer in the capsule filled with dilated blood vessels and newly formed small vessels.<sup>16</sup> Furthermore, the tough fibrous capsule that develops under an expanded flap may inhibit vessel retraction and contraction when it is cut. Emissary veins originating intracranially and perforating the skull to drain into the scalp venous system probably provided an entry point for VAE.<sup>17</sup> Finally, surgical manipulation of the tissue flap may have elevated an open vessel in relation to the level of the heart, producing a further drop in intravascular pressure at the surgical site, as well as mechanically stenting open the venous channels.

With the increasing use of tissue expansion in the head and neck region of children, an awareness of the increased risk of VAE is essential. Adequate hydration, blood replacement,<sup>18</sup> and being cognizant of the level of surgical manipulation in relation to the heart can decrease the risk of VAE. For immediate recognition of VAE, non-invasive monitoring is well justified. This should include ECG, esophageal stethoscope, precordial Doppler, and measurement of end-tidal CO<sub>2</sub> concentration. The roll of a CVP remains controversial at this time.<sup>19,20</sup>

Compressing the jugular veins can increase venous pressure at the site of the incision, limiting the entrance of air into the venous system. The application of PEEP, although somewhat controversial because of the risks of elevated right atrial pressure, can also raise venous

pressure, impeding the accumulation of air in the venous system.<sup>21</sup> Most importantly, however, is early recognition of air embolism to allow for rapid and effective treatment and to minimize postoperative sequelae.

### REFERENCES

1. Porter SS, Boyd RC, Albin MS: Venous air embolism in a child undergoing posterior fossa craniotomy: A case report. *Can Anaesth Soc J* 31:86-90, 1984
2. Cucchiara RF, Bowers B: Air embolism in children undergoing suboccipital craniotomy. *ANESTHESIOLOGY* 57:338-339, 1982
3. Alexander JI, Lewis AAM: Air embolism during mastectomy. *Anaesthesia* 24:618-619, 1969
4. Kornblut AD, Shumrick DA: Complications of head and neck surgery. *Arch Otolaryngol* 94:246-254, 1971
5. Durant TM, Long J, Oppenheimer MJ: Pulmonary (venous) air embolism. *Am Heart J* 33:3:269-281, 1947
6. Root B, Levy MN, Pollack S, Lubert M, Pathak K: Gas embolism death after laparoscopy delayed by "trapping" in portal circulation. *Anesth Analg* 57:232-237, 1978
7. Conahan III TJ: Air embolization during percutaneous Swan-Ganz catheter placement. *ANESTHESIOLOGY* 50:360-361, 1979
8. Leicht CH, Waldman J: Pulmonary air embolism in the pediatric patient undergoing central catheter placement: A report of two cases. *ANESTHESIOLOGY* 64:519-521, 1986
9. Cleveland JC: Fatal air embolism to the right side of the heart during pneumonectomy for carcinoma. *Chest* 71:4, 1977
10. Thomas AN, Stephens BG: Air embolism: A cause of morbidity and death after penetrating chest trauma. *J Trauma* 14:8:633-638, 1974
11. Andersen KH: Air aspirated from the venous system during total hip replacement. *Anaesthesia* 38:1175-1178, 1983
12. Symons NLP, Leaver HK: Air embolism during craniotomy in the seated position: A comparison of methods for detection. *Can Anaesth Soc J* 32:174-177, 1985
13. Harris M, Strafford MA, Rowe RW, Sanders SP, Winston KR, Rockoff MA: Venous air embolism and cardiac arrest during craniectomy in a supine infant. *ANESTHESIOLOGY* 65:547-550, 1986
14. Harris M, Yement T, Strafford M, Davidson A, Rowe R, Sanders SP, Rockoff MA: Venous air embolism during craniectomies in supine infants. *ANESTHESIOLOGY* 63:A424, 1985
15. Sasaki GH, Pang CY: Pathophysiology of skin flaps raised on expanded pig skin. *Plast Reconstr Surg* 74:59-65, 1984
16. Pasyk KA, Argenta LC, Austuu ED: Histopathology of human expanded tissue. *Clinics in Plastic Surgery*. Edited by Argenta LC, Austud ED. Philadelphia, W. B. Saunders, 1987, pp 435-445
17. Gray H: *Anatomy, Descriptive and Surgical*. Philadelphia, Running Press, 1974, pp 605-606
18. Colohan ART, Perkins NAK, Bedford RF, Jane JA: Intravenous loading as prophylaxis for paradoxical air embolism. *J Neurosurg* 62:839-842, 1985
19. Michenfelder JD: Central venous catheters in the management of air embolism: Whether as well as where. *ANESTHESIOLOGY* 55:339-341, 1981
20. Cucchiara RF, Bowers B: Air embolism in children undergoing suboccipital craniotomy. *ANESTHESIOLOGY* 57:338-339, 1982
21. Pearl RG, Larson CP Jr: Hemodynamic effects of positive end-expiratory pressure during continuous venous air embolism in the dog. *ANESTHESIOLOGY* 64:724-729, 1986

Anesthesiology  
68:459-462, 1988

## Effect and Interaction of pH and Lidocaine on Epinephrine Absorption

WASA UEDA, M.D., PH.D.,\* MASAHISA HIRAKAWA, M.D., PH.D.,† KOREAKI MORI, M.D., PH.D.‡

To achieve optimal local hemostasis in patients undergoing surgery, a dilute solution of epinephrine is often injected. We previously demonstrated that lidocaine added to an epinephrine solution accelerates the transfer of epinephrine to the blood in patients under-

going craniotomy.<sup>1</sup> Sosis *et al.* suggested that the pH of the solution might have been responsible for the variation in epinephrine uptake attributed to lidocaine,<sup>2</sup> because the pH of commercially available lidocaine with epinephrine is made acidic to achieve stability of the added epinephrine. Therefore, in this study, we investigated the effect of the pH of the solution on the absorption of locally injected epinephrine into the blood stream.

### MATERIALS AND METHODS

After obtaining approval of the committee for the protection of human subjects and informed consent, we studied 40 (18 males and 22 females) ASA I and II adult

\* Associate Professor of Anesthesiology.

† Professor of Anesthesiology.

‡ Professor of Neurosurgery.

Received from the Departments of Anesthesiology and Neurosurgery, Kochi Medical School, Kochi, 781-51, Japan. Accepted for publication October 12, 1987.

Address reprint requests to Dr. Ueda: Department of Anesthesiology, Kochi Medical School, Kochi, 781-51, Japan.

Key words: Acid-base equilibrium; pH. Anesthetics, local; lidocaine. Sympathetic nervous system; catecholamine; epinephrine