

iment above for an entire cardiac cycle; note that the portion used for analysis of T is between maximum negative dP/dt and mitral valve opening (estimated as LVEDP of the preceding beat). The plot was generated using high fidelity pressure transducers, an analog-to-digital convertor, and special computer software to produce dP/dt from the digitized pressures and time. Since Swanson and Muir did not use an A/D convertor to capture their data, the differentiated pressures might be difficult for them to obtain.

We cannot predict whether the method of analysis herein proposed would change the conclusions of Swanson and Muir. Thompson *et al.*⁵ have shown that techniques using equation 1 underestimate T. In fact, however, T may be either overestimated or underestimated, since the P axis intercept (P_{asym}) from equation 4 may be positive or negative (cf figure 3 from reference 6). The error becomes greater as P_{asym} becomes increasingly different from zero. When values of T are compared between interventions that may change P_{asym} (such as halothane or ischemia), both the absolute values and the conclusions may suffer. Most investigators have abandoned the equation 1 model.⁴⁻⁶

Finally, it is well to note that the pressure-time asymptote (or the pressure axis intercept in the dP/dt versus P plot) is not necessarily identical to the actual physiologic pressure to which the system decays. The issue is really the "apparent" value of P_{asym} that applies over the range of P(t) that is analyzed for T. Over another pressure range, different values of both P_{asym} and T may be obtained. An outstanding feature of the dP/dt versus P display is that simple inspection will reveal the extent to which any portion of the relationship does or does not follow the presumed monoexponential fall-off.

CHARLES BEATTIE, PH.D., M.D.
LINDA S. HUMPHREY, M.D.
GARY MARUSCHAK, R.C.P.T.

Anesthesiology
70:160, 1989

In Reply:—Drs Beattie *et al.* point out a fundamental analytical error in our data describing left ventricular relaxation. We agree with their assessment, and appreciate their critical reading of our manuscript.

CLIFFORD R. SWANSON, D.V.M., M.S.
Assistant Professor of Anesthesiology
Department of Anatomy,
Physiological Sciences, and Radiology
North Carolina State University
4700 Hillsborough Street at William
Moore Drive
Raleigh, North Carolina 27606

Anesthesiology
70:160-161, 1989

Use Caution when Extrapolating from a Small Sample Size to the General Population

To the Editor:—Sears *et al.* recently reported "that the administration of a second dose of succinylcholine to healthy adult patients after induction with ketamine is safe with respect to cardiac rate and rhythm."¹ They based this conclusion on the results of a study performed on eight patients. We believe their conclusion is too strong. Because they encountered no dysrhythmias and did not have a statistically significant

Department of Anesthesiology and Critical Care Medicine
The Johns Hopkins Medical Institutions
600 North Wolfe Street
Baltimore, Maryland 21205

REFERENCES

1. Swanson CR, Muir WW: Simultaneous evaluation of left ventricular end-systolic pressure-volume ratio and time constant of isovolumic pressure decline in dogs exposed to equivalent MAC halothane and isoflurane. *ANESTHESIOLOGY* 68:764-770, 1988
2. Simon AC, Safar ME, Lenenson JA, London GM, Levy BI, Chau HP: An evaluation of large arteries compliance in man. *Am J Physiol* 235:H550-H554, 1979
3. Liu Z, Brin KP, Yin FC: Estimation of total arterial compliance: An improved method and evaluation of current methods. *Am J Physiol* 251 (Heart Circ Physiol 20):H588-H600, 1986
4. Weisfeldt ML, Weiss JL, Frederiksen JT, Yin FCP: Quantification of incomplete left ventricular relaxation: Relationship to the time constant for isovolumic pressure fall. *Eur Heart J (Suppl A)*:119-129, 1980
5. Thompson DS, Waldron CB, Coltart DJ, Jenkins BS, Webb-Peploe MM: Estimation of time constant of left ventricular relaxation. *Br Heart J* 49:250-258, 1983
6. Humphrey LS, Topol EJ, Rosenfeld GI, Borkon AM, Baumgartner WA, Gardner TJ, Maruschak G, Weiss JL: Immediate enhancement of left ventricular relaxation by coronary artery bypass grafting: intraoperative assessment. *Circulation* 77:886-896, 1988

(Accepted for publication August 24, 1988.)

WILLIAM W. MUIR III, D.V.M., PH.D.
Professor and Chairman of Anesthesiology
Department of Veterinary Clinical Sciences
College of Veterinary Medicine
The Ohio State University
Columbus, Ohio 43210

(Accepted for publication August 24, 1988.)

decrease in heart rate does not imply the true incidence of these undesirable side effects is insignificant.

Whenever the numerator is zero in the incidence of an effect, the true incidence in the population at large represented by the group is:

$\sqrt[3]{p}$

where n is the sample size and p is the level of credibility, *i.e.*, if 95% confidence is required, then $p = 0.05$.² If $n = 8$ and $p = 0.05$ as used by Sears, the maximum risk for significant bradydysrhythmias in a very large group of similar patients is actually 31%. Since bradydysrhythmias are undesirable and the true incidence could be as high as 31% based on Sears' study, we believe their conclusion would be more correctly stated that the incidence of bradydysrhythmias with a second dose of succinylcholine after a ketamine induction is (with 95% confidence) no higher than 31%.

When n is greater than 30, the computation is considerably simplified, and is referred to as the rule of 3.³ When $p = 0.05$, "if none of n patients shows the event about which we are concerned, we can be 95% confident that the chance of this event is at most 3 in n , (*i.e.*, 3/ n)."² Thus, if Sears *et al.* had observed zero incidence in 50 subjects, they could have predicted the maximum incidence to be no higher than 6% (3/50) in a similar but large group.

When negative results are reported, the author should be careful about extrapolating results to the universe of patients as a whole.

DAVID J. BENEFIEL, M.D.
Director of Cardiovascular Anesthesia

Anesthesiology
70:161, 1989

In Reply:—We agree with Benefiel *et al.* that our statement "the administration of a second dose of succinylcholine to healthy adult patients after induction with ketamine is safe with respect to cardiac rate and rhythm" may be too strong if only statistical analysis is considered. However, in clinical situations, other factors, such as patient's physical status, the pharmacological properties of the drug, or combination of drugs, must also be considered in the clinical conclusion of a particular study. In our study,¹ although the number is small ($n = 8$), we believe that our conclusion may not be too strong. This conclusion is not only based on the statistical analysis but also on the physical status of the patients (healthy adults) and on the pharmacodynamic property of ketamine as a drug with a sympathomimetic effect.²

The practical problem is the number of patients one should study before concluding that a technique is safe or dangerous. Based on our study, we feel it is appropriate to continue to use a second dose of succinylcholine after ketamine where required and continue to observe the patient response. If any serious side effects are encountered, we will of course report them. Thus far we still have not seen any problems with a second dose of succinylcholine in patients who received ketamine.

IMAD H. ABDUL-RASOOL, M.D., PH.D.
Assistant Professor of Anesthesiology

Anesthesiology
70:161-162, 1989

Succinylcholine and Trismus

To the Editor:—The recent publication of various articles and letters to the editor about succinylcholine and trismus has resulted in a re-evaluation of the use of succinylcholine and a reassessment of our clinical practice when trismus occurs after the administration of succinylcholine.¹⁻⁴ The major problem in this whole issue centers around the clinical judgment issue of what constitutes trismus. Unfortunately, Dorland's *Medical Dictionary* does not clarify the issue, nor is it clarified in any of

the papers that have been written. The reason is that it is a clinical judgment call with a broad spectrum of possibilities. At one end of the spectrum, the masseter muscle tone may be so increased that there is a complete inability to open the mouth. At the other end of the spectrum, there may be a mild increase in muscle tone that can be easily overcome. Van Der Spek has certainly thrown an entirely different light on the issue of succinylcholine and "trismus," with the finding

EDWARD A. EISLER, M.D.
Director of Obstetrical Anesthesia

RODGER SHEPHERD, M.D., M.P.H.
Department of Epidemiology

*Pacific Presbyterian Medical Center
Clay at Buchanan Street
San Francisco, California 94120*

REFERENCES

1. Sears SH, Abdul-Rasool IH, Katz RL: The effect of a second dose of succinylcholine on cardiac rate and rhythm following induction of anesthesia with ketamine. *ANESTHESIOLOGY* 68:644, 1988
2. Hanley JA, Lippman-Hand A: If nothing goes wrong, is everything all right? *JAMA* 249:1743-1745, 1983
3. Rumke CL: Implications of the statement: No side effects were observed. *N Engl J Med* 292:372-373, 1975

(Accepted for publication September 14, 1988.)

DANIEL H. SEARS, M.D.
Staff Anesthesiologist

RONALD L. KATZ, M.D.
*Professor of Anesthesiology and Chairman
of Anesthesiology*

*UCLA School of Medicine
10833 LeConte Avenue
Los Angeles, California 90024-1778*

REFERENCES

1. Sears DH, Abdul-Rasool IH, Katz RL: The effect of a second dose of succinylcholine on cardiac rate and rhythm following induction of anesthesia with ketamine. *ANESTHESIOLOGY* 68:144, 1988
2. White PF, Way WL, Trevor AJ: Ketamine—Its pharmacology and therapeutic uses. *ANESTHESIOLOGY* 56:119-136, 1982

(Accepted for publication September 23, 1988.)