

Fentanyl-induced Muscle Rigidity in Unanesthetized and Ketamine- or Thiopental-anesthetized Rats

Ping-Wing Lui, M.D.,* Tak-Yu Lee, M.D.,† Samuel H. H. Chan, Ph.D.‡

This study was undertaken to search for an alternative experimental model in the evaluation of fentanyl-induced muscle rigidity. Unanesthetized, spontaneously ventilating Sprague-Dawley rats, and rats anesthetized with either ketamine or thiopental whose ventilation was mechanically controlled, were studied. Intravenous administration of fentanyl (25, 50, or 100 µg/kg) caused an increase in electromyographic (EMG) activity in both unanesthetized and ketamine-anesthetized, but not in thiopental-anesthetized, animals. Muscle rigidity was more prominently manifested in the gastrocnemius muscle, when compared with the rectus abdominis muscle. Hypoxemia was exhibited during the course of rigidity by both spontaneously ventilating and ketamine-anesthetized rats, but not by thiopental-anesthetized animals. In addition, unanesthetized, spontaneously ventilating rats developed hypercarbia and respiratory acidosis. The authors suggest that, in addition to using unanesthetized animals, EMG activity in the gastrocnemius muscle of rats anesthetized with ketamine in whom ventilation is controlled may provide an alternative approach in the evaluation of fentanyl-induced muscle rigidity. (Key words: Analgesics: fentanyl. Anesthetics, intravenous: ketamine; thiopental. Complications, fentanyl: rigidity. Measurement techniques: electromyography. Muscle: rigidity.)

HIGH DOSES OF POTENT opioids, including fentanyl^{1,2} and alfentanil,³ are frequently administered in clinical practice. This anesthetic technique provides good analgesia and stable hemodynamic responses. Unfortunately, it is also commonly accompanied by muscle rigidity, especially during the induction of anesthesia.^{2,4,5} This opiate-induced rigidity may in turn be associated with decrease in chest wall compliance,⁶ increase in intracranial pressure secondary to elevation in central venous pressure,^{7,8} and possible respiratory and acid-base disturbances.⁴

* Staff Anesthesiologist, Veterans General Hospital; Lecturer of Anesthesiology, and Ph.D. candidate, Institute of Clinical Medicine, National Yang-Ming Medical College.

† Director of Anesthesiology, Veterans General Hospital; and Associate Professor of Anesthesiology, National Yang-Ming Medical College.

‡ Professor and Director of Pharmacology, National Yang-Ming Medical College.

Received from the Institutes of Clinical Medicine and Pharmacology, National Yang-Ming Medical College, and Department of Anesthesiology, Veterans General Hospital, Taipei, Taiwan, Republic of China. Accepted for publication January 31, 1989. Supported in part by Clinical Research Center Grant #46500-005 from the Veterans General Hospital (PWL), NSC78-0412-B075-40 (TYL) and NSC77-0412-B010-32 (SHHC) from the National Science Council, and a Tjing-Ling Medical Foundation Professorship (SHHC), Taiwan, Republic of China.

Address reprint requests to Dr. Chan: Institute of Pharmacology, National Yang-Ming Medical College, Taipei 11221, Taiwan, Republic of China.

The precise mechanism(s) that underlies opiate-induced muscle rigidity has yet to be completely elucidated. This may be due in part to the lack of a suitable animal preparation for its evaluation. A frequently employed approach,⁹⁻¹⁵ first reported by Wand *et al.*,⁹ uses the electromyographic (EMG) activity in unanesthetized, spontaneously ventilating animals as the experimental model. Inherent in this approach are possible confounding factors of hypoventilation, hypoxemia, hypercarbia, and bradycardia, all of which are commonly associated with high doses of opiates.^{1,4,16} A possible solution is to control respiration during the experiment. However, an anesthetic state must be induced in the animal before ventilation can be controlled. Unfortunately, many anesthetics, such as barbiturates^{7,17,18} and benzodiazepines,¹⁹ are reported to attenuate opiate-induced rigidity.

The purpose of this study was to search for an alternative approach in the evaluation of fentanyl-induced muscle rigidity. We suggest that a combination of the advantages from both above mentioned methods may result in an acceptable experimental preparation. Using EMG activity in unanesthetized, spontaneously ventilating rats and animals anesthetized with either ketamine or thiopental whose ventilation was controlled, we thus sought answers to four basic questions. First, is rigidity related to the state of consciousness? Second, will different anesthetic agents influence rigidity differently? Third, is rigidity equally expressed in both extremity and trunk muscles? Fourth, what are the alterations in hemodynamic and blood gas parameters that may accompany fentanyl-induced muscle rigidity during consciousness and anesthesia?

Materials and Methods

SUBJECTS

Sixty-nine adult, male Sprague-Dawley rats (220-300 g) were used. They were housed three per cage with free access to food and water, and were maintained on a 12:12 h light-dark schedule in a temperature-controlled room (23 ± 1°C). All procedures described below were carried out in accordance with institutional guidelines.

SURGICAL PREPARATIONS

Unanesthetized Rats. During pentobarbital sodium (40 mg/kg, ip) anesthesia, an arterial cannula (Clay Adams,

PE-50) was inserted into the right femoral artery for systemic arterial pressure measurement and blood sampling. The right femoral vein was also cannulated for the injection of drugs. Both catheters were guided percutaneously to the neck of the animal, where they were filled with heparinized saline (100 U/ml) and sealed with metal plugs.

A pair of platinum needle electrodes (Grass, Type E2) was inserted intramuscularly into the bilateral rectus abdominis and left gastrocnemius muscles, respectively, and was secured to the skin of the animal by sutures and micro-pore tape. The distance between each pair of electrodes was about 0.5–1.0 cm. Procaine penicillin (30,000 U, im) was given at the end of surgery. Each animal was thereafter housed individually. At least 3 days were allowed for recovery before the commencement of the experimental session.

Anesthetized Rats. Animals were anesthetized with either ketamine (100–120 mg/kg, ip) or thiopental sodium (50 mg/kg, ip). The right femoral artery and vein were cannulated with heparinized saline-filled catheters (Clay Adams, PE-50). A pair of platinum needle electrodes was inserted intramuscularly into the bilateral rectus abdominis and left gastrocnemius muscles, respectively, for EMG recording.

The lungs were ventilated with ambient air *via* a tracheostomy, using a rodent ventilator (Harvard, 683). The end-tidal CO₂ was monitored with a Capnometer (Datex) and maintained between 4.0–5.0%. This corresponds to a PaCO₂ of 35–45 mmHg as obtained by blood gas analysis. The rectal temperature of the animal was maintained at 37 ± 0.5° C using a heating lamp. Supplemental ketamine (5 mg/kg, iv) or thiopental (2 mg/kg, iv) was administered whenever movement of the head, extremity, or tail was observed, to maintain the animal in an anesthetized state.

EMG, SYSTEMIC ARTERIAL PRESSURE, AND HEART RATE RECORDING

Raw EMG signals were bipolarly recorded from the gastrocnemius and rectus abdominis muscles and were differentially amplified by a Grass 7P3 preamplifier (bandpass: DC–3 Hz). They were also summated and integrated every 10 s, using an integrator preamplifier (Grass, 7P10). Systemic arterial pressure was measured from the cannulated femoral artery *via* a Statham pressure transducer (P23ID). Heart rate was determined by a biotachometer (Grass, 7P4) triggered by the arterial pulses.

BLOOD GAS ANALYSIS

Arterial blood samples (0.25 ml) were obtained 5 min before fentanyl injection, and 5, 10, and 15 min thereafter. Blood withdrawn was replaced with an equal volume of saline to avoid hypovolemia. PaO₂, PaCO₂, pH, and base

excess were measured using a Radiometer machine (Copenhagen, ABL 300), which was routinely calibrated daily.

EXPERIMENTAL PROTOCOLS

All recording session took place in a quiet room with minimal physical disturbance. The unanesthetized, spontaneously ventilating rats were placed in a cylindrical cage that permitted restricted but unrestrained movement. The anesthetized animals remained unrestrained in a prone position.

Raw and integrated EMG signals from the gastrocnemius and rectus abdominis muscles, systemic arterial pressure, heart rate, and end-tidal CO₂ concentration (anesthetized animals only) were continuously displayed on a multi-channel Grass 7D polygraph. The effects of intravenously administered fentanyl, delivered over 15 s, on these parameters were observed for 15 min. Correct placements of EMG electrodes were verified by necropsy immediately after each experiment.

An equal volume of venous blood (0.2 ml) was slowly withdrawn prior to fentanyl injection to avoid hypervolemia. Vehicle control groups received an equal amount of saline. To avoid the confounding actions of multiple dosing and tachyphylaxis, each animal received only a single injection of fentanyl or saline.

DRUGS

Drugs used in this study included fentanyl citrate (Janssen), ketamine HCl (Parke-Davis), and thiopental sodium (Lilly).

STATISTICS

The EMG activity from each experiment was first quantified by averaging the integrated signals recorded over each 5-min interval (30 datum points) during the observation period. These averaged values were further normalized to a percent of predrug control to compensate for interanimal variations. The effects of fentanyl or vehicle on these EMG signals from the gastrocnemius and rectus abdominis muscles, mean arterial pressure, heart rate, PaO₂, PaCO₂, pH, and base excess were statistically assessed using analysis of variance (ANOVA) with repeated measurements. This was followed by the Student-Newman-Keuls multiple range test for a *posteriori* analysis of individual means (fentanyl *versus* vehicle, high *versus* low dose of fentanyl) at 5, 10, and 15 min postinjection. A *P* value < 0.05 was taken to be statistically significant.

Results

EFFECTS OF FENTANYL IN UNANESTHETIZED RATS

Intravenous administration of fentanyl (25 µg/kg) was followed by a sustained and dramatic increase in EMG




FIG. 1. Representative data from a single animal demonstrating the effects of fentanyl (25 µg/kg, iv) on electromyographic (EMG) activity of left gastrocnemius (GC) and rectus abdominis (RA) muscles, systemic arterial pressure (SAP), and heart rate (HR) in unanesthetized, spontaneously ventilating rats. The raw EMG signals were summated and integrated over 10 s for quantitative analysis.

activity from the gastrocnemius muscle (figs. 1, 2). The onset of muscle rigidity (fig. 1), defined as the time taken (from the end of injection) for the EMG signal to increase by at least 100% over the control, was almost immediate, and the peak effect occurred between 1–2 min thereafter. The enhanced activity from the rectus abdominis muscle, on the other hand, was rather transient in duration (fig. 1) and mild in intensity (fig. 2).

Accompanying the increased muscle activity were severe and prolonged hypotension and bradycardia (fig. 2), which began immediately (fig. 1) following the administration of fentanyl. There was also an apneic period that

lasted 15–35 s, followed by 2–3 min of severe hypoventilation. Arterial blood gas analysis (table 1) documented the occurrence of respiratory acidosis with hypoxemia and hypercarbia.

It should be mentioned that a higher dose of fentanyl (50 µg/kg, iv) was administered in five unanesthetized, spontaneously ventilating animals. They were excluded from statistical analysis, since all of them succumbed to the abovementioned detrimental respiratory, hemodynamic, and acid-base variables, which were further intensified.

EFFECTS OF FENTANYL IN RATS ANESTHETIZED WITH KETAMINE

Two intravenous doses (50 and 100 µg/kg) of fentanyl were used in this series. A significant ($P < 0.05$) and persistent increase in EMG activity from the gastrocnemius muscle was noted 30–60 s after the administration of fentanyl (figs. 3, 4), in a dose-dependent manner. There was,




FIG. 2. Effects of intravenous administration of fentanyl (25 µg/kg, filled bar) and saline (open bar) on electromyographic (EMG) activity of left gastrocnemius and rectus abdominis muscles, mean arterial pressure (MAP), heart rate (HR), and PaO_2 in unanesthetized, spontaneously ventilating rats. Values are mean \pm SEM ($n = 5$ in each group). * $P < 0.01$, ** $P < 0.005$ versus saline in the Student-Newman-Keuls analysis.

TABLE 1. Arterial Blood Gas Parameters following Intravenous Administration of 25 µg/kg of Fentanyl (A) and Saline (B) in Conscious Rats. Controls were Obtained 5 Min before Fentanyl/Saline.

		Control	Postinjection		
			5 Min	10 Min	15 Min
pH	A	7.307 ± 0.012	7.271 ± 0.015†	7.313 ± 0.012*	7.359 ± 0.013
	B	7.410 ± 0.014	7.396 ± 0.021	7.370 ± 0.017	7.391 ± 0.021
PaO ₂	A	104 ± 2	51 ± 3†	71 ± 3†	78 ± 3†
	B	112 ± 3	107 ± 3	108 ± 3	105 ± 2
PaCO ₂	A	39.3 ± 0.8	57.7 ± 1.8†	48.1 ± 2.3†	44.0 ± 2.1
	B	38.8 ± 1.3	40.2 ± 1.2	40.4 ± 1.5	38.6 ± 1.6
BE	A	0.6 ± 0.4	-2.4 ± 0.5*	-3.8 ± 0.4†	-2.1 ± 0.3†
	B	0.7 ± 0.4	0.3 ± 0.1	-0.1 ± 0.4	-0.4 ± 0.3

All values are mean ± SEM (n = 5 in each group).

* P < 0.01, †P < 0.005 vs. saline in the Student-Newman-Keuls analysis.

on the other hand, no apparent alteration in the electric activity of the rectus abdominis muscle (figs. 3, 4). Hypotension and bradycardia were seen immediately after the injection of fentanyl. Blood gas analysis (table 2) indicated that both arterial PaCO₂ and pH were within normal ranges. The animals did manifest hypoxemia during the course of fentanyl-induced muscle rigidity (fig. 4), even when ventilation was controlled.

EFFECTS OF FENTANYL IN RATS ANESTHETIZED WITH THIOPENTAL

Not only did both doses of fentanyl (50 and 100 µg/kg, iv) fail to cause an increase in EMG activity from the gastrocnemius and rectus abdominis muscles, there was in fact a significant (P < 0.05) decrease in the activity of the former muscle (fig. 5) at the higher dose. A reduction in systemic arterial pressure and heart rate still occurred, with no apparent alterations in any respiratory or acid-base parameter (table 3).

Discussion

The present study revealed that intravenous administration of fentanyl promoted an increase in EMG activity in both unanesthetized, spontaneously ventilating rats and rats anesthetized with ketamine. Furthermore, this correlate of opiate-induced muscle rigidity was preferentially manifested by the gastrocnemius muscle, when compared to the rectus abdominis muscle. Both groups of animals exhibited hypoxemia during the course of rigidity. The former, in addition, manifested respiratory acidosis and hypercarbia. Fentanyl did not, on the other hand, evoke




FIG. 3. Representative data from a single animal demonstrating the effects of fentanyl (50 µg/kg, iv) on electromyographic (EMG) activity of left gastrocnemius (GC) and rectus abdominis (RA) muscles, systemic arterial pressure (SAP), heart rate (HR), and end-tidal CO₂ (ETCO₂) in rats anesthetized with ketamine and whose lungs were mechanically ventilated.




FIG. 4. Effects of intravenous administration of fentanyl at 50 (stippled bar) and 100 (filled bar) $\mu\text{g}/\text{kg}$, and saline (open bar) on electromyographic (EMG) activity of left gastrocnemius (GC) and rectus abdominis (RA) muscles, mean arterial pressure (MAP), heart rate (HR), and PaO_2 in rats anesthetized with ketamine and whose lungs were mechanically ventilated. Values are mean \pm SEM ($n = 10$ in each group). * $P < 0.01$ versus saline and † $P < 0.01$ versus 50 $\mu\text{g}/\text{kg}$ of fentanyl in the Student-Newman-Keuls analysis.

any apparent EMG activation, nor alterations in PaO_2 , PaCO_2 , $p\text{H}$, and base excess in rats anesthetized with thiopental.

Within the limits of comparing the effects of two similar opioids in different species, our data from unanesthetized, spontaneously ventilating rats are in partial agreement with the observations by Benthuisen *et al.*⁷ These authors reported that alfentanil induces in their patients marked rigidity in eight muscle groups, including the gastrocnemius and rectus abdominis muscles. Also present in their subjects were respiratory acidosis, hypoxemia, hypercarbia, and hypotension. Our results are also compatible with those from the report by Comstock *et al.*,⁴ who showed that truncal rigidity and hypercarbia ($\text{PaCO}_2 = 52.1 \pm 1.8$ mmHg) were induced in 20 out of 21 patients following high-dose fentanyl (19 ± 1.9 $\mu\text{g}/\text{kg}$, iv) induction of anesthesia.

Ketamine is unique in producing an unusual trance-like state known as dissociative anesthesia.^{20,21} Similar to unanesthetized rats, we observed that animals anesthetized with this agent responded to fentanyl with an increase in EMG activity, but, presumably because ventilation was controlled, failed to exhibit hypercarbia and acidosis. It might be argued that the muscular hypertonus inherent in ketamine anesthesia²⁰ could be a confounding factor. That this may not be the case is amply suggested by the profound enhancement in EMG activity, as opposed to the vehicle control, promoted by fentanyl. Nonetheless, an increase in oxygen consumption secondary to these heightened muscular activity may account for the hypoxemia developed in spite of controlled artificial ventilation.⁷

Kissin *et al.*²² suggested that the interactive effects of fentanyl and thiopental may be different, depending on the endpoint of measurement. Thus, these two pharmacologic agents are synergistic with reference to the loss of righting reflex,²³ but are antagonistic in relation to purposeful movement response to tail clamping.²² The present study provides an additional dimension to this

TABLE 2. Arterial Blood Gas Parameters following Intravenous Administration of 50 $\mu\text{g}/\text{kg}$ (A), 100 $\mu\text{g}/\text{kg}$ (B) of Fentanyl or Saline (C) in Rats Anesthetized with Ketamine and Whose Lungs Were Mechanically Ventilating.

		Control	Postinjection		
			5 Min	10 Min	15 Min
$p\text{H}$	A	7.384 ± 0.013	7.371 ± 0.010	7.357 ± 0.009	7.359 ± 0.010
	B	7.393 ± 0.009	7.394 ± 0.006	7.385 ± 0.012	7.390 ± 0.011
	C	7.387 ± 0.010	7.385 ± 0.008	7.381 ± 0.010	7.394 ± 0.008
PaO_2	A	116 ± 3	$68 \pm 3^\dagger$	$61 \pm 2^\dagger$	$62 \pm 3^\dagger$
	B	110 ± 2	$66 \pm 4^\dagger$	$60 \pm 2^\dagger$	$65 \pm 3^\dagger$
	C	108 ± 2	106 ± 2	105 ± 1	103 ± 1
PaCO_2	A	39.2 ± 1.3	39.2 ± 1.6	40.6 ± 1.7	41.9 ± 1.3
	B	41.1 ± 1.6	38.5 ± 2.4	38.0 ± 1.4	37.6 ± 1.2
	C	39.1 ± 1.0	39.5 ± 0.9	38.4 ± 0.7	38.6 ± 0.6
BE	A	-1.1 ± 0.4	-2.3 ± 0.5	$-2.5 \pm 0.4^*$	$-2.2 \pm 0.3^*$
	B	0.2 ± 0.4	-1.4 ± 0.5	$-2.6 \pm 0.5^*$	$-2.1 \pm 0.5^*$
	C	-0.7 ± 0.4	-0.9 ± 0.2	-0.7 ± 0.3	-0.9 ± 0.3

Controls were obtained 5 min before fentanyl/saline. All values are mean \pm SEM ($n = 10$ in each group).

* $P < 0.01$, † $P < 0.005$ vs. saline in the Student-Newman-Keuls analysis.

fentanyl-thiopental interaction. The barbiturate attenuated (and even reduced) the muscular rigidity induced by the narcotic agent. From the standpoint of clinical practice, it appears that thiopental may be a suitable adjunct to fentanyl anesthesia, since it removes a major side-effect of the opiate.^{5,18,19} On the other hand, it will be a poor choice of anesthetic agent because it eliminates the key experimental index in animals in which fentanyl-induced muscle rigidity is being studied.

As pointed out by many authors,^{7,18,19,24} we are aware that some general anesthetics may influence the occurrence of opiate-induced rigidity. We also noted that the magnitude of rigidity elicited by fentanyl (50 µg/kg, iv) in rats anesthetized with ketamine was less than that seen in the unanesthetized animals receiving only 25 µg/kg of the opiate. One may therefore argue that ketamine would not be an ideal anesthetic in an animal in which rigidity was being studied. This seeming deficit, however, must be weighed against the advantages of ketamine anesthesia. First, fentanyl still produces an elevation in EMG activity in animals anesthetized with ketamine that is significantly different from vehicle control, without the detrimental effects of hypercarbia and respiratory acidosis that were present in unanesthetized, spontaneously ventilating rats. Second, ketamine has the benefit of being synergistic with fentanyl in their drug actions. Radioligand binding and behavioral studies²⁵⁻²⁷ indicate that this dissociative anesthetic may interact synergistically with the opiate receptors, possibly the mu-subtype, in the elicitation of catalepsy and analgesia. Thus, it appears that EMG activity in the gastrocnemius muscle of rats anesthetized with ketamine in whom ventilation is controlled may provide an alternative approach in the investigation of opiate-induced rigidity.

The primary purpose of our study was to search for an alternative experimental approach in the evaluation




FIG. 5. Effects of intravenous administration of fentanyl at 50 (stippled bar) and 100 (filled bar) µg/kg, and saline (open bar) on electromyographic (EMG) activity of left gastrocnemius (GC) and rectus abdominis (RA) muscles, mean arterial pressure (MAP), heart rate (HR), and PaO₂ in rats anesthetized with thiopental and whose lungs were mechanically ventilated. Values are mean ± SEM (n = 8 in each group). *P < 0.05 versus saline and †P < 0.05 versus 50 µg/kg of fentanyl in the Student-Newman-Keuls analysis.

of fentanyl-induced muscle rigidity. To be acceptable, this animal model must demonstrate EMG activation by fentanyl that is at least qualitatively similar to that elicited in conscious animals, without, however, the undesirable effects of hypercarbia and respiratory acidosis. It appears that EMG activity in gastrocnemius muscle of rats anes-

TABLE 3. Arterial Blood Gas Parameters following Intravenous Administration of 50 µg/kg (A), 100 µg/kg (B) of Fentanyl and Saline (C) in Rats Anesthetized with Thiopental Whose Lungs Were Mechanically Ventilated.

		Control	Postinjection		
			5 Min	10 Min	15 Min
pH	A	7.395 ± 0.011	7.384 ± 0.012	7.378 ± 0.007	7.391 ± 0.011
	B	7.407 ± 0.016	7.391 ± 0.035	7.405 ± 0.027	7.402 ± 0.023
	C	7.372 ± 0.002	7.372 ± 0.006	7.379 ± 0.008	7.374 ± 0.014
PaO ₂	A	101 ± 2	93 ± 3	97 ± 4	102 ± 3
	B	105 ± 1	95 ± 2	96 ± 3	99 ± 3
	C	104 ± 2	101 ± 3	97 ± 5	108 ± 1
PaCO ₂	A	37.6 ± 0.9	39.5 ± 1.3	41.3 ± 1.2	41.4 ± 2.0
	B	37.2 ± 1.5	38.8 ± 2.2	38.9 ± 2.3	41.0 ± 2.6
	C	38.6 ± 1.1	38.3 ± 0.9	37.4 ± 0.8	37.5 ± 0.9
BE	A	-0.4 ± 0.3	-1.2 ± 0.5	-1.6 ± 0.3	-1.6 ± 0.2
	B	-0.6 ± 0.5	-0.4 ± 0.6	-1.0 ± 0.6	-0.5 ± 0.8
	C	0.3 ± 0.2	-0.7 ± 0.3	-0.8 ± 0.1	-1.1 ± 0.2

Controls were obtained 5 min before fentanyl/saline. All values are mean ± SEM (n = 8 in each group).

thetized with ketamine and in whom ventilation is controlled may satisfy all these stipulations. Based on this experimental model, we recently identified²⁸ that the pontine nucleus, locus coeruleus, and noradrenergic neurotransmission are critically involved in fentanyl-induced muscle rigidity in the rat.

The authors wish to thank Professor B. N. Chiang and Dr. Julie Y. H. Chan for their enthusiastic support.

References

- Stanley TH, Philbin DM, Coggins CH: Fentanyl-oxygen anaesthesia for coronary artery surgery: Cardiovascular and antidiuretic hormone responses. *Can Anaesth Soc J* 26:168-172, 1979
- Bailey PL, Wilbrink J, Zwanikken P, Pace NL, Stanley TH: Anesthetic induction with fentanyl. *Anesth Analg* 64:48-53, 1985
- de Lange S, Stanley TH, Boscoe MJ: Alfentanil-oxygen anaesthesia for coronary artery surgery. *Br J Anaesth* 53:1291-1296, 1981
- Comstock MK, Carter JG, Moyers JR, Stevens WC: Rigidity and hypercarbia associated with high dose fentanyl induction of anaesthesia. *Anesth Analg* 60:362, 1981
- Freud FG, Martin EW, Wong KC, Hornbein TF: Abdominal muscle rigidity induced by morphine and nitrous oxide. *ANESTHESIOLOGY* 38:358-362, 1973
- Scamman FL: Fentanyl-O₂-N₂O rigidity and pulmonary compliance. *Anesth Analg* 62:332-334, 1983
- Benthuysen JL, Smith NT, Sanford TJ, Head N, Dec-Silver H: Physiology of alfentanil-induced rigidity. *ANESTHESIOLOGY* 64:440-446, 1986
- Benthuysen JL, Kien ND, Quam DD: Intracranial pressure increases during alfentanil-induced rigidity. *ANESTHESIOLOGY* 68:438-440, 1988
- Wand P, Kuschinsky K, Sontag KH: Morphine-induced muscular rigidity in rats. *Eur J Pharmacol* 24:189-193, 1973
- Havemann U, Kuschinsky K: Further characterization of opioid receptors in the striatum mediating muscular rigidity in rats. *Naunyn-Schmiedeberg's Arch Pharmacol* 317:321-325, 1981
- De Ryck M, Teitelbaum P: Morphine versus haloperidol catalepsy in the rat: An electromyographic analysis of postural support mechanisms. *Exp Neurol* 79:54-76, 1983
- Blasco TA, Lee D, Amalric M, Swerdlow NR, Smith NT, Koob GF: The role of the nucleus raphe pontis and the caudate nucleus in alfentanil rigidity in the rat. *Brain Res* 386:280-286, 1986
- Rackham A: Opiate-induced muscle rigidity in the rat: Effects of centrally acting agents. *Neuropharmacology* 19:855-859, 1980
- Weinger MB, Cline EJ, Smith NT, Blasco TA, Koob GF: Localization of brainstem sites which mediate alfentanil-induced muscle rigidity in the rat. *Pharmacol Biochem Behav* 29:573-580, 1988
- Slater P, Starkie DA: Changes in limb tone produced by regional injections of opiates into rat brain. *Naunyn-Schmiedeberg's Arch Pharmacol* 335:54-58, 1987
- Hug CC, Murphy MR: Fentanyl disposition in cerebrospinal fluid and plasma and its relationship to ventilatory depression in the dog. *ANESTHESIOLOGY* 50:342-349, 1979
- Hill AB, Nahrwold ML, deRosayro AM, Knight PR, Jones RM, Bolles RE: Prevention of rigidity during fentanyl-oxygen induction of anaesthesia. *ANESTHESIOLOGY* 55:452-454, 1981
- Blasco TA, Smith NT, Sanford TJ, Benthuysen J, Dec-Silver H, Head N: A clinical study of the effects of various pretreatment agents on alfentanil-induced rigidity: EMG data (abstract). *ANESTHESIOLOGY* 63:A380, 1986
- Sanford TJ, Smith NT, Weinger MB, Benthuysen JL, Head N: The effect of midazolam pretreatment on alfentanil-induced muscle rigidity (abstract). *ANESTHESIOLOGY* 69:A556, 1988
- Winters WD, Ferrar-Allado T, Guzman-Flores C, Alcaraz M: The cataleptic state induced by ketamine: A review of the neuropharmacology of anaesthesia. *Neuropharmacology* 11:303-315, 1972
- White PF, Way WL, Trevor AJ: Ketamine—Its pharmacology and therapeutic uses. *ANESTHESIOLOGY* 56:119-136, 1982
- Kissin I, Mason JO, Bradley EL Jr: Morphine and fentanyl interactions with thiopental in relation to movement response to noxious stimulation. *Anesth Analg* 65:1149-1154, 1986
- Kissin I, Mason JO, Bradley EL Jr: Morphine and fentanyl hypnotic interactions with thiopental. *ANESTHESIOLOGY* 67:331-335, 1987
- Hug CC Jr: Anesthetic agents and the patient with cardiovascular disease, *Acute Cardiovascular Management: Anaesthesia and Intensive Care*. Edited by Ream AK, Fogdall RP. Philadelphia, JB Lippincott Co., 1982, pp 247-291
- Smith DJ, Pekoe GM, Martin LL, Coalgate B: The interaction of ketamine with the opiate receptor. *Life Sci* 26:789-795, 1980
- Finck AD, Ngai SH: Opiate receptor mediation of ketamine analgesia. *ANESTHESIOLOGY* 56:291-297, 1982
- Smith DJ, Bouchal RL, deSanctis CA, Monroe PJ, Amedro JB, Perrotti JM, Crisp T: Properties of the interaction between ketamine and opiate binding sites in vivo and in vitro. *Neuropharmacology* 26:1253-1260, 1987
- Lui PW, Lee TY, Chan SHH: Involvement of locus coeruleus and noradrenergic neurotransmission in fentanyl-induced muscular rigidity in the rat. *Neurosci Lett* 96:114-119, 1989