

RICHARD J. TRAYSTMAN, PH.D.
Professor

*Department of Anesthesiology/Critical Care Medicine
Johns Hopkins
600 North Wolfe Street/CMSC 7-110
Baltimore, Maryland 21205*

REFERENCES

1. Eger EI, Saidman LJ, Brandstater: Minimum alveolar anesthetic concentration: A standard of anesthetic potency. *ANESTHESIOLOGY* 26:756-763, 1965
2. Eger EI: MAC, Anesthetic Uptake and Action. Edited by Eger EI, Baltimore, Williams and Wilkins, 1974, pp 1-25
3. Shingu K, Eger EI, Johnson BH, Lurz FW, Hickey RF: MAC values of thiopental and fentanyl in rats. *Anesth Analg* 62:151-154, 1983
4. Bailey PL, Port JD, McJames S, Reinersman L, Stanley TH: Is fentanyl an anesthetic in the dog? *Anesth Analg* 66:542-548, 1987
5. Murphy MR, Hug CC Jr: The anesthetic potency of fentanyl in

- terms of its reduction of enflurane MAC. *ANESTHESIOLOGY* 57:485-488, 1982
6. Yaster M, Koehler RC, Traystman RJ: Effects of fentanyl on peripheral and cerebral hemodynamics in neonatal lambs. *ANESTHESIOLOGY* 66:524-530, 1987
7. Yaster M, Koehler RC, Traystman RJ: Interaction of fentanyl and pentobarbital on peripheral and cerebral hemodynamics in newborn lambs. *ANESTHESIOLOGY* 70:461-469, 1989
8. Steen PA, Michenfelder JD: Cerebral protection with barbiturates, relation to anesthetic effect. *Stroke* 9:140-142, 1971
9. Wang KC: Narcotics are not expected to produce unconsciousness and amnesia. *Anesth Analg* 62:625-626, 1983
10. Lowenstein E, Philbin DM: Narcotic "anesthesia" in the eighties. *ANESTHESIOLOGY* 55:195-197, 1981
11. Lunn JK, Stanley TH, Eisele J, Webster L, Woodward A: High dose fentanyl anesthesia for coronary artery surgery: Plasma fentanyl concentrations and influence of nitrous oxide on cardiovascular responses. *Anesth Analg* 58:390-395, 1979
12. Stanley TH, Webster LR: Anesthetic requirements and cardiovascular effects of fentanyl-oxygen and fentanyl-diazepam-oxygen anesthesia in man. *Anesth Analg* 57:411-416, 1978

(Accepted for publication December 29, 1989.)

Anesthesiology
72:769, 1990

Correction

To the Editor:—We failed to state in our article¹ that Ovassapian *et al.*² proposed that glycine may produce visual disturbance. It was our oversight, and there was no intention to not give credit where credit is due.

K. C. WONG, M.D., PH.D.
*Professor of Anesthesiology and Chairman,
Department of Anesthesiology*

JANICE MEI-LI WANG, M.D.
*Anesthesia Research Fellow, Department of
Anesthesiology*

DONNELL J. CREEL, PH.D.
Research Professor of Ophthalmology

*The University of Utah
50 North Medical Drive
Salt Lake City, Utah 84132*

REFERENCES

1. Wang JM-L, Creel DJ, Wong KC: Transurethral resection of the prostate, serum glycine levels, and ocular evoked potentials. *ANESTHESIOLOGY* 70:36-41, 1989
2. Ovassapian A, Joshi CW, Brunner, EA: Visual disturbances: An unusual symptom of transurethral prostatic resection reaction. *ANESTHESIOLOGY* 57:332-334, 1982

(Accepted for publication January 16, 1990.)

Anesthesiology
72:769-771, 1990

Cardiovascular Responses to Noxious Stimuli in Experimental Animals:
"Pressor or Depressor"?

To the Editor:—We read with interest the recent reports by Gibbs *et al.* of cardiovascular reflex responses to noxious stimuli in rats.^{1,2} In their study, noxious stimuli to the rat's tail were applied during halothane anesthesia by clamping with a rubber-shod clamp, a commonly

used and "standard" method of evaluating anesthetic potency for animal experiments.³ In rats, increasing depth of halothane anesthesia reversed the cardiovascular responses to the noxious stimulation from the pressor (hypertensive) to the depressor (hypotensive) responses.