

TITLE: COMPARISON OF TWO SEDATIVE-ANALGESIC TECHNIQUES FOR LITHOTRIPSY UNDER MAC

AUTHORS: TG Monk, MD, B Bouré, MD, PF White, PhD, MD
AFFILIATION: Department of Anesthesiology, Washington University, St. Louis, MO 63110

Immersion extracorporeal shock wave lithotripsy (ESWL) is widely used to treat urinary tract calculi. Although these procedures are usually performed under general or epidural anesthesia, we designed a study to compare the intraoperative effects and recovery profiles of two sedative-analgesic techniques for immersion ESWL.

44 consenting ASA I-III adult outpatients scheduled to undergo immersion ESWL with the Dornier-HM3 lithotripter were randomly assigned to one of two treatment groups according to an IRB-approved protocol. The first group (FP, n=21) received fentanyl, 2-3 µg/kg iv, followed by propofol, 0.75 mg/kg iv and a maintenance propofol infusion, 50 µg/kg/min. The second group (MA, n=23) received midazolam, 0.05-0.1 mg/kg iv, followed by alfentanil, 10 µg/kg iv, and a maintenance alfentanil infusion, 1.0 µg/kg/min. The infusion rates were varied to achieve adequate sedation and analgesia. Mean arterial pressure, heart rate, respiratory rate (RR), and O₂ saturation (SaO₂), were measured at predetermined time intervals. Postoperatively, the adequacy of the technique was evaluated by the anesthesiologist, urologist, and patient. Side effects and overall satisfaction with the sedation technique were assessed with a follow-up questionnaire. During the same time period, 29 patients who received epidural anesthesia for similar ESWL procedures were compared with the two sedation groups. Data were analyzed using ANOVA and Chi-square tests,

with p<0.05 considered significant.


The three groups were comparable with respect to demographic data and calculi fragmentation. Both FP and MA groups had similar anesthesia and recovery times (table 1). These times were significantly shorter compared to the epidural group. Hemodynamic parameters were comparable between FP and MA, however, both RR and SaO₂ were significantly lower in the MA group (fig. 1 and 2). Both groups were associated with high patient satisfaction and minimal postoperative complications.

In conclusion, the use of a sedation-analgesic technique is an acceptable alternative to epidural anesthesia for immersion lithotripsy. The fentanyl-propofol technique produced less respiratory depression than midazolam-alfentanil during ESWL.

Table 1: Duration of perioperative events (min)*

	FP	MA	EPIDURAL
ANESTHESIA TIME	65±20 †	56±20 †	105±35
PHASE I RECOVERY	43±16 †	52±27 †	109±43
DISCHARGE TIME	147±50 †	143±56 †	199±64

*Mean ± S.D. †p<0.05 vs epidural group



A388

TITLE: MIDAZOLAM PHARMACOKINETICS IN CHILDREN AFTER INTRANASAL ADMINISTRATION

AUTHORS: J.M. Malinovsky, M.D., Y. Le Normand*, M.D., C. de Dieuleveult, M.D., J.Y. Lepage, M.D., A. Cozian, M.D.

AFFILIATION: Dépt. d'Anesth., *Lab. Pharmacol., CHU, 44035 NANTES, France.

Midazolam (M) is often used as premedication. Recently, its intranasal (IN) administration¹ was tested in preschool children for premedication. The aim of the present study was to measure pharmacokinetic parameters of IN M.

Following a pilot study¹ that showed the efficient minimal dose was 0.2 mg.kg⁻¹, 23 healthy children (ASA 1), aged 2-9 yr, weighing 10-30 kg, were included in the study after informed consent of parents and institutional approval. Children, hospitalized for minor urological surgery, were randomly assigned to 2 groups: group IN (n=13) received IN 0.2 mg.kg⁻¹ M; group IV (n=10) received IV 0.2 mg.kg⁻¹ M. None of the children have rhinopharyngitis. Fourteen blood samples were collected before and at 2, 4, 6, 8, 10, 15, 20, 30, 45, 60, 120, 180, 240 and 360 min after M administration. Plasma concentration were measured by GLC with electron-capture detection² and analyzed using a triexponential model. Bioavailability was calculated from the measurement of area under the concentration-time curve (AUC_{IN}/AUC_{IV}). Statistics comparisons were carried out by contingency table.

Results (mean ± SD) are listed in table. No intranasal lesions and disturbances were observed postoperatively.

Important interindividual differences were observed in kinetic parameters, especially after IN administration. By this route, measured availability was 0.32. Reduced IN availability led to significant lower plasma peak, then to lower AUC_{0-∞}. Plasma peak was obtained after a significant longer time than IV administration. In this study, M showed a satisfactory sedative effect, appearing within 5 to 10 min: this seems to be related to plasma levels that reached hypnotic threshold (100 ng.ml⁻¹) within 6 min in all patients. Because it is efficient and not invasive, IN administration of M may be considered as a good route for premedication in children.

Groups	C _{max} ng.ml ⁻¹	T _{max} min	T _{1/2β} min	Vd _{ss} l.kg ⁻¹	Cl ml.min ⁻¹ .kg ⁻¹	AUC _{0-∞} ng.ml ⁻¹ .min
IN	182 ^{¶¶} ±60	13.0 ^{¶¶} ±6.26	88.6 ±54.7	0.704 ±0.481	76 ±53	14580 ^{¶¶} ±4846
IV	1477 ±1214	2 ±0	48.5 ±24.9	0.477 ±0.542	84 ±73	40010 ±28683

Table 1: Pharmacokinetic of M. ¶ p<0.05, ¶¶ p<0.01 vs IV

References

- 1- ANESTHESIOLOGY 69:972-975, 1988.
- 2- ANESTHESIOLOGY 65:536-538, 1986.