

TITLE SUPEROXIDE PRODUCTION BY STIMULATED ENDOTHELIAL CELLS EXPOSED TO VARYING HALOTHANE CONCENTRATIONS

AUTHORS SH Noorily, M.D., JR Shayevitz, M.D., J Varani, M.S., Ph.D.

AFFILIATION Department of Anesthesiology, University of Michigan Medical School, Ann Arbor, MI 48109.


Vascular endothelium produces superoxide anion (O_2^-) when exposed to soluble or particulate stimuli. Halothane (Hal) has been known to inhibit O_2^- production by stimulated neutrophils.¹ In contrast, volatile anesthetics have been found to enhance oxidant-induced endothelial injury.² Given this finding, and the assumption that endothelial cell production of O_2^- is related to injury, we expected to find that Hal enhanced O_2^- production.

Rat pulmonary artery endothelial cells (RPAECs) were isolated as described elsewhere,³ and cultured into monolayers on microcarriers (μ) in Minimal Essential Media containing 10% fetal calf serum. Ten million RPAECs added to two million μ took 5–6 days to grow to confluence. The final culture contained 160 million cells. At confluence, the μ were washed in Hanks Balanced Salt Solution, 0.2% BSA, and aliquoted to test tubes so that each contained 750,000 cells. Tubes were equilibrated with either carrier gas (5% CO_2 in air) or Hal (0.5, 2, or 3 MAC) for 10 minutes. Half of each group of cells were stimulated with phorbol myristate acetate (PMA), 1 μ g/cc. All tubes were then incubated for 1 hour at 37°C. O_2^- production was measured by reduction of ferricytochrome C, as described elsewhere, using an extinction coefficient of 18.5/cm \cdot mM.⁴

Values for O_2^- production were analyzed by 2-way ANOVA. If the F-ratio was significant ($p < 0.05$), between-group comparisons were made

using 1-way ANOVA and Dunnett's test. The O_2^- content in the medium was decreased significantly for PMA-stimulated cells exposed to Hal 2 and 3 MAC. Values shown in the figure represent mean \pm s.e. for 12 experiments.

In conclusion, Hal was found to inhibit the release of O_2^- into the medium by PMA-stimulated cells. This could be a result of inhibited O_2^- release or production. Possible mechanisms for inhibited O_2^- production are inhibition of the xanthine dehydrogenase to xanthine oxidase conversion or inhibition of purine metabolism.



¹Anesthesiology 64:4-12, 1986. ²Anesthesiology 71:A212, 1989.
³J Tissue Cult Methods 10:9-13, 1986. ⁴J Clin Invest 52:741-745, 1973.

A404

TITLE: EFFECTS OF HALOTHANE ON PHOSPHOLIPID N-METHYLATION IN RAT BRAIN SYNAPTOSOMES

Authors: JJ Franks, MD, BVR Sastry, Ph.D., JR Kambam, MD, MJ Surber, BS, R England, BS

Affiliation: Depts. of Anesthesiology and Pharmacology, Vanderbilt University, Nashville, TN 37232

INTRODUCTION: The mechanism of action of inhalational anesthetics is unknown, but neuronal membrane alteration is a favored hypothesis. Axelrod, Hirata and others¹ have demonstrated that enzymatic conversion (methylation) and translocation of inner membranal phosphatidylethanolamine (PE) to outer membranal phosphatidylcholine (PC) facilitates transduction of receptor-mediated signals through cell membranes. We recently have shown that PE methylation is doubled in brain synaptosomes taken from rats anesthetized with 1.4% halothane, returning to normal in rats allowed to recover.² We report here the effect of in vitro exposure of isolated synaptosomes to varying concentrations of halothane.

METHODS: Animal use was approved by the Animal Care Committee of Vanderbilt University. Male Sprague-Dawley rats weighing 285 to 460g were used. Synaptosomes were isolated from brain homogenates by differential centrifugation, as described by Cotman.³ Methylation was measured by the incorporation of tritiated methyl groups from S-adenosyl-L-[³H-methyl]methionine (SAM) into PE.⁴ The incubation mixture, consisting of 0.2mg of synaptosomal protein, buffers and 2 μ M ³H-SAM, was exposed to varying concentrations of halothane for 30 min in a Dubnoff shaker. (Delivered halothane concentrations were confirmed by gas chromatography.) The methylated phospholipids were extracted with chloroform:methanol:HCl (2:1:0.02, v/v) and separated by thin layer chromatography. The activity of PE-N-methyltransferase, the rate limiting enzyme in transmethylation, was expressed by the amount

of phosphatidyl-N-methylethanolamine (PME) formed in fmol/mg protein/30 min.

RESULTS AND DISCUSSION: Halothane in concentrations of 1.0% and 1.4% increased PME ($P < 0.01$) formation from 495 ± 27 fmols (control, $N=8$) to 1594 ± 107 fmols ($N=6$) and 1562 ± 68 fmols ($N=7$), respectively. Thus, halothane at 1.0-1.4% produced a three-fold increase in phospholipid methylation. Halothane concentration of 0.5% did not increase PME formation. PME formation was 1006 ± 47 fmols ($N=7$) and 821 ± 57 fmols ($N=7$), at halothane concentrations of 1.9 and 2.4%, respectively. These high concentrations of halothane caused significant but smaller increases in PME formations. These observations indicate that halothane exhibits a biphasic effect on PME formation. Halothane concentrations higher than 1.4% do seem to retard PME formation. Supported by the Study Center for Anesthesia Toxicology.

REFERENCES:

- 1 Science 209: 1082-1090, 1980.
- 2 FASEB J 4(4):A1007, 1990.
- 3 Methods in Enzymology, Vol. 31, pp 445-452, 1974.
- 4 J Neurochem 34: 1491-1498, 1980.