

## Differential Effect of Oncotic Pressure on Cerebral and Extracerebral Water Content during Cardiopulmonary Bypass in Rabbits

Bradley J. Hindman, M.D.,\* Naohiko Funatsu, M.D.,† Davy C. H. Cheng, M.D.,‡ Roy Bolles, C.C.P.,§ Michael M. Todd, M.D.,¶ John H. Tinker, M.D.\*\*

To study the effect of oncotic pressure on brain water content during cardiopulmonary bypass (CPB), 14 anesthetized New Zealand White rabbits underwent 60 min of nonpulsatile CPB at normothermia. Animals were grouped according to the composition of the circuit priming fluid. Group 1 animals ( $n = 7$ ) received a priming fluid (6.5% hydroxyethyl starch in 0.72 N NaCl;  $323 \pm 13$  mOsm/kg [mean  $\pm$  SD]) that maintained normal colloid oncotic pressure (COP) during CPB ( $19.0 \pm 1.5$  mmHg). Group 2 animals ( $n = 7$ ) received a priming fluid (0.9 N NaCl;  $324 \pm 23$  mOsm/kg) that led to a hyponcotic state (COP =  $6.2 \pm 1.2$  mmHg). Blood chemistries and hemodynamics were recorded every 15 min during CPB. Animals were given additional priming fluid and sodium bicarbonate during CPB to maintain a circuit flow of  $85 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  and arterial pH greater than 7.35. There were no significant differences between groups 1 and 2 with respect to temperature, central venous pressure, mean arterial pressure,  $\text{PaO}_2$ ,  $\text{PaCO}_2$ , plasma sodium concentration, or osmolality at any time during CPB, although osmolality increased in both groups. After 60 min of bypass, animals were killed and organ water contents were determined by wet/dry weight ratios. A separate group of nine similarly prepared and anesthetized animals that did not undergo cannulation or CPB also underwent measurement of plasma chemistries and tissue water contents and served as nonbypass controls (group 3). Brain and kidney water contents were unaffected by oncotic pressure, whereas duodenum and skeletal muscle had significantly greater water content ( $P = 0.003$  and  $P = 0.008$ , respectively) after hyponcotic CPB. To maintain flow and pH, group 2 (hyponcotic) animals required an

average of  $313 \pm 82$  ml additional fluid and  $14 \pm 7$  mEq bicarbonate, whereas group 1 (isooncotic) animals required only  $21 \pm 27$  ml additional fluid ( $P = 0.0001$ ) and  $3 \pm 4$  mEq bicarbonate ( $P = 0.0025$ ). All tissue water contents were identical between the control (group 3) and group 1 animals. These results indicate that the mechanisms that maintain brain fluid balance remain intact during nonpulsatile hyponcotic CPB. Fluid and bicarbonate requirements, and edema formation in other tissue beds can be minimized by maintenance of normal oncotic pressure during CPB. (Key words: Brain; edema. Cardiopulmonary bypass; complications. Fluids: colloid oncotic pressure; osmolality.)

SEVERAL STUDIES HAVE SHOWN that total body interstitial water is increased 15–30% for 24–48 h following cardiopulmonary bypass (CPB), whereas intracellular water content is unchanged.<sup>1–3</sup> Whether particular vital organs are more or less susceptible to interstitial fluid accumulation (edema formation) during bypass has received little attention, although the lung<sup>4</sup> and heart<sup>5</sup> have been the organs of greatest interest. The effects of CPB upon central nervous system (CNS) volume regulation and brain water content have not been studied extensively.

Cardiopulmonary bypass differs from normal circulation in several ways that may predispose to tissue edema. Hemodilution during CPB, usually achieved with hyponcotic priming solutions, decreases colloid oncotic pressure (COP) at least 25–50%.<sup>2,4,6,7</sup> This should theoretically result in increased fluid transfer into the interstitial space. Some data also suggest capillary permeability is increased during CPB.<sup>8</sup> If so, interstitial edema might occur during CPB even if hydrostatic and oncotic forces are maintained at normal values. Finally, early animal studies indicate that nonpulsatile flow impairs lymphatic function.<sup>9</sup> If so, decreased lymphatic clearance of interstitial water would augment the accumulation of edema fluid regardless of the mechanism of formation. Because cerebral edema may compromise cerebral blood flow, particularly in regions of cerebral ischemia or infarction,<sup>10,11</sup> these characteristics of CPB may contribute to the appearance of neurologic deficits following cardiac surgery.

With these considerations in mind, we posed the following questions regarding CNS volume regulation during CPB: 1) Does nonpulsatile CPB result in brain water

\* Associate in Anesthesia, Department of Anesthesia, University of Iowa Hospitals and Clinics.

† Research Fellow, Department of Anesthesia, University of Iowa Hospitals and Clinics.

‡ Research Fellow, Department of Anesthesia, University of Iowa Hospitals and Clinics. Current address: Department of Anaesthesia, University of Toronto, Toronto Western Hospital, Toronto, Ontario.

§ Chief Perfusionist, Division of Cardiothoracic Surgery, Department of Surgery, University of Iowa Hospitals and Clinics.

¶ Professor, Department of Anesthesia, University of Iowa Hospitals and Clinics.

\*\* Professor and Head, Department of Anesthesia, University of Iowa Hospitals and Clinics.

Received from the Cardiovascular Anesthesia Research Laboratory, Department of Anesthesia, College of Medicine, University of Iowa, Iowa City, Iowa. Accepted for publication June 4, 1990. Supported in part by Biomedical Research Support Grant RR 05372 from the Biomedical Research Support Branch, Division of Research Facilities and Resources, National Institutes of Health, to the University of Iowa College of Medicine.

Address reprint requests to Dr. Hindman: Department of Anesthesia, University of Iowa Hospitals and Clinics, Iowa City, Iowa 52242.

accumulation when hydrostatic, osmotic, and oncotic forces are maintained at approximately normal nonbypass values? and 2) Does nonpulsatile CPB predispose the brain to fluid accumulation in hyponcotic states?

## Materials and Methods

### BASIC PREPARATION

The experimental protocol was approved by the Animal Care Committee of the University of Iowa School of Medicine. Anesthesia was induced with 5% halothane in O<sub>2</sub> in 14 New Zealand White rabbits (weight, 4.5 ± 0.5 kg [mean ± SD]) in a plastic box. After cannulation of a marginal ear vein and tracheal intubation with a 3.0-mm ID cuffed tube (Mallinckrodt, Glens Falls, NY), the animals were paralyzed with 1 mg/kg succinylcholine and their lungs ventilated to achieve normocarbica (guided initially by capnography and later by arterial blood gas analysis) using a gas mixture containing 1.5% halothane in 33% O<sub>2</sub>/balance N<sub>2</sub>O. Paralysis was maintained with succinylcholine added to a maintenance lactated Ringer's solution (4 ml · kg<sup>-1</sup> · h<sup>-1</sup>) in amounts sufficient to ensure delivery of 3 mg · kg<sup>-1</sup> · h<sup>-1</sup>. Esophageal temperature was kept at 37–39° C with a servocontrolled heating pad. *Via* a groin incision, catheters were inserted into the abdominal aorta and inferior vena cava to allow continuous monitoring of arterial and central venous pressures and intermittent blood sampling. A 3-ml arterial sample was then obtained for measurement of baseline blood gases, pH, hematocrit, plasma Na<sup>+</sup> concentration (in duplicate *via* flame photometry), osmolality (freezing point depression, Osmette A®, Precision Systems Inc., Sudbury, MA), and oncotic pressure (Wescor 4400, SS-30 membrane, Logan, UT).

### STERNOTOMY AND CANNULATION

After initial preparation, a median sternotomy was performed, and the thymus and pericardium were reflected. Halothane, N<sub>2</sub>O, and maintenance fluids were then discontinued, and each animal received a loading dose of fentanyl and diazepam, followed by a continuous infusion of each for the remainder of the experiment (fentanyl: 100 µg/kg loading dose, 2.5 µg · kg<sup>-1</sup> · min<sup>-1</sup> infusion; diazepam: 2 mg/kg loading dose, 50 µg · kg<sup>-1</sup> · min<sup>-1</sup> infusion). Pancuronium (0.1 mg/kg) was used for subsequent muscle relaxation. The animal's blood was anticoagulated with 300 U/kg of heparin, achieving an activated clotting time greater than 900 s. An 18-French venous catheter (Polystan, Ballerup, Denmark) was placed through a purse-string suture in the right atrium, and a 10 cm long 10-G catheter (Deseret, Sandy, UT) was similarly placed in the ascending aorta (distal orifice ≈ 3 mm above the aortic valve) *via* the left ven-

tricular apex. Subsequent studies using radioactive microspheres have shown equal hemispheric perfusion with this system.

### CARDIOPULMONARY BYPASS

The CPB circuit consisted of a model 540® centrifugal blood pump with a BP-50® pump head (Biomedicus, Eden Prairie, MN), a Capiiox II 08® membrane oxygenator/heat exchanger (Terumo Corporation, Piscataway, NJ), and a heater/cooler (VWR Scientific, San Francisco, CA). Priming volume of the circuit was approximately 200 ml. (See below for details regarding priming fluids.)

Cardiopulmonary bypass was initiated at a flow rate of 85 ml · kg<sup>-1</sup> · min<sup>-1</sup> (reported previously to provide adequate tissue oxygen delivery in rabbits undergoing CPB<sup>12</sup>) and monitored with a calibrated in-line electromagnetic flow meter (Biomedicus® TX40P). The oxygenator was ventilated with a variable mixture of oxygen and nitrogen to achieve PaO<sub>2</sub> of near 250 mmHg and PaCO<sub>2</sub> near 40 mmHg. Because this perfusion system lacks an in-circuit venous reservoir, the animal acted as its own venous reservoir. When venous return decreased, the negative pressure created by the centrifugal pump caused the atrial walls to collapse upon the venous cannula, creating an intermittent, "staccato" venous return that limited bypass flow.<sup>13</sup> In each experimental group, the appearance of this sign was used as the indication for administration of additional priming fluid to maintain the desired flow rate of 85 ml · kg<sup>-1</sup> · min<sup>-1</sup>. At 15-min intervals, 3 ml arterial blood was withdrawn from the animal for measurement of blood gases, pH, hematocrit, plasma Na<sup>+</sup> concentration, and osmotic and oncotic pressure. Sodium bicarbonate was administered to maintain pH > 7.35.

### BYPASS FLUID GROUPS

Bypass animals were assigned to one of two groups based on the composition of the pump priming fluid. In group 1 animals (n = 7), the priming fluid consisted of 350 ml 6.5% hydroxyethyl starch (E. I. DuPont, Bannockburn IL) in 0.72 N sodium chloride to which was added 250 mg CaCl<sub>2</sub> and 1000 U heparin. In addition, 100–150 ml of filtered fresh-packed donor rabbit red blood cells (collected in citrate/phosphate/dextrose) were added, depending upon the volume available from the donor rabbit. In group 2 animals (n = 7), the priming fluid consisted of 350 ml 0.9 N sodium chloride with additional CaCl<sub>2</sub>, heparin, and rabbit red blood cells as per group 1. Prime pH was adjusted to approximately 7.4 with sodium bicarbonate, and the prime composition was measured (hematocrit, sodium, osmolality, and oncotic pressure) before bypass (table 1). Priming fluid in excess of that required for circuit priming was used for supple-

TABLE 1. Priming Fluid Composition

Variable	Group 1 Hydroxyethylstarch (n = 7)	Group 2 Saline (n = 7)
Oncotic pressure (mmHg)	20.7 ± 0.9	0.2 ± 0.2†
Osmolality (mOsm/kg)	323 ± 13	324 ± 23
Sodium* (mmol/l)	137 ± 7	160 ± 4‡
Hematocrit (%)	11 ± 2	12 ± 3
pH	7.32 ± 0.11	7.35 ± 0.04

Mean ± SD.

\* n = 4 for group 1; n = 6 for group 2. Samples not included had been accidentally discarded prior to analysis.

† Significantly different from Group 1, P = 0.0001.

‡ Significantly different from Group 1, P = 0.0007.

mental fluid administration during CPB as described above.

### TISSUE WATER CONTENT

After 60 min of bypass, the animals were killed by discontinuation of CPB and intracardiac administration of saturated KCl solution. One- to 2-g portions of the following structures were rapidly dissected and placed in dry, preweighed vials to determine wet weight: right and left cerebral hemispheres, cervical spinal cord, kidney, skeletal muscle (masseter), and duodenum. Tissue samples were then dried at 80° C over 3–5 days to constant weight.

Organ water content (per cent) was calculated as the difference between wet and dry weights divided by wet weight.

### NONBYPASS CONTROLS (GROUP 3)

To permit evaluation of the effects of CPB *per se*, plasma chemistries and tissue water contents were also measured in a group of nine rabbits studied in a separate protocol that did not involve either cardiac cannulation or CPB. Because the basic preparation, anesthetic and hemodynamic profile, and total experimental duration of 3–4 hours was equivalent to bypass animals, we have included their time-related results as nonbypass controls.

### STATISTICS

Tissue water content in the three groups was examined using a one-way analysis of variance (ANOVA). Hemodynamic and blood data were examined by two-way ANOVA (bypass group, time on CPB), with time on CPB treated as the repeated variable. Significance was assumed for P < 0.05. All results are expressed as mean ± SD.

### Results

Physiologic data from the bypass animals are shown in table 2. There were no significant differences between groups 1 and 2 with respect to temperature (38.2 ± 1.3 vs. 37.0 ± 1.0° C), central venous pressure (4 ± 3 vs. 5

TABLE 2. Blood and Hemodynamic Data of Groups 1 and 2 before and at Intervals during Cardiopulmonary Bypass

Variable	Group	Baseline	Duration of CPB (min)			
			15	30	45	60
Oncotic pressure (mmHg)	1 (n = 7)	18.7 ± 1.3	18.8 ± 1.6	19.2 ± 1.5	18.9 ± 1.7	18.9 ± 1.6
	2 (n = 7)	19.2 ± 1.5	6.1 ± 1.2	6.3 ± 1.1	6.4 ± 1.2	6.2 ± 1.4
Osmolality* (mOsm/kg)	1	304 ± 6	321 ± 10	327 ± 11	330 ± 11	335 ± 8
	2	311 ± 12	330 ± 11	334 ± 13	336 ± 11	336 ± 14
Sodium (mmol/l)**	1	138 ± 7	137 ± 6	137 ± 6	137 ± 7	137 ± 7
	2	136 ± 7	142 ± 7	140 ± 6	138 ± 6	144 ± 6
Hematocrit† (%)	1	41 ± 6	23 ± 2	21 ± 2	21 ± 2	21 ± 2
	2	43 ± 6	29 ± 3	28 ± 5	29 ± 4	27 ± 4
pH‡	1	7.44 ± .04	7.45 ± .06	7.39 ± .02	7.33 ± .05	7.32 ± .08
	2	7.45 ± .02	7.34 ± .07	7.32 ± .04	7.32 ± .05	7.34 ± .04
Mean arterial pressure (mmHg)	1	83 ± 18	67 ± 13	70 ± 12	71 ± 10	68 ± 10
	2	84 ± 19	60 ± 15	60 ± 13	63 ± 13	64 ± 12
Pump flow (ml · kg <sup>-1</sup> · min <sup>-1</sup> )	1	—	88 ± 3	87 ± 5	90 ± 4	90 ± 3
	2	—	86 ± 4	86 ± 4	87 ± 4	91 ± 5
Fluid required (ml)	1	—	—	—	—	21 ± 27
	2	—	—	—	—	313 ± 82§
Bicarbonate required (mEq)	1	—	—	—	—	3 ± 4
	2	—	—	—	—	14 ± 7¶

Mean ± SD.

CPB = cardiopulmonary bypass.

\* Osmolality increased significantly in both groups during CPB, P = 0.008.

† Groups 1 and 2 significantly different during CPB, P = 0.01.

‡ Group 2 significantly more acidotic than group 1 during CPB, P = 0.05

§ Significantly different from group 1, P = 0.0001.

¶ Significantly different from group 1, P = 0.0025.

\*\* n = 4 group 1, n = 6 group 2. Samples not included had been accidentally discarded prior to analysis.

$\pm 2$  mmHg),  $\text{PaO}_2$  ( $297 \pm 127$  vs.  $256 \pm 155$  mmHg), or  $\text{PaCO}_2$  ( $36 \pm 6$  vs.  $40 \pm 7$  mmHg) during CPB, respectively. Total flow was maintained at approximately  $88 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  and arterial  $\text{pH}$  at 7.35 for 60 min in each bypass group, although group 2 animals tended to be more acidotic during the early phase of CPB ( $P = 0.05$ ). Group 2 (hyponcotic) animals had numerically lower average mean arterial pressures at equivalent bypass flow rates than group 1 (isocotc) animals, but group means were not significantly different. To maintain flow and  $\text{pH}$ , group 2 animals required an average of  $313 \pm 82$  ml additional fluid and  $14 \pm 7$  mEq bicarbonate. In contrast, group 1 animals required only  $21 \pm 27$  ml additional fluid ( $P = 0.0001$ ) and  $3 \pm 4$  mEq bicarbonate ( $P = 0.0025$ ). Despite equivalent values for baseline (pre-CPB) and priming hematocrit, and much greater volume requirements, hyponcotic group 2 animals had significantly higher hematocrits during CPB than isocotc group 1 animals ( $P = 0.01$ ).

Oncotic pressure was stable during CPB, averaging  $19.0 \pm 1.5$  mmHg in group 1 animals and  $6.2 \pm 1.2$  mmHg in group 2 ( $P = 0.0001$ ). Osmolality was greater during CPB in groups 1 and 2 compared to prebypass baseline and increased over time ( $P = 0.008$ ). However, there was no significant difference in osmolality or plasma sodium concentration between groups 1 and 2 at any time. Control animals (group 3) had an average oncotic pressure of  $18.8 \pm 1.5$  mmHg, with osmolality increasing from  $304 \pm 10$  to  $318 \pm 13$  mOsm/kg over the 3-h fentanyl/diazepam infusion period ( $P = 0.025$ , paired  $t$  test).

Despite the marked differences in volume requirement and hematocrit, there were no significant differences in brain, spinal cord, or kidney water content among the three groups. In contrast, tissue water content was significantly greater in skeletal muscle (masseter;  $P = 0.008$ ) and duodenum ( $P = 0.003$ ) after 60 min of CPB in hyponcotic animals (group 2) when compared to isocotc bypass and nonbypass control groups (table 3). There were no differences in tissue water content in any organ between the isocotc bypass group (group 1) and nonbypass controls (group 3).

## Discussion

The few studies examining brain edema following bypass have yielded conflicting results. In 1958, Halley and co-workers measured brain volume using a volumetric displacement technique in dogs after CPB.<sup>14</sup> Thirteen of 27 dogs were found to have increased brain volume. Unfortunately, measurement of brain volume is not an accurate indicator of water accumulation because swelling may occur simply with vascular congestion and enlargement of the intravascular compartment. Obstruction of cerebral venous return can produce such a situation; in fact, caval obstruction from the venous cannula was noted in 31% of the affected animals. In contrast, Utley, using wet/dry weight ratios, found no increase in brain water content following CPB in either hemodiluted or nonhemodiluted animals.<sup>15</sup> Unfortunately, in neither study were hemodynamic, osmotic, or oncotic parameters quantified. Thus, the current report appears to be the first to examine the acute effect of nonpulsatile CPB and altered oncotic pressure on brain water content. Our results clearly show brain water content was not different from nonbypass controls following 60 min of CPB, *irrespective of COP*. In contrast, other tissue beds (skeletal muscle and duodenum) had significantly greater water contents following hyponcotic CPB.

We believe these findings are consistent with other experimental data and with known differences in structure between brain and peripheral capillaries. In the brain, endothelial tight junctions render the capillary bed remarkably impermeable to both ions and large molecules (colloids), although not to water or to lipid-soluble substances. Net water movement is largely dependent on changes in total osmotic pressure.<sup>16</sup> Because colloid osmotic pressure contributes, at most, the equivalent of 0.5–1 mOsm/kg to total osmolality,<sup>16</sup> even large changes in COP have a negligible effect on total osmolality. Thus, blood–brain osmotic gradients, rather than oncotic gradients, determine net water movement.<sup>16–18</sup> Peripheral capillaries differ from those in the brain in that they *are* permeable to ions but remain relatively impermeable to

TABLE 3. Tissue Water Content after 60 minutes of Cardiopulmonary Bypass

Group	Tissue Water Content (%)					
	Right Hemisphere	Left Hemisphere	Spinal Cord	Kidney	Duodenum	Skeletal Muscle
1 (n = 7)	78.4 $\pm$ 0.5	78.5 $\pm$ 0.4	67.9 $\pm$ 1.3	80.6 $\pm$ 0.6	76.5 $\pm$ 3.4	74.9 $\pm$ 1.4
2 (n = 7)	78.6 $\pm$ 0.4	78.5 $\pm$ 0.8	68.6 $\pm$ 0.9	80.6 $\pm$ 2.0	82.3 $\pm$ 3.3*	77.6 $\pm$ 1.2†
3 (n = 9)	78.4 $\pm$ 0.6	78.4 $\pm$ 0.6	68.0 $\pm$ 0.5	80.9 $\pm$ 1.9	75.5 $\pm$ 3.7	75.4 $\pm$ 1.8

Mean  $\pm$  SD.

Group 2 significantly different than groups 1 and 3 (one-way ANOVA): \*  $P = 0.003$ ; †  $P = 0.008$ .

oncologically active molecules. Under these circumstances, intravascular-extravascular/interstitial osmotic gradients are difficult to establish and maintain. As a consequence, oncotic and hydrostatic gradients become important determinants of water movement, as described by the Starling equation. One would therefore predict that changes in oncotic pressure might have a major influence on the formation of edema in peripheral tissue but not in the brain. Recently, using nonbypass models, Zornow *et al.*<sup>18</sup> and Kaieda *et al.*<sup>19</sup> found brain water content to be unaffected by changes in plasma oncotic pressure in both normal rabbits and those subjected to cryogenic brain injury. Our data indicate nonpulsatile CPB does not alter these relationships.

In this experiment, as in others,<sup>15,19,20</sup> skeletal muscle and the gastrointestinal tract (small bowel) appeared to be quite susceptible to edema formation in hyponcotic states. Oncotic pressure also influenced bicarbonate and fluid requirements during CPB, with hyponcotic animals requiring nearly 15 times the volume of fluid to maintain bypass flow and five times the amount of bicarbonate to maintain arterial pH. Despite these greater volume requirements, hyponcotic animals still had significantly greater hematocrits during bypass than did isoncotic animals. These findings are consistent with a marked translocation of fluid from the intravascular space, impaired tissue perfusion, and possibly, increased metabolic acid (lactic) production in hyponcotic animals. Our finding of unaltered tissue water content following isoncotic CPB suggests either: 1) nonpulsatile CPB does not result in measurable alteration of capillary permeability; or 2) lymphatic function was not significantly impaired and was therefore able to compensate for any increase in transcapillary fluid movement into the interstitium. This latter hypothesis is supported by findings of increased lymphatic flow during CPB in animals in both iso- and hyponcotic states.<sup>8,21</sup>

Although oncotic pressure remained stable during CPB, our unexpected finding was that osmolality increased during CPB in both groups by 25–30 mOsm/kg. Group 2 animals required, on average, 11 mEq more bicarbonate than those in group 1, which might be expected to create an osmolar load (22 mOsm), leading to “dehydration” of the brain. During CPB, the volume of extravascular fluid is increased by the volume of the CPB circuit, approximately 200 ml. If extracellular volume is 25–30% of body weight, this indicates the total volume of extracellular fluid during bypass was  $0.3 \times 4,500 + 0.200 = 1.5$  l. The additional bicarbonate given to group 2 animals in this volume would be expected to increase plasma sodium by  $\approx 8$  mmol/l, which was the observed increase, and intravascular osmolality by  $\approx 15$  mOsm/kg, which is less than the observed increase. Consequently, the increase

in osmolality during CPB cannot be completely ascribed to increased plasma sodium concentration. Glucose and lactate concentrations were not measured, but studies performed recently with this preparation revealed lactate accumulation (5–10 mmol/l) and hyperglycemia (7–14 mmol/l) during 60 min of CPB. Such increases may partially explain the increase in osmolality observed during CPB. Thus, as shown in Table 2, despite differences in bicarbonate requirement, there was no significant difference in plasma osmolality or sodium concentration at any time point between groups 1 and 2. Therefore, the absence of a difference in brain water content in the presence of low oncotic pressure cannot be ascribed to differences in osmolality.

Osmolality also increased significantly in nonbypass group 3 animals, and although well matched to group 1 with respect to oncotic pressure, there was a significant difference in the final value of measured osmolality ( $335 \pm 8$  vs.  $318 \pm 13$  mOsm/kg;  $P = 0.01$ ). Thus, it is at first surprising that there was no difference in brain water content between these groups when brain water content is so highly dependent on osmotic gradients.<sup>16–18</sup> However, for a substance to be osmotically active, a concentration gradient must exist across an impermeable barrier. Osmolality, measured *via* freezing point depression, depends upon the total number of solute molecules, regardless of their permeability characteristics. Because brain water content was equivalent in all groups, it would appear the observed increase in osmolality in both bypass and nonbypass animals was at least partially due to substance(s) that rapidly equilibrate across the blood–brain barrier (possibly propylene glycol,<sup>16</sup> the primary vehicle of the diazepam formulation used in these experiments, or glucose) such that osmotic gradients were not established.

We chose to study rabbits undergoing CPB for several reasons. First, rabbits are of sufficient size that a commercially available pediatric perfusion apparatus can be employed. Second, unlike dogs and cats, which possess rete mirabilia resulting in extensive collateralization between internal and external carotid systems, the rabbit brain is supplied exclusively by the internal carotid and vertebral arteries in a pattern similar to humans.<sup>22</sup> Although subhuman primates have comparable cerebrovascular anatomy, the low cost and ease of care of rabbits compared to primates make them an attractive experimental animal. Third, reproducible models of focal cerebral infarction, both microscopic (embolic)<sup>23,24</sup> and macroscopic (vessel occlusion),<sup>25,26</sup> have been developed for the rabbit such that future studies concerning the effect of CPB on outcome from neurologic injury should be possible. Fourth, the cerebrovascular responses to changes in  $P_{aCO_2}$ <sup>27,28</sup> and arterial pressure<sup>29</sup> are well

characterized in the rabbit and are virtually identical to human values. Resting cerebral blood flow<sup>25,27</sup> and cerebral metabolic rate for oxygen<sup>28,30</sup> also approximate human values.

The anesthetic chosen was based on the desire to approximate anesthetics used clinically for patients undergoing CPB. Based on clinical,<sup>31-33</sup> electroencephalographic,<sup>34,35</sup> pharmacokinetic,<sup>36,37</sup> and pharmacodynamic<sup>38</sup> data for fentanyl and diazepam in the rabbit, we devised a constant infusion technique that would be anticipated to maintain plasma concentrations of at least two times the ED<sub>95</sub> concentration of each agent. Studies in unparalyzed animals in our laboratory have shown that this regimen completely eliminated all somatic and hemodynamic responses to clamping of the hindpaw web space over a 3-h period without evidence of acute tolerance or tachyphylaxis.

In summary, in rabbits undergoing nonpulsatile CPB, we found that brain water content was unaffected by marked reductions in oncotic pressure produced by a crystalloid priming solution. Hypooncotic animals had exaggerated fluid and bicarbonate requirements during CPB and significant increases in muscle and duodenal water contents as compared to isoncotic animals. Nonpulsatile CPB *per se*, irrespective of oncotic pressure, did not appear to result in brain edema, nor edema in other organs, when oncotic pressure was maintained at normal levels.

---

†† Bergman SA, Wynn RL, Williams G: Diazepam enhances fentanyl and diminishes meperidine antinociception. *Anesth Prog* 35: 190-194, 1988.

The authors wish to thank E. I. DuPont De Nemours and Company for kindly providing Hetastarch® powder, and Terumo Corporation for generously providing many of the oxygenators used in these experiments.

### References

- Brans YW, Dweck HS, Harris HB, Parr GVS, Bailey PE, Kirklin JW, Cassady G: Effect of open-heart surgery on the body composition of infants and young children. *Pediatr Res* 15:1024-1028, 1981
- Beattie HW, Evans G, Garnett ES, Webber CE: Sustained hypovolemia and extracellular fluid volume expansion following cardiopulmonary bypass. *Surgery* 71:891-897, 1972
- Pacifico AD, Digerness S, Kirklin JW: Acute alterations of body composition after open intracardiac operations. *Circulation* 41: 331-341, 1970
- Byrick RJ, Kay JC, Noble WH: Extravascular lung water accumulation in patients following coronary artery surgery. *Can Anaesth Soc J* 24:332-345, 1977
- Laks H, Standeven J, Blair O, Hahn J, Jellinek M, Willman VL: The effects of cardiopulmonary bypass with crystalloid and colloid hemodilution on myocardial extravascular water. *J Thorac Cardiovasc Surg* 73:129-138, 1977
- English TAH, Digerness S, Kirklin JW: Changes in colloid osmotic pressure during and shortly after open intracardiac operation. *J Thorac Cardiovasc Surg* 61:338-341, 1971
- Sade RM, Stroud MR, Crawford FA, Kratz JM, Dearing JP, Bartles DM: A prospective randomized study of hydroxyethyl starch, albumin, and lactated Ringer's solution as priming fluid for cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 89:713-722, 1985
- Baue AE, Nusbaum M, Anstadt G, Blakemore WS: The pattern of lymphatic flow during extracorporeal circulation. *J Thorac Cardiovasc Surg* 50:648-657, 1965
- Parsons RJ, McMaster PD: The effect of the pulse upon the formation and flow of lymph. *J Exp Med* 68:353-376, 1938
- Iannotti F, Hoff JT, Schielke GP: Brain tissue pressure in focal cerebral ischemia. *J Neurosurg* 62:83-89, 1985
- Meyer FB, Sundt TM, Yanagihara T, Anderson RE: Focal cerebral ischemia: Pathophysiologic mechanisms and rationale for future avenues of treatment. *Mayo Clin Proc* 62:35-55, 1987
- Wabeke E, Elstrodt JM, Mook PH, Gathier S, Wildevuur RH: Clear prime for infant cardiopulmonary bypass: A miniaturized circuit. *J Cardiovasc Surg* 29:117-122, 1988
- Wenger RK, Bavaria JE, Ratcliffe MB, Bogen D, Edmunds LH: Flow dynamics of peripheral venous catheters during extracorporeal membrane oxygenation with a centrifugal pump. *J Thorac Cardiovasc Surg* 96:478-484, 1988
- Halley MM, Reemtsma K, Creech O: Cerebral blood flow, metabolism, and brain volume in extracorporeal circulation. *J Thorac Surg* 36:506-518, 1958
- Utley JR, Todd EP, Wachtel CC, Cain RB, Collins J: Effect of hypothermia, hemodilution, and pump oxygenation on organ water content and blood flow. *Surg Forum* 27:217-219, 1976
- Fenstermacher JD: Volume regulation of the central nervous system, Edema. Edited by Staub NC, Taylor AE. New York: Raven Press, 1984, pp 383-404
- Todd MM, Tommasino C, Moore S: Cerebral effects of isovolemic hemodilution with a hypertonic saline solution. *J Neurosurg* 63:944-948, 1985
- Zornow MH, Todd MM, Moore SS: The acute cerebral effects of changes in plasma osmolality and oncotic pressure. *ANESTHESIOLOGY* 67:936-941, 1987
- Kaieda R, Todd MM, Warner DS: Prolonged reduction in colloid oncotic pressure does not increase brain edema following cryogenic injury in rabbits. *ANESTHESIOLOGY* 71:554-560, 1989
- Rosenkranz ER, Utley JR, Menninger FJ, Dembitsky WP, Hargens AR, Peters RM: Interstitial fluid pressure changes during cardiopulmonary bypass. *Ann Thorac Surg* 30:536-542, 1980
- Anabtawi IN, Womack CE, Ellison RG: Thoracic duct lymph flow during pulsatile and nonpulsatile extracorporeal circulation. *Ann Thorac Surg* 2:38-43, 1966
- Scremin OU, Sonnenschein RR, Rubinstein EH: Cerebrovascular anatomy and blood flow measurements in the rabbit. *J Cereb Blood Flow Metab* 2:55-66, 1982
- Lyden PD, Zivin JA, Kochhar A, Mazzarella V: Effects of calcium channel blockers on neurologic outcome after focal ischemia in rabbits. *Stroke* 19:1020-1026, 1988
- Zivin JA, Lyden PD, DeGirolami U, Kochhar A, Mazzarella V, Hemenway CC, Johnston P: Tissue plasminogen activator. Reduction of neurologic damage after experimental embolic stroke. *Arch Neurol* 45:387-391, 1988
- Meyer FB, Anderson RE, Sundt TM, Yaksh TL: Intracellular brain pH, indicator tissue perfusion, electroencephalography, and histology in severe and moderate focal cortical ischemia in the rabbit. *J Cereb Blood Flow Metab* 6:71-78, 1986
- Yamamoto K, Yoshimine T, Yanagihara T: Cerebral ischemia in

- rabbit: A new experimental model with immunohistochemical investigation. *J Cereb Blood Flow Metab* 5:529-536, 1985
27. Orr JA, DeSoignie RC, Wagerle LC, Fraser DB: Regional cerebral blood flow during hypercapnia in the anesthetized rabbit. *Stroke* 14:802-807, 1983
  28. Pearce WJ, Scremin OU, Sonnenschein RR, Rubinstein EH: The electroencephalogram, blood flow, and oxygen uptake in rabbit cerebrum. *J Cereb Blood Flow Metab* 1:419-428, 1981
  29. Tuor UI, Farrar JK: Pial vessel caliber and cerebral blood flow during hemorrhage and hypercapnia in the rabbit. *Am J Physiol* 247:H40-H51, 1984
  30. Heistad DD, Marcus ML, Gourley JK, Busija DW: Effect of adenosine and dipyridamole on cerebral blood flow. *Am J Physiol* 240:H775-H780, 1981
  31. Flecknell PA, John M, Mitchell M, Shurey C, Simpkin S: Neuroleptanalgesia in the rabbit. *Lab Anim* 17:104-109, 1983
  32. Flecknell PA, Mitchell M: Midazolam and fentanyl-fluanisone: Assessment of anaesthetic effects in laboratory rodents and rabbits. *Lab Anim* 18:143-146, 1984
  33. Mero M, Makela A, Vainionpaa S, Vihtonen K, Rokkanen P: The use of neuroleptanaesthesia for experimental orthopaedic surgery in the rabbit. *Acta Vet Scand* 28:251-252, 1987
  34. Kubicki S, Freund G, Henschel FW, Schoppenhorst M: Fentanyl und sulfentanil im elektroenzephalographischen vergleich. *Anaesthesist* 26:333-342, 1977
  35. Yamamoto J: Characteristics of the cortical and hippocampal EEG power spectra of rabbits during normal behavioral states and after administration of CNS acting drugs. *Jpn J Pharmacol* 37: 227-234, 1985
  36. Rigg JRA, Wong TY, Horsewood P, Hewson JR: Steady-state plasma fentanyl in the rabbit. *Br J Anaesth* 53:1337-1345, 1983
  37. Klotz U, Antonin KH, Bieck PR: Pharmacokinetics and plasma binding of diazepam in man, dog, rabbit, guinea pig and rat. *J Pharmacol Exp Ther* 199:67-73, 1976
  38. Hess R, Herz A, Friedel K: Pharmacokinetics of fentanyl in rabbits in view of the importance for limiting the effect. *J Pharmacol Exp Ther* 179:474-484, 1971