

Anesthesiology
76:150, 1992

In Reply:—We appreciate the interest of Rooke regarding our work¹ on myocardial oxygen demand indices. Of course regression analysis of pooled data may lead to a result different than that using regression analysis performed separately for each patient. However, the difference in results cannot be predicted because of the intersubject variability. Although in Rooke and Feigl's dog studies² the use of pooled values *versus* individual regression analysis resulted in an underestimation of the relative value of external work term, with different data an overestimation might have resulted. We would agree that an individual regression analysis for each patient would be interesting; however, due to the nature of our data such an approach would seem not to be valid. It is not possible under clinical conditions to perform abundant coronary blood flow measurements in patients. Therefore, in most patients only three measurements were made, and in some of the patients only two measurements were possible. Thus, an individual regression analysis does not make much sense in our data.

We appreciate the careful reading of Rooke and agree that the coefficient for the external work term in PWI modified for clinical use (PWI_{mod}) should be 8×10^{-3} and not 8×10^{-6} when cardiac index is expressed in liters per squared meter. The calculations were performed with cardiac index expressed in milliliters per squared meter, which explains the difference.

Anesthesiology
76:150, 1992

An Armored Laryngeal Mask Airway

To the Editor:—Wilson and Eastley¹ have been forward-thinking in their use of an armored Mallinckrodt tube to reinforce a size-2 laryngeal mask airway. We are at present using a prototype armored laryngeal mask airway in the Cambridge Day Surgery Unit (fig. 1).

A. P. L. GOODWIN, F.C.ANAES.
Napp Research Fellow

T. W. OGG, M. A., F.C.ANAES.
*Consultant Anaesthetist
Director of Day Surgery*

*Day Surgery Unit
Addenbrooke's Hospital
Hills Road
Cambridge CB2 2QQ
United Kingdom*

REFERENCE

1. Wilson IG, Eastley R: A modification of the laryngeal mask airway. ANESTHESIOLOGY 74:1157, 1991

(Accepted for publication September 20, 1991.)

ANDREAS HOEFT, M.D.
*Visiting Assistant Professor
Department of Anesthesiology
University of Texas
Health Science Center at Houston
6341 Fannin, 5.020 MSMB
Houston, Texas 77030*

REFERENCES

1. Hoeft A, Sonntag H, Stephan H, Kettler D: Validation of myocardial demand indices in patients awake and during anesthesia. ANESTHESIOLOGY 75:49–56, 1991
2. Rooke GA, Feigl EO: Work as a correlate of canine left ventricular oxygen consumption, and the problem of catecholamine oxygen wasting. Circ Res 50:273–286, 1982

(Accepted for publication September 18, 1991.)




FIG. 1. The armored laryngeal mask airway (Intavent, UK).