

Anesthesiology
79:313-322, 1993
© 1993 American Society of Anesthesiologists, Inc.
J. B. Lippincott Company, Philadelphia

Mechanism of the Direct, Negative Inotropic Effect of Ketamine in Isolated Ferret and Frog Ventricular Myocardium

Suneerat Kongsayreepong, M.D.,* David J. Cook, M.D.,† Philippe R. Housmans, M.D., Ph.D.‡

Background: Ketamine exerts both an indirect, positive inotropic effect and a direct, negative inotropic effect in isolated ferret ventricular myocardium. This negative inotropic effect becomes apparent after inactivation of the sympathetic neuroeffector junction. The aim of this study was to investigate the mechanisms of ketamine's intrinsic negative inotropic effect.

Methods: The authors analyzed the effects of ketamine after β -adrenoceptor blockade on variables of contractility and relaxation, and on the free intracellular Ca^{++} transient detected with the Ca^{++} -regulated photoprotein aequorin. Ketamine's effects were also evaluated in a preparation in which the sarcoplasmic reticulum (SR) function was impaired by ryanodine, and in frog ventricular myocardium in which the SR is poorly developed.

Results: Ketamine at concentrations $\geq 3.3 \times 10^{-5}$ M decreased contractility and the amplitude of the intracellular Ca^{++} transient. After inactivation of sarcoplasmic reticulum Ca^{++} release with 10^{-6} M ryanodine, a condition in which myofibrillar activation depends almost exclusively on transsarcolemmal Ca^{++} influx, ketamine caused a decrease in contractility and in the amplitude of the intracellular Ca^{++} transient, and ketamine's relative negative inotropic effect was not different from that in control muscles not exposed to ryanodine. Furthermore, $\geq 10^{-4}$ M ketamine decreased contractility in frog ventricular myocardium, a species that is almost entirely dependent on transsarcolemmal Ca^{++} influx for its myofibrillar activation.

Conclusions: These findings indicate that the direct negative inotropic effect of ketamine results from a decrease in intracellular Ca^{++} availability with no changes in myofibrillar Ca^{++} sensitivity. At least part of ketamine's action is caused by inhibition of transsarcolemmal Ca^{++} influx. (Key words: Aequorin. Anesthetics, intravenous: ketamine. Heart: contractility; intracellular Ca^{++} transient.)

* Resident, Department of Anesthesiology, Mayo Foundation.

† Instructor in Anesthesiology, Mayo Medical School.

‡ Associate Professor of Anesthesiology and Pharmacology, Mayo Medical School.

Received from the Departments of Anesthesiology and Pharmacology, Mayo Clinic and Mayo Foundation, Rochester, Minnesota. Accepted for publication March 31, 1993. Supported by the United States Public Health Service Grant GM 36365 and by Mayo Foundation.

Address reprint requests to Dr. Housmans: Department of Anesthesiology, Mayo Clinic, 200 First Street SW, Rochester, Minnesota 55905.

quorin. Anesthetics, intravenous: ketamine. Heart: contractility; intracellular Ca^{++} transient.)

KETAMINE has useful cardiostimulatory effects¹ that result mostly from central² and peripheral³ stimulation of the sympathetic nervous system. Results of two previous studies^{4,5} on isolated ferret papillary muscle indicated that the positive inotropic effect of ketamine results from the inhibition of neuronal and extraneuronal catecholamine uptake. In muscles from animals treated with reserpine or after β -adrenoceptor blockade,⁴ ketamine did not exert a positive inotropic effect, but demonstrated a negative inotropic effect at concentrations equal to or greater than 10^{-4} M. The aim of this study was to examine the mechanism(s) of the direct inotropic effects of ketamine after β -adrenoceptor blockade. The results indicate that ketamine decreases intracellular $[\text{Ca}^{++}]$, does not change myofibrillar Ca^{++} sensitivity, and decreases net transsarcolemmal Ca^{++} influx.

Materials and Methods

This study was approved by the Animal Care and Use Committee of the Mayo Foundation. We used papillary muscles from the right ventricle of adult male ferrets (weight 1,100–1,500 g, age 16–19 weeks). The animals were anesthetized with sodium pentobarbital (100 mg/kg, intraperitoneally), and the heart was quickly removed through a left thoracotomy. The right ventricle was opened, and suitable papillary muscles were excised and mounted in a temperature-controlled (30° C) muscle chamber that contained a physiologic salt solution of the following composition (mM): Na^{+} 135; K^{+} 5; Ca^{++} 2.25; Mg^{2+} 1; Cl^{-} 103.5; HCO_3^{-} 24; HPO_4^{2-} 1; SO_4^{2-} 1; acetate 20; and glucose 10. This solution was equilibrated with 95% O_2 and 5% CO_2 (pH = 7.4). Suitable preparations were selected on the basis of previously used criteria.⁴ The muscles were

held between the lever of a force-length transducer (Innovi, Antwerp, Belgium) and a miniature Lucite® clip with a built-in stimulation electrode. Muscles were stimulated at a stimulus frequency of 0.25 Hz, with rectangular pulses of 5-ms duration and an intensity 10% above threshold. Muscles were made to contract in alternating series of four isometric and four isotonic twitches at preload only during a 2-h period of stabilization before the onset of the experiment. All experiments were carried out with the initial muscle length set at L_{\max} , *i.e.*, the muscle length at which active force development is maximal. All ferret papillary muscles were pretreated with 10^{-7} M (\pm)-bupranolol HCl⁶ before the onset of the experiment to abolish the indirect effects of ketamine.^{4,5}

Contractile Variables

Isometric and isotonic twitches against the L_{\max} preload only were recorded during steady state at each drug concentration.⁴ Peak developed force (DF), time to peak force (TPF), and time to 50% isometric relaxation measured from time to peak force (RTH) were measured from isometric twitches. Peak isotonic shortening (DL) was measured from isotonic twitches at the preload of L_{\max} . Maximal unloaded velocity of shortening (MUVS) was measured from "zero-load-clamped" twitches, *i.e.*, an isotonic twitch at L_{\max} in which load is rapidly (<3 ms) decreased electronically to zero load during the latent period.

Detection of the Intracellular Ca^{++} Transient

In experiments in which the intracellular Ca^{++} transient was measured, muscles were prepared as follows. Electrical stimulation was stopped and multiple superficial cells were microinjected with the Ca^{++} -regulated photoprotein aequorin,⁷ to allow for subsequent detection of the intracellular Ca^{++} transient. It was usually necessary to microinject 30–100 cells. After microinjection, muscles were allowed to rest for 1 h, and were then carefully transferred to a vertical muscle chamber that allowed for simultaneous detection of variables of contractility and of aequorin luminescence.^{8,9} Muscles were made to contract isometrically at L_{\max} throughout experiments in which aequorin luminescence was measured. It was usually necessary to average luminescence and force signals of 16–256 twitches to obtain a satisfactory signal-to-noise ratio in aequorin luminescence signals. This was accomplished on a digital oscilloscope (Nicolet 4094B, Madison, WI). We quantified diastolic aequorin luminescence, peak

systolic aequorin luminescence, and time to peak aequorin luminescence. The time from the stimulus to the time when aequorin luminescence had decreased to 25% of its peak value (during the decline of the aequorin signal) ($t_{1.25}$) was taken as a measure of the time course of the decline of the aequorin signal.

Experimental Design

Four protocols were used to examine the mechanism of ketamine's intrinsic, direct inotropic effect; each muscle served as its own control.

In group 1 muscles ($n = 10$), we determined possible changes in the intracellular Ca^{++} transient during the direct, intrinsic effect of ketamine. After β -adrenoceptor blockade with 10^{-7} M (\pm)-bupranolol HCl, a dose-response curve to ketamine was obtained in each of ten muscles (group 1). The following steps were used: control; 10^{-6} M, 10^{-5} , 10^{-4} , and 3.3×10^{-4} M ketamine; and washout. The muscles were exposed to each concentration of ketamine until a steady state of at least 5 min was achieved before contractile response was recorded. To determine whether muscle performance and aequorin luminescence decayed during the duration of the experiment, ketamine was washed out and control measurements were repeated at the end of each experiment.

Total serum concentrations of ketamine were $60 \mu\text{M}$ 5 min after intravenous administration of 2 mg/kg ketamine, and reached steady levels of 7–10 μM in patients.¹⁰ Because ketamine's plasma protein binding is 12%,¹¹ one can calculate peak free plasma ketamine concentrations of 5.3×10^{-5} M and sustained free plasma concentrations of 6.2 to 8.8×10^{-6} M. These calculations indicate that we studied ketamine concentrations (10^{-6} M to 3.3×10^{-4} M) that encompass the clinically relevant free (unbound) drug concentrations in plasma.

In group 2 muscles ($n = 9$), we determined whether ketamine possibly alters myocardial relaxation or myofibrillar Ca^{++} sensitivity. Each of nine muscles were pretreated with 10^{-7} M (\pm)-bupranolol, and subjected to "Ca⁺⁺-back titration" experiments: after measurement of control variables of the isometric twitch, muscles were exposed to 3.3×10^{-5} M ketamine. Extracellular $[Ca^{++}]$ was then rapidly raised by adding small aliquots of a concentrated $CaCl_2$ solution (112.5 mM) to the bathing solution, until peak developed force was equal to that in the control twitch. This protocol allowed us to compare relaxation and time variables ($n = 9$) and aequorin luminescence signals ($n = 5$) in

KETAMINE AND MYOCARDIAL CONTRACTILITY

control and in the presence of 3.3×10^{-5} M ketamine at equal peak developed force. Similar "Ca⁺⁺-back titration" experiments were carried out in muscles exposed to 10^{-4} M (n = 5) and 3.3×10^{-4} M (n = 5) ketamine.

In group 3 muscles (n = 11), we attempted to determine whether transsarcolemmal Ca⁺⁺ exchange is affected by ketamine, by excluding the contribution of the sarcoplasmic reticulum to Ca⁺⁺ release by pretreatment with the plant alkaloid ryanodine. Each of 11 muscles was exposed to 10^{-7} M (\pm)-bupranolol and 10^{-6} M ryanodine. The effects of 10^{-6} M, 10^{-5} M, 10^{-4} M, and 3.3×10^{-4} M ketamine and washout on contractile variables (n = 11) and aequorin luminescence (n = 5) were assessed.

In group 4 muscles (n = 8), we assessed the effects of ketamine on frog ventricular myocardium, a species that is primarily dependent on transsarcolemmal Ca⁺⁺ exchange for activation.¹² Ventricular strips were cut from the ventricle of pithed frogs (*Rana pipiens*), and were mounted vertically in the muscle chamber for measurements of contractility variables during ketamine dose-response experiments. The physiological salt solution was diluted to 80% of its original composition with distilled water to approximate the composition of extracellular fluid in frogs.¹³ Frog ventricular strip experiments were carried out at 25°C. In an additional seven frog ventricular strips, we assessed the effects of 10^{-4} M ketamine after exposure to 10^{-6} M ryanodine.

All waveforms of aequorin luminescence, force, length, and velocity were displayed as a function of time on a four-channel digital oscilloscope (Nicolet 4094B), stored permanently on 5¼" floppy disks and recorded at slow speed on a four-channel pen recorder (Honeywell 1400, Denver, CO). All waveforms of interest recorded on the digital oscilloscope were transferred to a computer (Reason Technology 486/33 MHz, Minneapolis, MN), on which variables of contraction and relaxation, aequorin luminescence, and corresponding time values were automatically determined.

Statistical Analysis

The contractile responses to ketamine and peak aequorin luminescence were assessed with repeated measures analysis of variance. When appropriate, Dunnett's test was used to compare effects of individual ketamine concentrations with control. A *P* value < 0.05 was considered significant.

Results

Table 1 shows the values of aequorin luminescence and of measurements of contractility in the ketamine dose-response experiments (group 1). Figures 1 and 2 illustrate that, after β -adrenoceptor blockade, ketamine did not significantly alter DF or aequorin luminescence at concentrations $\leq 10^{-5}$ M. At concentrations $\geq 10^{-4}$ M, ketamine caused a concentration-dependent decrease of both peak developed force and peak aequorin luminescence. Diastolic aequorin luminescence, time to peak aequorin luminescence, time to 25% decline in the peak luminescence signal (t_{L25}), time to peak force (TPF), and time to half isometric relaxation (RTH) were not significantly changed at any ketamine concentration. In 3.3×10^{-4} M ketamine, the peak of aequorin luminescence occurred later (66.6 ± 22.7 ms) than in control (52.1 ± 9.7 ms), and the Ca⁺⁺ transient lasted longer (t_{L25} 183.7 ± 61.8 ms *vs.* 139.4 ± 24.7 ms), yet these changes were not statistically significant. All variables of contractility and aequorin luminescence after washout of ketamine did not differ significantly from control.

To determine whether ketamine alters myofibrillar Ca⁺⁺ responsiveness, aequorin luminescence signals were measured in five muscles (group 2) and were compared at equal peak developed force in control (fig. 3, *left*) and after exposure to 3.3×10^{-5} M ketamine in elevated [Ca⁺⁺]_o (fig. 3, *right*). Table 2 lists the values of aequorin luminescence and of peak developed force in control; in (3.3×10^{-5} M, 10^{-4} M, and 3.3×10^{-4} M) ketamine; and in ketamine in higher [Ca⁺⁺]_o. Figure 4 summarizes the changes in peak aequorin luminescence during the Ca⁺⁺-back titration experiments for each of the three ketamine concentrations tested. When ketamine's negative inotropic effect on DF was corrected by raising [Ca⁺⁺]_o, peak aequorin luminescence was not significantly different from control (*P* > 0.05, n = 5, Student's paired *t* test). In the presence of $\geq 10^{-4}$ M ketamine in higher [Ca⁺⁺]_o at equal peak force as in the control, time to peak aequorin luminescence, TPF, and RTH were significantly decreased (table 2).

To assess the effects of ketamine on contractility independent of the sarcoplasmic reticulum Ca⁺⁺ release, aequorin luminescence (n = 5) and contractility were measured under isometric conditions in ferret papillary muscle after exposure to 10^{-6} M ryanodine (fig. 5). Consistent with its effects on the SR, 10^{-6} M ryanodine decreased developed force and peak aequorin lumi-

Table 1. Aequorin Luminescence and Variables of Contractility during Control Conditions in Ferret Papillary Muscles (Group 1, n = 10; mean ± SD)

	Control	Ketamine 10 ⁻⁶ M	Ketamine 10 ⁻⁵ M	Ketamine 10 ⁻⁴ M	Ketamine 3.3 × 10 ⁻⁴ M	Wash
Diastolic aequorin luminescence (nA)	0.45 ± 0.16	0.49 ± 0.16	0.45 ± 0.17	0.47 ± 0.22	0.50 ± 0.37	0.48 ± 0.30
Peak systolic aequorin luminescence (nA)	1.01 ± 0.76	1.20 ± 0.79	0.96 ± 0.71	0.49 ± 0.36*	0.16 ± 0.11*	1.00 ± 0.76
Time to peak aequorin luminescence (ms)	52.1 ± 9.7	56.8 ± 9.9	53.3 ± 10.9	60.3 ± 15.5	66.6 ± 22.7	53.9 ± 13.0
Peak developed force (mN/mm ²)	20.73 ± 8.12	22.5 ± 7.7	20.6 ± 8.4	14.4 ± 5.1*	6.5 ± 4.3*	23.0 ± 8.9
Time to peak force (ms)	237.6 ± 70.6	247.6 ± 77.4	241.6 ± 76.2	225.5 ± 82.2	229.1 ± 90.1	237.9 ± 73.9
Time to half isometric relaxation (ms)	122.2 ± 30.1	131.8 ± 30.1	125.7 ± 32.2	115.6 ± 28.4	108.4 ± 25.9	119.1 ± 26.4
Time to 25% luminescence (t ₂₅)	139.4 ± 24.7	143.5 ± 23.9	141.6 ± 30.4	152.3 ± 45.3	183.7 ± 61.8	137.5 ± 30.8

* P < 0.01 for comparison with control by repeated-measures analysis of variance and Dunnett's test.

nescence from control conditions (table 3, fig. 5, left). In muscles pretreated with ryanodine 10⁻⁶ M, force and aequorin luminescence were further decreased by ketamine at concentrations ≥ 10⁻⁴ M (fig. 6).

The role of the sarcoplasmic reticulum in ketamine's negative inotropic effect was further evaluated in six additional experiments summarized in figure 7. Contractility under isometric and isotonic conditions was compared in papillary muscles in which SR function was poisoned by ryanodine, and those in which SR function remained intact (historical control reported in Cook *et al.*⁴). Correcting for ryanodine's effect by expressing values as a percent of control reveals a virtually superimposable negative inotropic effect of ketamine in the presence or absence of SR function. The relative effects of ketamine on DF and DL with or without functional sarcoplasmic reticulum did not differ (one-way ANOVA, P > 0.2, n = 6).


In frog ventricular muscle, ketamine had no significant effect on either peak force of isometric twitches (DF) or on peak shortening of isotonic twitches (DL) at concentrations up to 10⁻⁴ M (figs. 8 and 9). Ketamine, at concentrations of 10⁻⁴ M and higher caused a marked negative inotropic effect (figs. 8 and 9). Peak developed force was decreased from a control value of 14.26 ± 7.79 mN · mm⁻² to 11.57 ± 6.77 mN · mm⁻² in 10⁻⁴ M ketamine, and to 7.30 ± 4.82 mN · mm⁻² in 3.3 × 10⁻⁴ M ketamine (all values mean ± SD; n = 8). Ketamine at concentrations of 10⁻⁴ M and 3.3 × 10⁻⁴ M decreased DL from a control value (mean ± SD; n = 8) of 0.125 ± 0.066 L/L_{max} to 0.101 ± 0.061 L/L_{max} and 0.060 ± 0.042 L/L_{max} in 10⁻⁴ M and 3.3 × 10⁻⁴ M ketamine, respectively. To further ascertain that the effects of ketamine on frog ventricular myocardium were independent of any potential effect on the sarcoplasmic reticulum, an additional seven frog ventricular strips were exposed to 10⁻⁶ M ryanodine and then to 10⁻⁴ M ketamine. Ryanodine, 10⁻⁶ M, did not change DF, DL, or MUVS. Subsequent exposure to 10⁻⁴ M ketamine significantly decreased these contractile variables (fig. 10).

Discussion

The inotropic effects of ketamine on mammalian myocardium have been reported to be positive, negative, or biphasic, depending on the species studied and on the experimental conditions. Variances of this kind have been extensively discussed by others.^{4,5,14} Previous studies have demonstrated that, at clinically rel-

KETAMINE AND MYOCARDIAL CONTRACTILITY

Fig. 1. Effects of ketamine on force and aequorin luminescence during isometric twitch contractions after β -adrenoceptor blockade. Each panel shows superimposed traces of force and of aequorin luminescence. One hundred and twenty-eight contractions were averaged.



evant concentrations, ketamine exerts an indirect positive inotropic effect on ferret ventricular myocardium.^{4,5} This was shown to result from inhibition of neuronal catecholamine uptake mechanisms.^{4,5} The overall effect of ketamine on contractility is the net result of this indirect positive inotropic effect and of ketamine's intrinsic negative inotropic effect on ventricular myocardium. The purpose of this study was to investigate the mechanisms of ketamine's direct negative inotropic effect.

This *in vitro* study clearly demonstrates the direct negative inotropic effects of ketamine at concentrations of 3.3×10^{-5} M or greater; this approximates the plasma concentrations of ketamine measured after intravenous induction doses.¹⁰ The negative inotropic effect of ketamine is accompanied by a decrease in peak aequorin luminescence without any change in the kinetics of the aequorin signal. This concomitant decrease in the amplitude of contraction and of the intracellular Ca^{++} transient is most likely caused by a reduced availability of intracellular Ca^{++} . A decreased intracellular Ca^{++} availability could result from decreased Ca^{++} release by the sarcoplasmic reticulum or decreased net transsarcolemmal Ca^{++} entry, or from an enhanced Ca^{++} uptake by the SR. A significant alteration of Ca^{++} sequestration by the SR is unlikely, because the time courses of relaxation and of aequorin luminescence were not affected by ketamine.

To determine possible effects of ketamine on mechanisms other than Ca^{++} release from the SR, we studied the effects of ketamine in myocardium after SR Ca^{++} release was completely abolished by ryanodine pretreatment.^{15,16}

The plant alkaloid ryanodine binds to specific myocardial receptors¹⁷ and reduces the availability of Ca^{++} from the SR for contractile activation.¹⁸ Ryanodine markedly decreases the amount of Ca^{++} released from the SR, and has no effect on the SR Ca^{++} uptake pump,¹⁹ or the $\text{Na}^{+}\text{-Ca}^{++}$ exchanger. The significant reduction in all contractile variables from control measurements after 10^{-6} M ryanodine is similar in magnitude to that reported previously in similar²⁰ and identical²¹ experimental conditions, and reflects the substantial contribution of the SR to the activator Ca^{++} pool in ferret ventricular myocardium. In the presence of 10^{-6} M




Fig. 2. Effects of ketamine on peak developed force (●) and peak aequorin luminescence (○) after β -adrenoceptor blockade. Values in ketamine (mean \pm SD, $n = 10$) are expressed as percent of control. ** $P < 0.01$ versus control.




Fig. 3. Ca^{2+} -back titration experiments (see text for details). Two hundred and fifty-six contractions were averaged. The timing of the stimulus is indicated on the short horizontal bar at the lower left corner of each panel.

ryanodine, ketamine caused a further reduction in contractile variables and of peak aequorin luminescence; this probably reflects a depressant effect of ketamine on an alternative source of activator Ca^{2+} , most likely transsarcolemmal Ca^{2+} influx.

Frog ventricular myocardium has a poorly developed SR, and is almost entirely dependent on transsarcolemmal Ca^{2+} influx to activate the myofibrillar apparatus.¹² Indeed, 10^{-6} M ryanodine failed to change myocardial contractility in frog ventricle, an observation that indicates that the SR in frog ventricle does not regulate contractility. The ketamine-induced decrease in contractility in frog ventricle with or without ryanodine

pretreatment is, therefore, consistent with an effect of ketamine to decrease net transsarcolemmal Ca^{2+} influx.

Our findings are in agreement with those of Rusy *et al.*,²² who attributed the negative inotropic effect of ketamine in isolated rabbit papillary muscle primarily to inhibition of transsarcolemmal Ca^{2+} influx with little or no effect on Ca^{2+} release by the sarcoplasmic reticulum. More recently, Baum and Tecson²³ observed that 10^{-5} M and 10^{-4} M ketamine decreased peak inward calcium current in single enzymatically dissociated guinea pig myocardial cells.

Endou *et al.*²⁴ studied the correlation between inotropic and electrophysiologic effects of ketamine in

Table 2. Aequorin Luminescence and Developed Force in Ferret Papillary Muscle during Ca^{2+} -Back Titration Experiments (Group 2; mean \pm SD)

	Control $[\text{Ca}^{2+}]_o$ = 2.25 mM	Ketamine 3.3×10^{-5} M $[\text{Ca}^{2+}]_o$ = 2.25 mM	Ketamine 3.3×10^{-5} M $[\text{Ca}^{2+}]_o > 2.25$ mM	N
Aequorin luminescence (nA)	0.58 \pm 0.13	0.42 \pm 0.06	0.58 \pm 0.09	5
Time to peak aequorin luminescence (ms)	50.6 \pm 3.7	50.0 \pm 4.0	49.4 \pm 2.9	5
Peak developed force (mN/mm ²)	30.38 \pm 3.13	26.60 \pm 3.88†	30.53 \pm 3.27	9
Time to peak force (ms)	216.9 \pm 8.3	219.6 \pm 14.5†	215.0 \pm 8.3	9
Time to half isometric relaxation (ms)	127.4 \pm 6.6	122.4 \pm 7.7	127.8 \pm 7.2	9
	Control $[\text{Ca}^{2+}]_o$ = 2.25 mM	Ketamine 10^{-4} M $[\text{Ca}^{2+}]_o$ = 2.25 mM	Ketamine 10^{-4} M $[\text{Ca}^{2+}]_o > 2.25$ mM	
Aequorin luminescence (nA)	4.07 \pm 2.08	1.74 \pm 1.32*	3.85 \pm 2.73	5
Time to peak aequorin luminescence (ms)	103.4 \pm 20.2	114.0 \pm 17.5	52.2 \pm 2.7†	5
Peak developed force (mN/mm ²)	31.80 \pm 20.46	23.77 \pm 15.65*	34.55 \pm 23.86	5
Time to peak force (ms)	402.8 \pm 37.8	378.2 \pm 63.7	203.4 \pm 18.6†	5
Time to half isometric relaxation (ms)	230.0 \pm 34.6	197.8 \pm 42.3*	128.8 \pm 32.1*	5
	Control $[\text{Ca}^{2+}]_o$ = 2.25 mM	Ketamine 3.3×10^{-4} M $[\text{Ca}^{2+}]_o$ = 2.25 mM	Ketamine 3.3×10^{-4} M $[\text{Ca}^{2+}]_o > 2.25$ mM	
Aequorin luminescence (nA)	3.21 \pm 1.71	0.61 \pm 0.50*	3.83 \pm 1.24	5
Time to peak aequorin luminescence (ms)	91.6 \pm 13.2	109.8 \pm 28.5	49.0 \pm 3.6†	5
Peak developed force (mN/mm ²)	34.46 \pm 20.47	16.17 \pm 12.83*	39.63 \pm 27.84	5
Time to peak force (ms)	374.4 \pm 25.8	335.8 \pm 96.7	217.2 \pm 25.3‡	5
Time to half isometric relaxation (ms)	204.0 \pm 40.0	168.2 \pm 42.7*	133.4 \pm 26.2†	5

* $P < 0.05$, † $P < 0.01$, ‡ $P < 0.001$, for comparison with control by Student's paired *t* test.

KETAMINE AND MYOCARDIAL CONTRACTILITY




Fig. 4. Summary of Ca^{++} -back titration experiments in 3.3×10^{-5} M (○), 10^{-4} M (●), and 3.3×10^{-4} M (△) ketamine ($n = 5$ each). Peak aequorin luminescence (peak light) is plotted as percent of control (mean \pm SD; $n = 5$ in each group) during exposure to ketamine in $[\text{Ca}^{++}]_o = 2.25$ mM and at $[\text{Ca}^{++}]_o > 2.25$ mM at equal peak force as in the control. * $P < 0.05$ versus control.

atrial and ventricular myocardium of the guinea pig and rat, species in which ketamine exerts, respectively, a negative and a positive inotropic effect. The increase in force after ketamine in rat myocardium was found to result from ketamine's inhibition of the transient outward current (I_{to}) with minimal effect on I_{Ca} , whereas, in guinea pig myocardium, the decrease in I_{Ca} after ketamine was more important than the positive inotropic effect supported by a small decrease in I_{to} . In view of these results, it is not surprising that ketamine exerts a positive inotropic effect in rat myocardium^{24,25} that depends mostly on Ca^{++} release from the sarcoplasmic reticulum for its activation. The results of Endou *et al.*²⁴ are also consistent with ketamine's negative inotropic effect in rabbit²² and ferret myocardium, species in which transsarcolemmal Ca^{++} entry plays a larger role for activation than it does in rat myocardium.

Table 3. Aequorin Luminescence and Developed Force in Ferret Papillary Muscle before and after Ryanodine 10^{-6} M Group ($n = 5$; mean \pm SD)

	Control	Ryanodine 10^{-6} M
Aequorin luminescence (nA)	0.93 ± 0.62	$0.10 \pm 0.07^*$
Time to peak aequorin luminescence (ms)	93.0 ± 96.7	$163.0 \pm 73.3^\dagger$
Developed force (mN/mm ²)	24.25 ± 12.26	$4.30 \pm 2.19^*$
Time to peak force (ms)	252.6 ± 93.8	$303.6 \pm 104.7^*$
Time to half isometric relaxation (ms)	110.4 ± 25.1	99.0 ± 9.6

* $P < 0.05$, $^\dagger P < 0.01$, for comparison with control by Student's paired t test.

To determine possible effects of ketamine on myofibrillar Ca^{++} responsiveness (Ca^{++} sensitivity), Ca^{++} -back titration experiments were designed to produce isometric twitches of equal peak amplitude with and without ketamine, and to compare the $[\text{Ca}^{++}]_i$ transients and mechanical relaxation in those conditions. The assumption implicit to this type of analysis is that the Ca^{++} occupancy of troponin C at peak force is the same in either condition, so that myofibrillar Ca^{++} sensitivity can be assessed from the relationship between $[\text{Ca}^{++}]_i$ and Ca^{++} occupancy of troponin C. If ketamine alters reaction mechanisms "downstream" from the binding of Ca^{++} to troponin C, and modifies the relationship between Ca^{++} occupancy of troponin C and force, our approach would be invalid. Yet, so far, there is no evidence that this occurs. Moreover, it is difficult to determine, in twitch contractions of intact living muscle fibers, whether a particular intervention changes myofibrillar Ca^{++} responsiveness by comparison of force and Ca^{++} transients alone, because the relationship between force and $[\text{Ca}^{++}]_i$ in twitch contractions does not reach steady state. When the kinetics of the $[\text{Ca}^{++}]_i$ tran-




Fig. 5. Effects of ketamine on force and aequorin luminescence after β -adrenoceptor blockade and inactivation of SR Ca^{++} release by 10^{-6} M ryanodine. The left panel shows superimposed traces of force and of aequorin luminescence during isometric twitches before and after 10^{-6} M ryanodine. Note alteration in vertical scales in the post-ryanodine treated figures. One hundred and twenty-eight contractions were averaged in control conditions; 512 twitches were averaged in ryanodine.




Fig. 6. Effects of ketamine on peak developed force and on peak aequorin luminescence after β -adrenoceptor blockade and inactivation of SR Ca^{++} release with 10^{-6} M ryanodine. Values in ketamine (mean \pm SD, n = 5) are expressed as percents of control. * $P < 0.05$; ** $P < 0.01$ versus control.

sient are dramatically altered by the intervention, the changes may be impossible to interpret in terms of changes of myofibrillar Ca^{++} responsiveness.⁸ In our




Fig. 7. Effects of 10^{-4} and 3×10^{-4} M ketamine after β -adrenoceptor blockade on peak developed force (DF) and peak isotonic shortening (DL) in ferret right ventricular papillary muscles (mean \pm SD; n = 9) with an intact SR (upper graph, data from Cook *et al.*⁴) and after inactivation of SR Ca^{++} release (lower graph) in muscles (mean \pm SD; n = 6) exposed to 10^{-6} M ryanodine before the ketamine dose-response experiment. ** $P < 0.01$ compared to respective control.




Fig. 8. Effects of ketamine of force of frog ventricular strips. Force traces of isometric twitch contractions before and during a ketamine dose-response experiment are superimposed.

experiments, because the kinetics of the aequorin luminescence signals were not changed in the Ca^{++} -back titration experiments in 3.3×10^{-5} M ketamine, the conclusions based on our experimental data should be valid for that ketamine concentration.

Because the amplitude and time course of the intracellular Ca^{++} transients at equal peak developed force were not different with or without ketamine, we conclude that ketamine has minimal, if any, effects on myofibrillar responsiveness to Ca^{++} . The absence of




Fig. 9. Effects of ketamine on peak developed force (DF) and peak isotonic shortening (DL) of frog ventricular strips. Values in ketamine (mean \pm SD, n = 8) are expressed as percent of control. ** $P < 0.01$ versus control.

KETAMINE AND MYOCARDIAL CONTRACTILITY




Fig. 10. Effects of 10^{-6} M ryanodine (left) and of 10^{-6} M ryanodine and 10^{-4} M ketamine (right) on amplitude variables of contractility (DF, DL, MUVS) in frog ventricular strips. Data are mean \pm SD ($n = 7$). * $P < 0.05$ versus respective control; # $P < 0.01$ versus respective control.

specific effects of ketamine on isometric relaxation is consistent with this view. However, because $\geq 10^{-4}$ M ketamine in higher $[Ca^{++}]_o$ in the Ca^{++} -back titration experiments abbreviated the time course of both aequorin luminescence and isometric force (table 2), inferences about the possible absence of changes in myofibrillar Ca^{++} sensitivity in $\geq 10^{-4}$ M ketamine are less firm.

The direct negative inotropic effect of ketamine can, therefore, be attributed to an interference with cellular mechanisms that regulate intracellular Ca^{++} availability. The evidence from the current study indicates that, at clinically relevant concentrations, ketamine exerts a mild-to-moderate direct negative inotropic effect by decreasing net transsarcolemmal Ca^{++} influx. The depressant effect may be caused by an inhibition of the sarcolemmal slow inward L-type Ca^{++} current.^{23,24} Possible effects of ketamine on other membrane Ca^{++} exchange mechanisms (Na^+/Ca^{++} exchange, Ca^{++} ATPase export pump, etc.) cannot be excluded, and may necessitate further study.

In conclusion, ketamine exerts both a direct negative inotropic effect and an indirect positive inotropic effect, the latter resulting from inhibition of neuronal and extraneuronal norepinephrine uptake.^{4,5} The net inotropic effect in ferret ventricular myocardium was a complex dose-response curve (positive inotropic effect at low ketamine concentrations, and a smaller positive inotropic effect at high ketamine concentrations).

We speculate that the inotropic impact of ketamine in other species, or in human myocardium, will depend on the state of sympathetic neuroeffector junction,^{4,5} on the relative importance of intracellular Ca^{++} delivery mechanisms (SR Ca^{++} release and transsarcolemmal Ca^{++} influx⁴), and on specific electrophysiologic effects on I_{to} and I_{Ca} .²⁴

References

1. Tweed WA, Minuck M, Mymin D: Circulatory responses to ketamine anesthesia. *ANESTHESIOLOGY* 37:613-619, 1972
2. Ivankovich AD, Miletich DJ, Reimann C, Albrecht RF, Zahed B: Cardiovascular effects of centrally administered ketamine in goats. *Anesth Analg* 53:924-931, 1974
3. Lundy PM, Gverzdys S, Frew R: Ketamine: Evidence of tissue specific inhibition of neuronal and extraneuronal catecholamine uptake processes. *Can J Physiol Pharmacol* 63:298-303, 1985
4. Cook DJ, Carton EG, Housmans PR: Mechanism of the positive inotropic effect of ketamine in isolated ferret ventricular papillary muscle. *ANESTHESIOLOGY* 74:880-888, 1991
5. Cook DJ, Housmans PR, Roric DK: Effect of ketamine HCl on norepinephrine disposition in isolated ferret ventricular myocardium. *J Pharmacol Exp Ther* 261:101-107, 1992
6. Kaumann AJ, McInerney TK, Gilmour DP, Blinks JR: Comparative assessment of β -adrenoceptor blocking agents as simple competitive antagonists in isolated heart muscle. *Naunyn Schmiedebergs Arch Pharmacol* 311:219-236, 1980
7. Blinks JR, Mattingly PH, Jewell BR, van Leeuwen M, Harrer GC, Allen DG: Practical aspects of the use of aequorin as a calcium indicator: Assay, preparation, microinjection, and interpretation of signals. *Methods Enzymol* 57:292-328, 1978
8. Blinks JR, Endoh M: Modification of myofibrillar responsiveness to Ca^{++} as an inotropic mechanism. *Circulation* 73:III-85-III-98, 1986
9. Carton EG, Wanek LA, Housmans PR: Effects of nitrous oxide on contractility, relaxation and the intracellular calcium transient of isolated mammalian ventricular myocardium. *J Pharmacol Exp Ther* 257:843-849, 1991
10. Idvall J, Ahlgren I, Aronsen KF, Stenberg P: Ketamine infusions: Pharmacokinetics and clinical effects. *Br J Anaesth* 51:1167-1173, 1979
11. Wieber J, Gugler R, Hengstmann JH, Dengler HJ: Pharmacokinetics of ketamine in man. *Anaesthetist* 24:260-263, 1975
12. Fabiato A: Calcium-induced release of calcium from the cardiac sarcoplasmic reticulum. *Am J Physiol* 245:C1-C14, 1983
13. Lockwood APM: "Ringer" solutions and some notes on the physiological basis of their ionic composition. *Comp Biochem Physiol* 2:241-289, 1961
14. Pagel PS, Kampine JP, Schmeling WT, Warltier DC: Ketamine depresses myocardial contractility as evaluated by the preload recruitable stroke work relationship in chronically instrumented dogs with autonomic nervous system blockade. *ANESTHESIOLOGY* 76:564-572, 1992
15. Jenden DJ, Fairhurst AS: The pharmacology of ryanodine. *Pharmacol Rev* 21:1-25, 1969
16. Sutko JL, Willerson JT, Templeton GH, Jones LR, Besch HR Jr: Ryanodine: Its alterations of cat papillary muscle contractile

state and responsiveness to inotropic interventions and a suggested mechanism of action. *J Pharmacol Exp Ther* 209:37-47, 1979

17. Inui M, Saito A, Fleischer S: Isolation of the ryanodine receptor from cardiac sarcoplasmic reticulum and identity with feet structures. *J Biol Chem* 262:15637-15642, 1987

18. Inui M, Wang S, Saito A, Fleischer S: Characterization of junctional and longitudinal sarcoplasmic reticulum from heart muscle. *J Biol Chem* 263:10843-10850, 1988

19. Feher JJ, Lipford GB: Mechanism of action of ryanodine on cardiac sarcoplasmic reticulum. *Biochim Biophys Acta* 813:77-86, 1985

20. Marban E, Wier WG: Ryanodine as a tool to determine the contributions of calcium entry and calcium release to the calcium transient and contraction of cardiac Purkinje fibers. *Circ Res* 56:133-138, 1985

21. Carton EG, Housmans PR: Role of transsarcolemmal Ca^{2+} entry in the negative inotropic effect of nitrous oxide in isolated ferret myocardium. *Anesth Analg* 74:575-579, 1992

22. Rusy BF, Amuzu JK, Bosscher HA, Redon D, Komai H: Negative inotropic effect of ketamine in rabbit ventricular muscle. *Anesth Analg* 71:275-278, 1990

23. Baum VC, Tecson ME: Ketamine inhibits transsarcolemmal calcium entry in guinea pig myocardium: Direct evidence by single cell voltage clamp. *Anesth Analg* 73:804-807, 1991

24. Endou M, Hattori Y, Nakaya H, Gotoh Y, Kanno M: Electrophysiologic mechanisms responsible for inotropic responses to ketamine in guinea pig and rat myocardium. *ANESTHESIOLOGY* 76:409-418, 1992

25. Riou B, Lecarpentier Y, Viars P: Inotropic effect of ketamine on rat cardiac papillary muscle. *ANESTHESIOLOGY* 71:116-125, 1989