

CORRESPONDENCE

Anesthesiology
81:264, 1994
© 1994 American Society of Anesthesiologists, Inc.
J. B. Lippincott Company, Philadelphia

Postanesthetic Apnea in Full-term Infants

To the Editor:—Recently Andropoulos *et al.*¹ described four full-term infants who had postoperative apnea after pyloromyotomy. The cases are well presented, and the discussion of the possible mechanisms for the apnea in these infants is comprehensive.

As correctly indicated by the authors, the possibility of apnea immediately after pyloromyotomy in otherwise healthy infants has been described previously. This series of four infants is a good addition to the existing body of knowledge. The whole report, however, must be examined in the proper perspective before generalized recommendations are made.

The authors correctly acknowledge that, in the absence of an apnea-bradycardia monitor, the duration of apnea in patients 1, 2, and 4 may have been overestimated and actually may have been within what would be considered normal limits for this postconceptional age. Patient 3 had significant intraoperative respiratory events, including hemoglobin oxygen desaturation and possible bronchospasm requiring metaproterenol inhalation. Patient 4 had documented preoperative apnea and hypochloremic metabolic alkalosis.

Some aspects of the anesthetic management of these infants require closer examination. These infants underwent a rapid-sequence induction of anesthesia to protect against the risk of aspiration. Although the case report does not indicate the dose of thiopental used in each patient, it generally is acknowledged that infants in this age group have greater thiopental requirements than do older children.² Thiopental, however, has a markedly prolonged serum half-life in newborns compared with adults.³ It was shown recently that the incidence of respiratory complications in otherwise healthy infants 1–6 months of age is greater after thiopental induction when compared with other agents, such as halothane or propofol.⁴

This case report adds to our knowledge of the effect of general anesthesia on the incidence and severity of apnea in infants undergoing pyloromyotomy. To extrapolate the findings in these special patients to all other, presumably healthy, full-term infants undergoing routine elective surgery under an anesthetic technique that does not include barbiturates or opioids is not warranted. The conclusion that "*consideration should be given to short-term apnea monitoring in young full-term infants who undergo general anesthesia*" is

not supported by the data in this report. This recommendation would add unnecessary expense, liability, and anxiety to those caring for these infants and to their families. Until further investigations substantiate the authors' belief, this recommendation should be rescinded.

Raafat S. Hannallah, M.D.
Professor of Anesthesiology and Pediatrics

Leila G. Welborn, M.D.
Professor of Anesthesiology and Pediatrics

Willis A. McGill, M.D.
Professor of Anesthesiology and Pediatrics
Chairman, Department of Anesthesiology

Children's National Medical Center
111 Michigan Avenue NW
Washington, D.C. 20010

References

1. Andropoulos DB, Heard MB, Johnson KL, Clarke JT, Rowe RW: Postanesthetic apnea in full-term infants after pyloromyotomy. *ANESTHESIOLOGY* 80:216–219, 1994
2. Jonmarker C, Westrin P, Larsson S, Werner O: Thiopental requirements for induction of anesthesia in children. *ANESTHESIOLOGY* 67:104–107, 1987
3. Coté CJ: Practical pharmacology of anesthetic agents, narcotics, and sedatives, *A Practice of Anesthesia for Infants and Children*. 2nd edition. Edited by Coté CJ, Ryan JF, Todres ID, Goudsouzian NG. Philadelphia, WB Saunders, 1993, p 108
4. Schrum SF, Hannallah RS, Verghese PM, Welborn LG, Norden JM, Ruttiman U: Comparison of propofol and thiopental for rapid anesthesia induction in infants. *Anesth Analg* 78:482–485, 1994

(Accepted for publication April 4, 1994.)

Anesthesiology
81:264–265, 1994
© 1994 American Society of Anesthesiologists, Inc.
J. B. Lippincott Company, Philadelphia

In Reply:—Hannallah *et al.* have raised several issues concerning our case report that deserve comment. The dose of thiopental used for rapid-sequence induction of anesthesia in our patients ranged from 5.1 to 6.4 mg/kg. It is acknowledged that the serum half-life of thiopental in newborns may be prolonged considerably and that it is possible that this may have contributed to the central apnea seen in our patients within 1 h of the conclusion of surgery.

As to the contention that our conclusion is not supported by the data in the case report, it must be emphasized that the four patients in our report, combined with the previous four case reports, as well as knowledge of the development of respiratory drive in young infants, form the basis for this recommendation. Our four patients were inpatients, mildly to moderately dehydrated on admission. Of the four other reports of apnea in full-term infants, three were scheduled for

CORRESPONDENCE

outpatient elective surgery and were otherwise completely healthy. The anesthetic technique in these infants consisted of an inhalation induction with oxygen, nitrous oxide, and halothane. None of the three received a barbiturate or an opioid.¹⁻³

As mentioned in our discussion,⁴ respiratory drive, even in the full-term infant, undergoes a maturational process, with normal "adult" breathing patterns and carbon dioxide response curves not achieved until 42-43 weeks postconceptional age. Before this age, "abnormalities" in respiratory drive are of similar type but lesser in magnitude and frequency when compared with the premature infant of a younger postconceptional age.^{5,6} Postanesthetic apnea in the former preterm infant is well documented, and most pediatric anesthesiologists admit such patients for 12-24 h postoperatively for apnea monitoring up to a postconceptional age varying between 46 and 60 weeks.^{7,8} As Hannallah *et al.* are aware, attention to the problem of postanesthetic apnea in expremature infants was focused originally by case reports⁹ and a retrospective study,¹⁰ and similar recommendations for inpatient admission and careful postoperative monitoring to ours were made at the time, before careful prospective studies were performed. Then, when careful studies using polysomnography by Welborn *et al.*⁷ and Kurth *et al.*⁸ were published, recommendations based on more objective data could be made. It is clear from those careful studies that many of the apneic spells were clinically unrecognized, detected only by the apnea monitor.

Although we do not think that postanesthetic apnea in full-term infants is nearly as frequent as in the expremature infant, nevertheless, the problem exists. Clearly, a large-scale prospective study of young, healthy full-term infants with polysomnography and preferably standardized anesthetic techniques as well as similar surgeries should be performed before more objective guidelines can be established. Until that time, we still recommend consideration of short-term postanesthetic apnea monitoring for young full-term infants, even if healthy and undergoing uncomplicated anesthetics.

Dean B. Andropoulos, M.D.
Maurine B. Heard, M.D.
Kristen L. Johnson, M.D.

Anesthesiology
 81:265-266, 1994
 © 1994 American Society of Anesthesiologists, Inc.
 J. B. Lippincott Company, Philadelphia

Sympathetically Maintained Pain after Surgery May Be Prevented by Regional Anesthesia

To the Editor:—Rocco¹ described two patients who had recurrence of their sympathetically maintained pain (SMP) after surgery under general anesthesia. The author then postulated that general anesthesia may cause a recurrence of SMP and suggested that regional anesthesia rather than general anesthesia be used in patients who had suffered previously from reflex sympathetic dystrophy. We strongly agree with Rocco and share his opinion concerning the possible role of general anesthesia, not only in "releasing a dormant pain," but also in favoring the appearance of SMP after surgery.

Anesthesiology, V 81, No 1, Jul 1994

Jonathan T. Clarke, M.D.
Richard W. Rowe, M.D.
 Department of Anesthesiology
 Children's Hospital, Oakland
 747 52nd Street
 Oakland, California 94609

References

1. Tetzlaff JE, Annand DW, Pudimat MA, Nicodemus HF: Postoperative apnea in a full-term infant. *ANESTHESIOLOGY* 69:426-428, 1988
2. Coté CJ, Kelley DH: Postoperative apnea in a full-term infant with a demonstrable respiratory pattern abnormality. *ANESTHESIOLOGY* 72:559-561, 1990
3. Karayan J, LaCoste L, Fuscuardi J: Postoperative apnea in a full-term infant. *ANESTHESIOLOGY* 75:375, 1991
4. Andropoulos DB, Heard MB, Johnson KL, Clarke JT, Rowe RW: Postanesthetic apnea in full-term infants after pyloromyotomy. *ANESTHESIOLOGY* 80:216-219, 1994
5. Albani M, Bentele KHP, Budde C, Schulte FJ: Infant sleep apnea profile: Preterm vs. term infants. *Eur J Pediatr* 143:261-268, 1985
6. Stebbins VA, Poets CF, Alexander JA, Arrowsmith WA, Southall DP: Oxygen saturation and breathing patterns in infancy: I. Full term infants in the second month of life. *Arch Dis Child* 66:569-573, 1991
7. Welborn LG, Ramirez N, Oh TH, Ruttimann UE, Fink R, Guzzeta P, Epstein BS: Postanesthetic apnea and periodic breathing in infants. *ANESTHESIOLOGY* 65:658-661, 1986
8. Kurth CD, Spitzer AR, Broennle AM, Downes JJ: Postoperative apnea in preterm infants. *ANESTHESIOLOGY* 66:483-488, 1987
9. Gregory GA: *Outpatient anesthesia, Anesthesia*. 1st edition. Edited by Miller RD. New York, Churchill Livingstone, 1981, p 1329.
10. Steward DJ: Preterm infants are more prone to complications following minor surgery than are term infants. *ANESTHESIOLOGY* 56:304-306, 1982

(Accepted for publication April 4, 1994.)

In our experience with patients presenting with SMP after surgery, we are confronted by the striking fact that most of them had surgery under general anesthesia. Two recent cases emphasize this point. The first patient was a 26-yr-old man who had severe multiple injuries of the right lower limb and developed SMP after surgery performed under general anesthesia. SMP was relieved with intravenous guanethidine blocks.² Two subsequent surgical procedures were performed using epidural anesthesia, with no relapse of SMP. Eighteen months later, this patient underwent a short-duration general anes-