

Intracerebroventricular Morphine Produces Antinociception by Evoking γ -Aminobutyric Acid Release through Activation of 5-Hydroxytryptamine 3 Receptors in the Spinal Cord

Tomoyuki Kawamata, M.D.,* Keiichi Omote, M.D.,† Masaki Toriyabe, M.D.,‡ Mikito Kawamata, M.D.,§ Akiyoshi Namiki, M.D.||

Background: It has been generally considered that supraspinal morphine activates the serotonergic descending inhibitory system and releases serotonin (5-hydroxytryptamine [5-HT]) in the spinal cord, producing antinociception through activation of 5-HT receptors. The involvement of a spinal γ -aminobutyric acid–mediated (GABAergic) system is also suggested in supraspinal morphine antinociception. It has been reported that spinal GABAergic system contributes to 5-HT₃ receptor-mediated antinociception. In this study, the authors investigated the contribution of spinal 5-HT₃ receptor and the GABAergic system in the intracerebroventricular morphine-induced antinociception.

Methods: Male Sprague-Dawley rats were used. Using the spinal microdialysis method, concentrations of 5-HT and GABA were measured after intracerebroventricular morphine administration. The effect of intracerebroventricular naloxone or spinal perfusion of a selective 5-HT₃ receptor antagonist 3-tropanyl-indole-3-carboxylate methiodide on the spinal release of GABA after intracerebroventricular morphine administration was also examined. In the behavioral study, involvement of 5-HT₃ receptors or GABA_A receptors in the intracerebroventricular morphine-induced antinociceptive effect was investigated using the tail-flick test.

Results: Intracerebroventricular morphine (40 nmol) significantly increased spinal GABA and 5-HT release. Evoked spinal GABA release was reversed by intracerebroventricular naloxone (40 nmol) or spinal perfusion of 3-tropanyl-indole-3-carboxylate methiodide (1 mM). In the behavioral study, intracerebroventricular morphine produced significant antinociception. Intrathecal administration of either GABA_A receptor antagonist bicuculine or 3-tropanyl-indole-3-carboxylate methiodide but not vehicle reversed the morphine-induced antinociceptive effect.

Conclusion: Intracerebroventricular morphine evokes spinal GABA release *via* the activation of 5-HT₃ receptors in the spinal cord, resulting in antinociceptive effect.

It is generally considered that supraspinal morphine produces its antinociceptive effect through the descending serotonergic and noradrenergic neurons projecting to spinal cord. Several spinal mechanisms of supraspinal morphine-induced antinociception have been proposed. In addition to direct inhibition of spinal neurons or primary afferent nerve terminals by noradrenaline and

serotonin (5-hydroxytryptamine [5-HT]), indirect mechanisms *via* spinal interneurons have been suggested.^{1,2} It has been reported that a spinal γ -aminobutyric acid type A (GABA_A) receptor antagonist inhibits the supraspinal morphine-induced antinociception.¹ Furthermore, a spinal GABA-mediated (GABAergic) system contributes to spinally administered 5-HT₃ receptor agonist-mediated antinociception.³ These observations suggest a possible interaction between the serotonergic descending system and a spinal GABA system in supraspinal morphine-induced antinociception.

Among 5-HT receptor subtypes, only the 5-HT₃ receptor is a ligand-gated cation channel.⁴ Electrical stimulation in the periaqueductal gray (PAG) produces inhibition of dorsal horn neuronal responses, and this PAG-induced inhibition is blocked by spinally administered 5-HT₃ receptor antagonists.⁵ Behavioral studies indicated that intrathecal 5-HT-induced antinociception was mediated by 5-HT₃ receptors.⁶ However, it is unlikely that the activation of 5-HT₃ receptors directly inhibits neuronal responses because the activation of 5-HT₃ receptors causes a depolarization by means of increased membrane permeability to monovalent cation.⁴ Supraspinal morphine could evoke GABA release from interneurons through the activation of 5-HT₃ receptors in the spinal cord, resulting in antinociception. However, there has been no direct evidence of supraspinal morphine-induced 5-HT₃ receptor activation and GABA release in the spinal cord.

In the current study, we examined whether supraspinally administered morphine would evoke GABA release through the activation of 5-HT₃ receptors in the spinal cord. We also examined the functional roles of these neurotransmitters in supraspinal morphine-induced antinociception.

Materials and Methods

The protocol for this study was approved by Sapporo Medical University Animal Care and Use Committee (Sapporo, Japan). Experiments were conducted on male Sprague-Dawley rats (weight, 250–350 g; Japan SLC, Hamamatsu, Japan) housed individually in a temperature-controlled (21 ± 1°C) room with a 12-h light–dark cycle and given free access to food and water. Each animal was used in only one experiment.

* Instructor, † Associate Professor, ‡ Postgraduate Student, § Assistant Professor, || Professor and Chairman.

Received from the Department of Anesthesiology, Sapporo Medical University School of Medicine, Sapporo, Japan. Submitted for publication August 10, 2001. Accepted for publication January 3, 2002. Support was provided solely from institutional and/or departmental sources.

Address reprint requests to Dr. Kawamata: Department of Anesthesiology, Sapporo Medical University School of Medicine, South 1, West 16, Chuo-ku, Sapporo, Hokkaido 060-8543, Japan. Address electronic mail to: kawamata@sapmed.ac.jp. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

Intracerebroventricular Cannula and Lumbar Intrathecal Catheter

Rats were anesthetized with pentobarbital (50 mg/kg administered intraperitoneally) and were placed in a stereotaxic head holder to implant an intracerebroventricular guide cannula. The skull was exposed, and a guide cannula equipped with a 32-gauge stylet was directed into the left lateral ventricle according to Paxinos and Watson⁷ (anterior-posterior +0.8 mm, midline +1.4 mm, dorsal-ventral 3.8 mm with respect to bregma) and was attached to the skull with three anchoring screws and cemented in place. For intracerebroventricular injection, the stylet was removed and the injection cannula was inserted.

For the behavioral study, a lumbar intrathecal catheter was implanted in the rats with an intracerebroventricular guide cannula. A polyethylene intrathecal catheter (PE-10; Becton Dickinson, Franklin Lakes, NJ) was inserted 15 mm cephalad into the lumbar subarachnoid space at the L4-L5 intervertebral level, with the tip of the catheter located near the lumbar enlargement of the spinal cord to administer the drugs intrathecally. The catheter was tunneled subcutaneously and externalized through the skin in the neck region. At least 6 days of postsurgical recovery were allowed before animals were used in the behavioral study.

Microdialysis Catheter Implantation

The spinal cord dialysis probe was prepared according to our modification of the method described by Skilling *et al.*⁸ The dialysis probe was constructed from a 1-cm length of dialysis fiber (an ID of 200 μm , an OD of 220 μm , and 50-kd molecular weight cutoff; DM-22, Eicom Co., Kyoto, Japan) which was coated with a thin layer of epoxy glue (Devcon Co., Danvers, MA) along the whole length, except for a 2-mm region in the middle. To make the fiber firm enough for implantation, a Nichrome-Formvar wire with a 78- μm ID (A-M Systems, Inc., Everett, WA) was passed through the fiber. Each end of the fiber was attached to 2-cm polyethylene catheters (PE-10; Becton Dickinson, Franklin Lakes, NJ), and each end of the polyethylene catheters was then attached to an 8-cm teflon tube with an ID of 100 μm and an OD of 400 μm (JT-10; Eicom). Seven days after an intracerebroventricular guide cannula implantation, rats were anesthetized with pentobarbital (50 mg/kg administered intraperitoneally) and incised along the dorsal midline from T2 to L2. The lateral surfaces of vertebra L1 were exposed, and bilateral holes were carefully made through the bone, exposing the spinal cord laterally at the level of the dorsal horn. A dialysis tube was placed through the holes by hand, passing transversely through the dorsal spinal cord. The two distal ends of the probe were tunneled subcutaneously and externalized through the skin in the neck region. The experiments were performed 18–24 h after the implantation of the dialysis

probe. After a recovery period, the animals showing any signs of limb paralysis or impaired movement were excluded in this study. After each experiment, we perfused methylene blue dye through the dialysis probe to verify the position of the dialysis fiber, and then rats were killed with an overdose of pentobarbital. The data to be reported are from rats in which methylene blue dye remained in the area of the dorsal half of the dorsal horn.

Microdialysis Study

The animals freely moved in a plastic cage with dimensions of 30 \times 30 \times 35 cm during the dialysis experiments. A liquid switch (SI-50; Eicom) was placed between the syringe pump and the dialysis probe to enable a different drug to be administered locally *via* the probe. The dialysis probe was perfused with artificial cerebrospinal fluid (ACSF; 140 mM NaCl, 4.0 mM KCl, 1.26 mM CaCl_2 , 1.15 mM MgCl_2 , 2.0 mM Na_2HPO_4 , 0.5 mM NaH_2PO_4 , pH 7.4) at a constant flow rate of 3 $\mu\text{l}/\text{min}$. The samples were collected as 20-min fractions and divided into two samples: 20 μl dialysate collected for analysis of GABA and 20 μl for 5-HT. The samples were frozen at -80°C until analysis. Three consecutive samples were collected for determination of basal concentrations 180 min after starting perfusion of ACSF. After obtaining three consecutive samples, 40 nmol/3 μl of intracerebroventricular morphine (Sankyo Co., Tokyo, Japan) was administered over a 120-s period. The samples were collected up to 360 min after morphine administration. To examine the naloxone sensitivity and the involvement of spinal 5-HT₃ receptor activation in morphine-induced spinal GABA release, either intracerebroventricular naloxone (Sigma, St. Louis, MO) or the selective 5-HT₃ receptor antagonist, 3-tropanyl-indole-3-carboxylate methiodide (TICM; Research Biochemical International, Natick, MA) was administered 180 min after morphine administration. Naloxone 40 nmol was injected in a volume of 3 μl over a 120-s period. TICM (1 mM) was perfused through the dialysis catheter for 60 min. All drugs used in microdialysis study were prepared in ACSF.

Analysis of Serotonin

Serotonin in the dialysate was analyzed using high-performance liquid chromatography with electrochemical detection (Degasser, DG-100; liquid chromatograph, EP-100; electrochemical detector, ECD-100; Eicom). The chromatographic conditions were as follows: column, Eicompak (CA-5ODS 2.1 \times 150 mm; Eicom); mobile phase, 0.1 M phosphate buffer (pH 6.0) containing 20.0% methanol, 0.02 mM EDTA, and 0.72 mM sodium 1-octanesulfonate for the analysis of 5-HT; working electrode, glassy carbon (WE-3G, Eicom); flow rate, 0.23 ml/min. Detector voltage and detector temperature were set at 0.45 V and at 25.0°C, respectively. Retention time for

5-HT was 14.85 min. The detection limit for 5-HT analysis is 0.05 fg/20 μ l.

Analysis of γ -Aminobutyric Acid

γ -Aminobutyric acid in the dialysate was analyzed using high-performance liquid chromatography with fluorescence detection (Degasser, DG-100; liquid chromatograph, EP-100; fluorescence detector, FLD-370; Eicom) after derivatization with o-phthalaldehyde (Sigma). The o-phthalaldehyde derivatizing reagent was prepared by dissolving o-phthalaldehyde (54 mg) in absolute methanol (1 ml) and adding 2-mercaptoethanol (40 μ l) and sodium carbonate (0.1 M, pH 9.5, 99 ml). Ten microliters of this derivatizing reagent was mixed with 20 μ l of the dialysate and allowed to react for 2.5 min. Twenty microliters of this solution was analyzed. The chromatographic conditions were as follows: column, Eicompak (SC-50DS 2.1 \times 150 mm; Eicom); mobile phase, 0.1 M sodium dihydrogenphosphate and 0.1 M disodium hydrogenphosphate (pH 3.5) containing 50.0% methanol and 0.1 mM disodium EDTA; flow rate, 0.23 ml/min. Detector temperature was set at 30°C. Retention time for GABA was 14.83 min. The detection limit for GABA analysis is 0.2 pg/20 μ l.

Behavioral Study

A behavioral study was performed separately from the microdialysis study. The tail-flick test was used to assess thermal nociceptive threshold. Tail-flick testing was performed by monitoring latency to withdrawal from the heat source (a 50-W projection lamp bulb) focused on a distal segment of the tail, using a thermal analgesimeter (KN-205E; Natsume, Tokyo, Japan). A cutoff time of 15.0 s was used to minimize damage to the skin of the tail. After determination of baseline tail-flick latencies, 40 nmol/3 μ l of intracerebroventricular morphine was administered. The tail-flick latencies were measured at 30, 60, 120, 180, 240, 300, and 360 min after the intracerebroventricular injection. To examine the involvement of spinal 5-HT₃ receptors and GABA_A receptors in morphine analgesia, either TICM or a GABA_A receptor antagonist bicuculline (Sigma) was intrathecally administered 170 min after morphine administration, and the effect of the antagonist on tail-flick latencies was evaluated 10 min later. TICM and bicuculline were freshly dissolved in ACSF in concentrations that allowed intrathecal injections in 10- μ l volumes. Intrathecal drug administration was accomplished by using a microinjection syringe (Hamilton, Reno, NV) connected to the intrathecal catheter in awake, briefly restrained rats. All intrathecal drugs were administered manually over 10 s.

Data Analysis

In the microdialysis study, the changes of GABA and 5-HT concentrations are presented as mean \pm SD of percentage of basal concentrations. Data were com-




Fig. 1. Changes of 5-hydroxytryptamine (5-HT) concentration in the spinal cord after administration of 40 nmol intracerebroventricular morphine. Data are presented as mean \pm SD of the percent of basal concentration, $n = 8$. * $P < 0.05$ compared with basal concentration.

pared with basal concentration using a one-way analysis of variance for repeated measures followed by Dunnett test within a single group, and were analyzed using a two-way analysis of variance for repeated measures followed by the Tukey Kramer test for between-group comparisons. In the behavioral study on morphine antinociception, the values of withdrawal latency were converted to percent maximum possible effect:

$$\%MPE = \frac{(\text{postdrug latency} - \text{baseline latency})}{(\text{cutoff latency} - \text{baseline latency})} \times 100.$$

Changes in percent maximum possible effect were compared with baseline using a one-way analysis of variance for repeated measures followed by Dunnett test within a single group, and analyzed using a paired t test for the comparison between pretreatment and posttreatment in the behavioral study. $P < 0.05$ was considered statistically significant.

Results

Microdialysis Study

Serotonin Analysis. The basal concentration of 5-HT was 2.57 ± 1.18 pg/20 μ l. The coefficient of variation among the three consecutive samples for determination of basal concentrations was less than 7%. Intracerebroventricular morphine (40 nmol) evoked a significant increase in spinal 5-HT concentration ($P < 0.01$; fig. 1). Significant increases in 5-HT concentration were observed 40–240 min after morphine administration.

γ -Aminobutyric Acid Analysis. The basal concentration of GABA was 0.27 ± 0.14 pg/20 μ l. The coefficient of variation among the three consecutive samples for determination of basal concentrations was less than 5%.




Fig. 2. Changes of γ -aminobutyric acid (GABA) concentration in the spinal cord. Open squares indicate GABA concentrations in the rats with administration of 40 nmol intracerebroventricular morphine ($n = 8$). Open circles indicate GABA concentrations in the rats with intracerebroventricular morphine followed by intracerebroventricular naloxone ($n = 6$). Intracerebroventricular naloxone was administered 180 min after morphine administration. Closed circles indicate GABA concentrations in the rats with intracerebroventricular morphine followed by spinal perfusion of 3-tropanyl-indole-3-carboxylate methiodide (TICM; $n = 7$). TICM was perfused 180–240 min after morphine administration. Data are presented as mean \pm SD of the percent of basal concentration. # $P < 0.05$ between intracerebroventricular morphine and intracerebroventricular morphine followed by TICM. * $P < 0.05$ between intracerebroventricular morphine and intracerebroventricular morphine followed by intracerebroventricular naloxone. i.c.v. = intracerebroventricular.

Figure 2 shows the effect of 40 nmol intracerebroventricular morphine on spinal GABA concentration. GABA concentration was gradually increased after intracerebroventricular morphine administration and reached a plateau from 180 to 260 min after morphine administration. Significant increases in GABA concentration were observed from 80 up to 360 min after administration of morphine compared with the basal concentration ($P < 0.01$). Intracerebroventricular naloxone (40 nmol) administered at 180 min after morphine significantly reversed the morphine-induced increase in GABA concentration ($P < 0.01$). Spinal perfusion of 1 mM TICM for 60 min (from 180 to 240 min after morphine) also reversed morphine-induced increase in GABA concentration ($P < 0.01$). GABA concentrations 180 min after morphine administration were comparable among the three different treatments. TICM (1 mM) alone did not




Fig. 3. Effect of spinal perfusion of 3-tropanyl-indole-3-carboxylate methiodide (TICM) on the basal γ -aminobutyric acid (GABA) concentration. TICM was perfused for 240 min. Data are presented as mean \pm SD of the percent of basal concentration, $n = 5$.

significantly affect the basal spinal GABA concentration for at least 240 min (fig. 3).

Behavioral Study

Effects of Intrathecal Bicuculline or TICM. Intrathecal administration of bicuculline at doses of greater than 0.02 μ g shortened the tail-flick latency in the tail-flick test. For example, 0.2 μ g bicuculline shortened the tail-flick latency, which was observed within 5 min after the administration and continued for 30 min (table 1). Doses of intrathecal TICM greater than 1.0 μ g also shortened the tail-flick latency. For example, 10.0 μ g TICM shortened the tail-flick latency that was observed within 5 min and continued for 30 min (table 2). Neither 0.02 μ g intrathecal bicuculline nor 1.0 μ g TICM affected the basal tail-flick latency (tables 1 and 2). Therefore, these doses were used in the behavioral study, and the antagonistic effects of these drugs were assessed at 10 min after intrathecal administration.

Antagonistic Effects of Intrathecal Bicuculline and TICM on the Intracerebroventricular Morphine-induced Antinociception

The mean baseline tail-flick latency in this experiment was 5.7 s (range, 5.1–6.2 s). Figure 4 shows the time course of the change in percent maximum possible effect after

Table 1. Change of Tail-flick Latency after Intrathecal Bicuculline

	Time after Intrathecal Administration (min)								
	0	5	10	15	20	30	45	60	
0.02 μ g Bicuculline	5.5 \pm 0.2	5.4 \pm 0.2	5.3 \pm 0.2	5.4 \pm 0.3	5.4 \pm 0.2	5.5 \pm 0.3	5.6 \pm 0.1	5.4 \pm 0.5 (s)	
0.2 μ g Bicuculline	5.6 \pm 0.1	4.8 \pm 0.4*	3.3 \pm 0.5*	3.6 \pm 0.4*	3.8 \pm 0.5*	4.2 \pm 0.6*	4.8 \pm 0.6	5.4 \pm 0.5 (s)	

Values are mean \pm SD of tail-flick latency ($n = 6$ in each dose).

* $P < 0.05$ compared with control (value at 0 min).

Table 2. Changes of Tail-flick Latency after Intrathecal TCM

	Time after Intrathecal Administration (min)							
	0	5	10	15	20	30	45	60
1.0 μ g TCM	5.6 \pm 0.1	5.5 \pm 0.2	5.4 \pm 0.2	5.4 \pm 0.3	5.6 \pm 0.2	5.5 \pm 0.2	5.6 \pm 0.4	5.5 \pm 0.1 (s)
10.0 μ g TCM	5.7 \pm 0.2	3.8 \pm 0.4*	3.6 \pm 0.4*	3.9 \pm 0.4*	4.5 \pm 0.6*	4.7 \pm 0.3*	5.4 \pm 0.3	5.6 \pm 0.1 (s)

Values are mean \pm SD of tail-flick latency (n = 6 in each dose).

* $P < 0.05$ compared with control (value at 0 min).

TCM = 3-tropanyl-indole-3-carboxylate methiodide.

intracerebroventricular morphine administration. Intracerebroventricular morphine (40 nmol) produced significant antinociceptive effects from 30 min up to 360 min after its administration. The maximal antinociceptive effect was observed 30–180 min after morphine administration. Figure 5 shows the effects of intrathecal bicuculline and TCM on the intracerebroventricular morphine-induced antinociceptive effect. We evaluated the antagonistic effects of bicuculline and TCM 180 min after morphine administration, because at this time point, the maximal effect of morphine in GABA analysis was observed (fig. 2). Intrathecal bicuculline significantly reversed the morphine-induced antinociception ($P < 0.05$). Intrathecal TCM also significantly reversed the morphine-induced antinociception ($P < 0.05$). On the other hand, intrathecal ACSF alone did not produce any significant changes in the tail-flick latency. There were no significant differences in percent maximum possible effect before intrathecal administration among the three different treatments.

Discussion

Intracerebroventricular Morphine-induced Spinal Serotonin Release

The current study showed that intracerebroventricular morphine evoked spinal 5-HT release. Previous reports




Fig. 4. Time course of percent maximum possible effect (%MPE) after administration of 40 nmol intracerebroventricular morphine in the tail-flick test. Data are presented as mean \pm SD, n = 12.

also found morphine-induced spinal 5-HT release.⁹⁻¹¹ The major source of serotonergic projections to the dorsal horn are neurons in the rostral ventromedial medulla, which contains both opioid peptides and opioid receptors.^{12,13} Several investigators found that electrical or chemical stimulation of the dorsal lateral funiculus, bulbospinal neurons, or the nucleus raphe magnus evoked 5-HT release in the dorsal horn of the spinal cord.¹⁴⁻¹⁶ Therefore, it would be considered that morphine evokes spinal 5-HT release through activation of rostral ventromedial medulla serotonergic neurons by the inhibition of inhibitory interneurons. However, this may not be always consistent. First, Matos *et al.*¹⁷ concluded that intracerebroventricular morphine at an analgesic dose did not necessarily increase the extracellular concentration of 5-HT in the spinal dorsal horn using high-performance liquid chromatography with electrochemical detection, because morphine increased 5-HT




Fig. 5. Effects of intrathecal bicuculline, 3-tropanyl-indole-3-carboxylate methiodide (TCM), and artificial cerebrospinal fluid (ACSF) on the antinociceptive effect of intracerebroventricular morphine. Pre values indicate percent maximum possible effects (%MPE) before intrathecal drug administration (170 min after intracerebroventricular morphine). Post values indicate percent maximum possible effects 10 min after intrathecal drug administration (180 min after intracerebroventricular morphine). Data are presented as mean \pm SD. * $P < 0.05$ between pre and post values. n.s. = no statistical significance. ACSF-treated rats, n = 6; TCM-treated rats, n = 8; bicuculline-treated rats, n = 8.

release in only two of four rats. The discrepancy between their results and ours may be caused by the sensitivity of assay system for 5-HT. Baseline 5-HT concentrations were close to the sensitivity limit in the high-performance liquid chromatography with electrochemical detection, and hence it is difficult to measure small changes reliably.¹⁸ We used a smaller diameter (2.1 mm) detector column in the current study, compared with that used in the study by Matos *et al.*¹⁷ (4.6 mm). This column enabled us to analyze 5-HT with high sensitivity. Using this assay system, morphine at the dose we used (40 nmol = 11.4 μ g), which was almost comparable to the dose used by Matos *et al.* (10 μ g), consistently increased spinal 5-HT concentration in all eight rats in the current study, and the degree of increase in 5-HT concentration was approximately 150% of the basal concentration. We performed the experiments 18–24 h after the implantation of the dialysis probe because of prevention of the decrease in recovery rate caused by gliosis. Therefore, there may be the possibility that the acute trauma of probe implantation modified the effect of morphine on 5-HT release. Second, electrophysiologic studies have demonstrated that most serotonergic rostral ventromedial medulla neurons are not affected by morphine.^{19–21} However, those studies used the anesthetized animal. There is a possibility that anesthesia reduced serotonergic neuronal activity,^{18,22} and serotonergic neuronal activity is lower during sleep than wakefulness.²³ We used the awake free-moving rats implanted with microdialysis probe in this study.

Intracerebroventricular Morphine-induced Activation of Spinal 5-HT₃ Receptors and Spinal γ -Aminobutyric Acid Release

Several lines of study have suggested that supraspinal morphine or PAG stimulation activates a spinal GABAergic system.^{1,3,5,24–26} However, there has not been direct evidence that supraspinal morphine evokes spinal GABA release. In the current study, we clearly demonstrated that intracerebroventricular morphine evokes spinal GABA release concomitant with 5-HT release. There was the discrepancy in time course between intracerebroventricular morphine-induced increases in 5-HT and GABA concentrations in our results. Spinal microdialysis does not directly measure neurotransmitter concentrations at synaptic sites, but rather neurotransmitter concentrations that have diffused into the extracellular space. Diffusion into the extracellular space is likely to involve processes of uptake and metabolism, which can modify the amount of neurotransmitter in the dialysate. Therefore, it is possible that diffusion into the extracellular space would contribute to the discrepancy in time course between 5-HT and GABA concentration.

There are at least four subtypes of 5-HT receptors (5-HT₁, 5-HT₂, 5-HT₃, and 5-HT₄) in the spinal cord.^{27,28} Among the subtypes of 5-HT receptors, 5-HT₃ receptors

are unique because G proteins are not affected by agonist binding to 5-HT₃ receptors. 5-HT₃ receptors are directly linked to the opening of nonselective monovalent cation channels⁴ and should mediate neuronal excitation. Alhaider *et al.*³ demonstrated that the activation of the GABAergic system contributed to spinal 5-HT₃ receptor-mediated antinociception. Therefore, we focused on 5-HT₃ receptors. TICM used in the current study is a potent and selective 5-HT₃ receptor antagonist and has been commonly known as IC 205-930. Radioligands of IC 205-930 have been used to detect 5-HT₃ receptor binding site in the central nervous system.²⁹ In the current study, blockade of spinal 5-HT₃ receptors inhibited intracerebroventricular morphine-induced spinal GABA release. This indicates that morphine activates supraspinal serotonergic neurons projecting to the spinal cord, and then released 5-HT excites spinal GABAergic interneuron through 5-HT₃ receptors. Immunohistochemical and autoradiographic studies revealed that 5-HT₃ receptors are restricted to the superficial layers of the dorsal horn in the spinal cord.^{29–31} Intense GABA immunoreactivities are also found within laminae I–III of the spinal cord.^{32,33} It would be supposed that the GABAergic neuron with 5-HT₃ receptors are directly activated or activation of presynaptic 5-HT₃ receptors indirectly induce the activation of GABAergic neuron. There is a report that the 5-HT₃ receptors mainly were located in the primary afferent terminal in the dorsal horn.³⁴ In addition, Peng *et al.*³⁵ recently suggested that 5-HT₃ receptors located in central terminals of primary afferents might contribute to the generation of dorsal root reflex, leading to presynaptic inhibition. Therefore, further study is necessary to clarify the mechanism of 5-HT₃ receptor activation-induced GABA release.

Contributions of 5-HT₃ Receptor Activation and γ -Aminobutyric Acid Release to Supraspinal Morphine-induced Antinociceptive Effect

In a behavioral study, we examined whether spinal GABA release through the activation of 5-HT₃ receptors observed in our microdialysis study would be a component of supraspinal morphine-induced antinociception. GABA and 5-HT concentrations were measured within dorsal horn at the level of the L1 vertebra, and the tail-flick test was used in the behavioral study. The primary afferent neurons responsible for tail-flick response largely innervate in the dorsal horn of the sacrococcygeal spinal cord. In this study, GABA and 5-HT concentrations were measured within dorsal horn at the level of the L1 vertebra at which L4 nerve root mainly terminated. Therefore, results of the current microdialysis study may not directly reflect the changes of GABA and 5-HT in the sacral spinal cord. However, neurons originating in the nucleus raphe magnus, on which intracerebroventricular morphine acts, project to all levels of the spinal cord.^{36–38} Therefore, intracerebroventricular

morphine-induced GABA and 5-HT release we observed should contribute to antinociception in the tail-flick test.

It has been generally considered that supraspinal morphine produces antinociceptive effects through the serotonergic and noradrenergic descending inhibitory systems. In respect to the serotonergic mechanism, the analgesic action of morphine is reduced by destruction of the serotonergic innervation of the spinal cord with selective neurotoxins³⁹ or lesions of the medullary raphe,^{40,41} which contains the cell of the origin of the bulbospinal 5-HT pathway innervating the spinal dorsal horn. Intrathecal injection of a 5-HT receptor antagonist also attenuates the analgesia produced by microinjection of morphine into the PAG.⁴² In addition, intrathecal 5-HT produces antinociceptive effects. However, the nature of the receptors involved in the 5-HT-induced modulation of pain in the spinal cord remains to be elucidated. To the best of our knowledge, there has been no report that examined the contribution of spinal 5-HT₃ receptors to intracerebroventricular morphine-induced antinociception in behavioral tests. The results of the current study have shown that spinal 5-HT₃ receptors are involved in supraspinal morphine-induced antinociceptive effects. In contrast, several studies found only a minor role of the serotonergic system in supraspinal morphine-induced antinociceptive effect.⁴³⁻⁴⁵ However, because these studies used methysergide as the 5-HT receptor antagonist, which blocks 5-HT_{1/2}, not 5-HT₃ receptors, the role of a serotonergic mechanism may be underestimated. The current study showed that TICM did not completely reverse the morphine-induced antinociceptive effect. This suggests the involvement of the other subtypes of 5-HT receptors or noradrenergic system in intracerebroventricular morphine-induced analgesia. Indeed, it has been reported that GABA_A receptor response is potentiated by the activation of 5-HT₂ receptor,⁴⁶ and that 5-HT₁ receptor activation inhibits the nociceptive transmission.⁴⁷ The contribution of noradrenergic system in intracerebroventricular morphine-induced analgesia was also well demonstrated.

Several investigators reported that the spinal GABA system contributes to nucleus raphe magnus or PAG stimulation- and supraspinal morphine-induced antinociceptive effects.^{1,3,5,25,26,48,49} Although our microdialysis study demonstrated that intracerebroventricular morphine gradually increased spinal GABA release with a maximal effect at 180-240 min after administration, an antinociceptive effect was observed 30 min after the administration and thereafter. Because we did not assess the antagonistic effect of bicuculline at the early phase of morphine-induced antinociception, the involvement of GABA in morphine analgesia at the early phase is unclear. We speculate that another mechanism such as noradrenergic or direct serotonergic inhibitions, in addition to GABAergic inhibition, might be involved in the antinociceptive effect observed in the early phase.

In conclusion, the current study showed that intracerebroventricular morphine produced antinociceptive effect, in part through GABA release by the activation of 5-HT₃ receptors in the spinal cord.

References

1. Suh HW, Kim YH, Choi YS, Choi SR, Song DK: Effects of GABA receptor antagonists injected spinally on antinociception induced by opioids administered supraspinally in mice. *Eur J Pharmacol* 1996; 307:141-7
2. Sweeney MI, White TD, Sawynok J: Intracerebroventricular morphine releases adenosine and adenosine 3',5'-cyclic monophosphate from the spinal cord via a serotonergic mechanism. *J Pharmacol Exp Ther* 1991; 259:1013-8
3. Alhaider AA, Lei SZ, Wilcox GL: Spinal 5-HT₃ receptors-mediated antinociception: Possible release of GABA. *J Neurosci* 1991; 11:1881-8
4. Derkach V, Surprenant A, North RA: 5-HT₃ receptors membrane ion channels. *Nature* 1989; 339:706-9
5. Peng YB, Lin Q, Willis WD: The role of 5-HT₃ receptors in periaqueductal gray-induced inhibition of nociceptive dorsal horn neurons in rats. *J Pharmacol Exp Ther* 1996; 276:116-24
6. Bardin L, Lavarenne J, Eschaler A: Serotonin receptor subtypes involved in the spinal antinociceptive effect of 5-HT in rats. *Pain* 2000; 86:11-8
7. Paxinos G, Watson C: *The Rat Brain in Stereotaxic Coordinates*, 2nd edition. Edited by Paxinos G, Watson C. Australia, Academic Press, 1986, pp 21-2
8. Skilling SR, Smullin DH, Beitz AJ, Larson AA: Extracellular amino acid concentrations in the dorsal spinal cord of freely moving rats following veratridine and nociceptive stimulation. *J Neurochem* 1988; 51:127-32
9. Yaksh TL, Tyce GM: Microinjection of morphine into the periaqueductal gray evokes the release of serotonin from spinal cord. *Brain Res* 1979; 171:176-81
10. Bardon T, Ruckebusch: Changes in 5-HIAA and 5-HT levels in lumbar CSF following morphine administration to conscious dogs. *Neurosci Lett* 1984; 49:147-51
11. Suzuki Y, Taguchi K: In vivo voltammetric studies of the effect of morphine on the serotonergic system in the cat spinal cord. *Brain Res* 1986; 398:413-8
12. Oliveras JL, Bourgoin S, Hery F, Besson JM, Hamon M: The topographical distribution of serotonergic terminals in the spinal cord of the cat: Biochemical mapping by the combined use of microdissection and microassay procedures. *Brain Res* 1977; 138:393-406
13. Kwiat GC, Basbaum AI: The origin of brainstem noradrenergic and serotonergic projections to the spinal cord dorsal horn in the rat. *Somatosens Mot Res* 1992; 9:157-73
14. Sorkin LS, McAdoo DJ, Willis WD: Raphe magnus stimulation-induced antinociception in the cat is associated with release of amino acids as well as serotonin in the lumbar dorsal horn. *Brain Res* 1993; 618:95-108
15. Pilowsky PM, Kapoor V, Minson JB, West MJ, Chalmers JP: Spinal cord serotonin release and raised blood pressure after brainstem kainic acid injection. *Brain Res* 1986; 366:354-7
16. Sorkin LS, Steinman JL, Hughes MG, Willis WD, McAdoo DJ: Microdialysis recovery of serotonin released in spinal cord dorsal horn. *J Neurosci Methods* 1988; 23:131-8
17. Matos FF, Rollema H, Brown JL, Basbaum AI: Do opioids evoke the release of serotonin in the spinal cord? An in vivo microdialysis study of the regulation of extracellular serotonin in the rat. *Pain* 1992; 48:439-47
18. Tao R, Auerbach SB: Anesthetics block morphine-induced increases in serotonin release in rat CNS. *Synapse* 1994; 18:307-14
19. Gao K, Chen DO, Genzen JR, Mason P: Activation of serotonergic neurons in the raphe magnus is not necessary for morphine analgesia. *J Neurosci* 1998; 18:1860-8
20. Potrebic SB, Fields HL, Mason P: Serotonin immunoreactivity is contained in one physiological cell class in the rat rostral ventromedial medulla. *J Neurosci* 1994; 14:1655-65
21. Chiang CY, Pan ZZ: Differential responses of serotonergic and non-serotonergic neurons in nucleus raphe magnus to systemic morphine in rats. *Brain Res* 1985; 337:146-50
22. Rivot JP, Pointis D, Besson JM: Morphine increases 5-HT metabolism in the nucleus raphe magnus: An in vivo study in freely moving rats using 5-hydroxyindole electrochemical detection. *Brain Res* 1988; 446:333-42
23. Jacobs BL, Fornal CA, Wilkinson LO: Neurophysiological and neurochemical studies of brain serotonergic neurons in behaving animals. *Ann N Y Acad Sci* 1990; 600:260-8
24. Xu TL, Pang ZP, Li JS, Akaike N: 5-HT potentiate of the GABA response in the rat sacral dorsal commissural neurons. *Br J Pharmacol* 1998; 124:779-87
25. Rady JJ, Fujimoto JM: Supraspinal delta₂ opioid agonist analgesia in swiss-webster mice involves spinal GABA receptors. *Pharmacol Biochem Behav* 1996; 54:363-9
26. Lin Q, Peng Y, Willis WD: Glycine and GABA antagonists reduce the inhibition of primate spinothalamic tract neurons produced by stimulation in periaqueductal gray. *Brain Res* 1994; 654:286-302
27. Hoyer D, Clarke DE, Fozard JR, Hartig PR, Martin GR, Mylecharane EJ, Saxena PR, Humphrey PP: International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (serotonin). *Pharmacol Rev* 1994; 46:157-203

28. Frazer A, Hensler JM: Serotonin, Basic Neurochemistry, 5th edition. Edited by Siegel GJ, Agranoff BW, Albers RW, Molinoff PB. New York, Raven, 1993, pp 283-308
29. Waeber C, Hoyer D, Palacios JM: 5-hydroxytryptamine₃ receptors in the human brain: Autoradiographic visualization using [³H]ICS 205-930. *Neuroscience* 1989; 31:393-400
30. Morales M, Battenberg E, Bloom FE: Distribution of neurons expressing immunoreactivity for the 5HT₃ receptor subtype in the rat brain and spinal cord. *J Comp Neurol* 1998; 402:385-401
31. Laporte AM, Koscielniak T, Ponchant M, Verge D, Hamon H, Gozlan H: Quantitative autoradiographic mapping of 5HT₃ receptors in the rat CNS using [¹²⁵I]iodo-zacopride and [³H]zacopride as radioligands. *Synapse* 1992; 10: 271-81
32. Magoul R, Onteniente B, Geffard M, Calas A: Anatomical distribution and ultrastructural organization of the GABAergic system in the rat spinal cord: An immunocytochemical study using anti-GABA antibodies. *Neuroscience* 1987; 20:1001-9
33. Todd AJ, McKenzie J: GABA-immunoreactive neurons in the dorsal horn of the rat spinal cord. *Neuroscience* 1989; 31:799-806
34. Kidd EJ, Laporte AM, Langlois X, Fattaccini CM, Doyen C, Lombard MC, Gozlan H, Hamon M: 5-HT₃ receptors in the rat central nervous system are mainly located on nerve fibres and terminals. *Brain Res* 1993; 612:289-98
35. Peng YB, Wu J, Willis WD, Kenshalo DR: GABA_A and 5-HT₃ receptors are involved in dorsal root reflexes: Possible role in periaqueductal gray descending inhibition. *J Neurophysiol* 2001; 86:49-58
36. Basbaum AI, Clanton CH, Fields HL: Three bulbospinal pathways from the rostral medulla of the cat: An autoradiographic study of pain modulating systems. *J Comp Neurol* 1978; 178:209-24
37. Holstege G, Kuypers HG: The anatomy of brain stem pathways to the spinal cord in cat: A labeled amino acid tracing study. *Prog Brain Res* 1982; 57:145-75
38. Martin RF, Haber LH, Willis WD: Primary afferent depolarization of identified cutaneous fibers following stimulation in medial brain stem. *J Neurophysiol* 1979; 42:779-90
39. Vogt M: The effect of lowering the 5-hydroxytryptamine content of the rat spinal cord on analgesia produced by morphine. *J Physiol* 1974; 236:483-98
40. Yaksh TL, Plant RL, Rudy TA: Studies on the antagonism by raphe lesions of the antinociceptive action of systemic morphine. *Eur J Pharmacol* 1977; 41:399-408
41. Proudfit HK: Effects of raphe magnus and raphe pallidus lesions on morphine-induced analgesia and spinal cord monoamines. *Pharmacol Biochem Behav* 1980; 13:705-14
42. Yaksh TL: Direct evidence that spinal serotonin and noradrenaline terminals mediate the spinal antinociceptive effects of morphine in the periaqueductal gray. *Brain Res* 1979; 160:180-5
43. Tseng LL, Tang R: Differential actions of the blockade of spinal opioid, adrenergic and serotonergic receptors on the tail-flick inhibition induced by morphine microinjected into dorsal raphe and central gray in rats. *Neuroscience* 1989; 33:93-100
44. DeLander GE, Wahl JJ: Morphine (intracerebroventricular) activates spinal systems to inhibit behavior induced by putative pain neurotransmitters. *J Pharmacol Exp Ther* 1989; 251:1090-5
45. Wigdor S, Wilcox GL: Central and systemic morphine-induced antinociception in mice: Contribution of descending serotonergic and noradrenergic pathways. *J Pharmacol Exp Ther* 1987; 242:90-5
46. Li H, Lang B, Kang JF, Li YQ: Serotonin potentiates the response of neurons of the superficial laminae of the rat spinal dorsal horn to gamma-aminobutyric acid. *Brain Res Bull* 2000; 52:559-65
47. el-Yassir N, Fleetwood-Walker SM, Mitchell R: Heterogeneous effects of serotonin in the dorsal horn of rat: The involvement of 5-HT1 receptor subtypes. *Brain Res* 1988; 456:147-58
48. McGowan MK, Hammond DL: Antinociception produced by microinjection of L-glutamate into the ventromedial medulla of the rat: mediation by spinal GABAA receptors. *Brain Res* 1993; 620:86-96
49. Lin Q, Peng TB, Willis WD: Role of GABA receptor subtypes in inhibition of primate spinothalamic tract neurons: Difference between spinal and periaqueductal gray inhibition. *J Neurophysiol* 1996; 75:109-23