

Isoflurane Pretreatment Inhibits Cytokine-induced Cell Death in Cultured Rat Smooth Muscle Cells and Human Endothelial Cells

Manuela J. M. de Klaver, M.D.,* Lee Manning, M.D.,† Lisa A. Palmer, Ph.D.,‡ George F. Rich, M.D., Ph.D.§

Background: Anesthetics are protective during ischemic-reperfusion injury and associated inflammation; therefore, the authors hypothesized that anesthetic pretreatment may provide protection in culture from cytokine-induced cell death.

Methods: Rat vascular smooth muscle (VSM) cell and human umbilical vascular endothelial cell (HUVEC) cultures were used to determine whether pretreatment with 30 min of isoflurane decreases cell death from tumor necrosis factor α (TNF- α), interleukin 1 (IL-1 β), and interferon (IFN- γ) alone or in combination. Cell survival and viability were determined by trypan blue staining and cell proliferation assay, as well as by DNA fragmentation assays. The roles of protein kinase C (PKC) and adenosine triphosphate-sensitive potassium (K_{ATP}) channels in mediating isoflurane (and halothane) protection were evaluated with the antagonists staurosporine or glibenclamide in cytokine- and also hydrogen peroxide (H_2O_2)-induced cell death.

Results: Pretreatment with 1.5% isoflurane immediately prior to cytokine exposure increased cell survival and viability from cytokines by 10–60% for 24, 48, 72, and 96 h in VSMs and up to 72 h in HUVECs. DNA fragmentation (TUNEL) was also attenuated by isoflurane. Isoflurane was equally effective in VSMs at 0.75, 1.5, and 2.5%, whereas in HUVECs, 1.5 and 2.5% were more effective than 0.75%. In VSMs, isoflurane administered 1 h prior to or simultaneously with cytokines was also effective, whereas isoflurane 2 h prior to cytokines was less effective, and either 4 h prior to or 30 min after cytokines was not effective. In both cytokine- and H_2O_2 -induced cell death, isoflurane and halothane pretreatment were equally protective, and staurosporine and glibenclamide attenuated the protective effect.

Conclusions: Thirty minutes of isoflurane attenuates cytokine-induced cell death and increases cell viability in VSMs for 96 h and in HUVECs for 72 h. Isoflurane must be administered less than 2 h prior to or simultaneously with the cytokines to be protective. These initial inhibitor studies suggest involvement of PKC and K_{ATP} channels in isoflurane and halothane protection against both cytokine- and H_2O_2 -induced cell death of VSMs and HUVECs.

ENDOTHELIAL and vascular smooth muscle (VSM) cells play key roles in the physiologic function of the pulmonary and systemic vasculature. The endothelium synthesizes and releases vasoactive mediators that modulate vascular resistance by altering VSM tone.¹ This important

function may be altered by clinical conditions that cause ischemic-reperfusion injury and inflammation of the vasculature.² Cell death during ischemia-reperfusion and inflammation may result from necrosis or increased apoptosis (programmed cell death).³ Pro-inflammatory cytokines, such as tumor necrosis factor α (TNF- α), interleukin 1 (IL-1 β), and interferon (IFN- γ) are released during reperfusion injury and induce apoptosis.⁴ Increased apoptosis results in endothelial cell up-regulation of adhesion molecules resulting in hyperadhesiveness of leukocytes and release of oxygen derived radicals, which causes necrosis of the endothelium.⁵ Apoptosis of VSM cells in regions of arteriosclerotic lesions may impede repair and precede plaque rupture.⁶ The final extent of endothelial and VSM injury after ischemia-reperfusion injury is dependent on the degree of apoptosis in addition to necrosis.⁷ Therefore, protection of the endothelium and VSM by inhibiting apoptosis and/or necrosis has important clinical implications in treatment of diseases associated with ischemic-reperfusion injury and inflammation.

In 1986, Murry *et al.*⁸ reported that a 5-min period of ischemia decreased the injury produced by a subsequent ischemic-reperfusion insult. Since then, studies have indicated that ischemic preconditioning (IPC) protects myocardial cells *in vivo*.⁹ Studies have also demonstrated that there are multiple ways to induce preconditioning similar to that seen with IPC, including transient elevations of calcium,^{10,11} heat shock,^{12,13} the administration of adenosine,¹⁴ opioids,¹⁵ protein kinase C (PKC) activators,¹⁶ adenosine triphosphate-sensitive potassium (K_{ATP}) channel agonists,¹⁷⁻¹⁹ and inhalational anesthetics.^{20,21} Volatile anesthetics appear to mimic IPC and have been shown to decrease myocardial infarction and preserve coronary endothelial function in coronary ligation models.^{20,21} Protective effects of continuously administered volatile anesthetics also have been demonstrated in cell culture models of induced cell death and apoptosis. Johnson *et al.*²² demonstrated that isoflurane and halothane decrease cell death in endothelial cells after exposure to hydrogen peroxide (H_2O_2). Similarly, Zaugg *et al.*²³ showed that volatile anesthetics inhibit norepinephrine-induced apoptosis in myocytes. Although the mechanisms are complicated and not entirely understood, there is evidence that anesthetics and IPC protect cells in both *in vivo* and *in vitro* models by regulation of K_{ATP} channels²⁴⁻²⁶ and PKC.^{27,9}

While continuously administered volatile anesthetics appear to be protective in cell culture,^{22,23} the effects of

* Post Doctoral Fellow, † Resident, Department of Anesthesiology, ‡ Associate Professor, Departments of Anesthesiology and Pediatrics, § Professor, Departments of Anesthesiology and Biomedical Engineering.

Received from the Departments of Anesthesiology, Pediatrics, and Biomedical Engineering, University of Virginia Health System, Charlottesville, Virginia. Submitted for publication November 28, 2001. Accepted for publication February 23, 2002. Supported by the Department of Anesthesiology, University of Virginia Health System, Charlottesville, Virginia. Presented at the annual meeting of the American Society of Anesthesiologists, New Orleans, Louisiana, October 15, 2001.

Address reprint requests to Dr. Rich: Department of Anesthesiology, PO Box 800710, University of Virginia Health System, Charlottesville, Virginia 22908-0710. Address electronic mail to: gfr2f@virginia.edu. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

preconditioning or anesthetic pretreatment on cell death and viability in cell culture have not been studied extensively. Cytokines have been shown to induce apoptosis in cell culture⁴; therefore, we used this model to test the hypothesis that a brief period of isoflurane may protect VSM cells and endothelial cells. We investigated whether 30 min of pretreatment with 1.5% isoflurane inhibits cytokine-induced cell death and increases cell viability in rat VSM and human umbilical vascular endothelial cells (HUVEC). Since previous *in vivo* studies have indicated that the timing of IPC and anesthetic preconditioning is important,²⁶ we evaluated the timing required for anesthetic administration to attenuate cytokine-induced cell death. To understand the mechanisms, we evaluated whether the effects of isoflurane pretreatment were similar to those with halothane, whether the pretreatment effects also occurred with H₂O₂-induced cell death, and whether PKC and/or K_{ATP} are involved in the mechanisms associated with anesthetic pretreatment.

Materials and Methods

Cell Cultures

Rat aorta VSMs were donated by Dr. L. Palmer, Ph.D. (Associate Professor, Departments of Anesthesiology and Pediatrics, University of Virginia Health System, Charlottesville, Virginia). The HUVECs were a gift from Dr. Tom G. O'Brig, Ph.D. (Department of Internal Medicine and Immunology, University of Virginia Health System). The SMCs were grown in Dulbecco's minimal essential growth medium (Gibco, Rockville, MD) supplemented with 10% fetal bovine serum (FBS). HUVECs were grown in medium bought from the Roswell Park Memorial Institute (Buffalo, New York), supplemented with 50 µg/ml endothelial mitogen (Biomedical Technologies, Fishers, IN), 29.2 mg/ml L-glutamine, 76 µg/ml heparin (Sigma, St. Louis, MO), and 20% FBS. The cytokines TNF-α, IFN-γ, and IL-1β were purchased from R&D Systems (Minneapolis, MN) and diluted into phosphate-buffered saline (PBS; Gibco) containing 0.1% bovine serum albumin (BSA; Sigma). The cells, passages 5-9, were plated at a density of 4 × 10⁵/ml.

Pretreatment and Cytokine Treatment

The initial experimental groups were pretreated with 1.5% isoflurane for 30 min in an airtight chamber before incubation with the cytokines at 37°C for periods of 24, 48, 72, or 96 h. Assessment was not possible for HUVECs at 96 h because of the need to change the medium for control cell survival. Isoflurane, 1.5%, was administered with an agent-specific vaporizer using a gas mixture of 95% O₂ and 5% CO₂ at a flow rate of 5 l/min. The isoflurane concentration in the chamber was measured with a gas analyzer (Datex Ohmeda, Finland). When the concentration remained constant at 1.5%, the chamber

inlet and outlet were closed, and the chamber was placed in the incubator at 37°C for 30 min. The medium was then replaced with 1 ml fresh medium in which the cytokines were dissolved at the following concentrations: 0.1 ng/ml TNF-α, 5.0 ng/ml IFN-γ, and 5.0 ng/ml IL-1β. The cultures were incubated with no cytokines (negative control), one cytokine, or combinations of two or three cytokines. After the different incubation periods (24, 48, 72, or 96 h), the cells were washed twice with PBS (Gibco) and were then further treated according to the specific assay protocols

Measurement of Cell Survival and Viability

Dead cells are incapable of excluding trypan blue; therefore, the uptake of this dye was used as a marker of cell death. Trypan blue (10 µl) was mixed with 50 µl cell suspension. For each sample, at least 100 cells per field were counted with a hemocytometer under a light microscope. Cell survival is the number of total cells minus the number of dead cells.

Cell viability was determined by proliferation assay (Chemicon International, Temecula, CA) and is based on cleavage of the tetrazolium salt WST-1 to formazan by cellular mitochondrial dehydrogenase as described by Krczeniewski *et al.*²⁸ Briefly, 100 µl of each sample was incubated with 10 µl of assay reagent in a 96-well plate for 2 h. The absorbance was measured with a spectrometer using a 450-nm filter. The viability of each sample was calculated as percentage of the absorbance of the sample cultured without cytokines.

Assessment of DNA Fragmentation Assays by TUNEL and ELISA

An apoptag plus peroxidase detection kit (Intergen, Purchase, NY) was used to detect DNA fragmentation, by means of nick-end labeling (TUNEL) fragmented DNA with biotinylated deoxynucleotide triphosphate (dUTP) as previously described by Wang *et al.*²⁹ Briefly, the cells were washed and treated with 20 µg/ml protein kinase K (Sigma). Endogenous peroxide was blocked by covering the sections with 3% H₂O₂ (Sigma). The biotinylated dUTP was applied to the slides, which then were incubated in a humidified chamber at 37°C for 1 h. The labeled DNA fragments were then stained with an anti-digoxigenin monoclonal antibody, and the cells were counterstained by immersion of the slides in 0.5% methyl green (Sigma). For each sample, two slides were made, and the total of stained cells were counted.

The enzyme-linked immunosorbent assay kit (Roche Molecular Biochemicals, Mannheim, Germany) was used to characterize cell death by measurement of 5'-Bromo-2'-deoxy-Uridine (BrdU)-labeled DNA fragments in culture supernatants and cell lysate. Proliferating VSMs were incubated with BrdU, which is incorporated into the genomic DNA. The cells were exposed to all three cytokines or H₂O₂ (1 mM) for periods of 2, 4, 6, and 8 h.

Aliquots (100 μ l) from the supernatant of each sample were transferred to an anti-BrdU antibody-coated 96-well plate. The remaining cells were lysed with a solution containing ethylenediaminetetraacetic acid, BSA, and poloxyethylenesorbitan. The lysate was also transferred into a BrdU antibody-coated well plate. After incubation for 90 min at room temperature, the samples in both plates were heated by microwave irradiation to denature DNA. Anti-BrdU peroxidase conjugate was added followed by a 90-min incubation period at room temperature. The samples were incubated on a shaker in darkness with 3,3'-5,5'-tetra-methyl-benzidine until fluorescence developed. This reaction was stopped by 5.5% sulfuric acid, and the absorbance was measured with a spectrometer at 450 nm.

Statistical Analysis

Cell counting was performed in a blinded manner. Comparisons between groups ($n = 6$) were made with an analysis of variance and a Tukey *post hoc* test. Statistical analysis was performed with SigmaStat 2.0 (Jandel Scientific Software, San Rafael, CA). Data are presented as mean \pm SD. $P < 0.05$ was considered significant.

Results

Effect of Isoflurane Pretreatment on Cytokine-induced Cell Death

The objective of the first experiment was to determine the effect of 30 min of pretreatment with 1.5% isoflurane on cell survival and viability. Isoflurane, 1.5%, was administered for 30 min in VSM and HUVEC cultures immediately prior to replacing the media with media in which TNF- α , IFN- γ , and IL-1 β alone, in combinations of two, and all three together, were dissolved. Control cultures not exposed to isoflurane or cytokines showed less than 10% cell death and greater than 90% viability over 72 h for HUVECs and 96 h for VSMs. Isoflurane had no effect on cell viability or cell survival in cultures not exposed to cytokines. Cytokines decreased cell viability and cell survival with all combinations of cytokines and for all time periods. The effects of the cytokines were increased at longer time periods and in the presence of greater combinations of cytokines.

In VSMs, pretreatment with 1.5% isoflurane for 30 min immediately prior to cytokine increased cell survival and increased cell viability for all multiple combinations of cytokines at all time periods by 25–60%. In single cytokine experiments using VSMs, isoflurane pretreatment increased cell survival at all time points except for IL-1 β at 24 h, TNF- α at 48 h, and IFN- γ at 72 h and increased cell viability for all time points except TNF- α alone and IL-1 β alone at 24 h and TNF- α at 48 h. In HUVECs, isoflurane pretreatment increased cell survival in the presence of IFN- γ , IL-1 β , and all three cytokines in com-




Fig. 1. The effect of isoflurane (1.5%) delivered for 30 min immediately prior to exposure to cytokines on cell survival (A) and viability (B) after 96 h in cultured vascular smooth muscle cells. Groups are no cytokines (neg), tumor necrosis factor α (TNF), interferon γ (IFN), interleukin 1 β (IL1), or combinations of two or three of these. # Denotes that isoflurane significantly ($P < 0.05$) increased cell survival and cell viability compared to cytokines alone (control). Data are mean \pm SD.

bination at 24 h, for all cytokine combinations at 48 h, and all combinations except TNF- α alone at 72 h by 10–45%. Cell viability was increased for all combinations except IL-1 β at 72 h. Isoflurane protection is illustrated at 96 h for VSMs (fig. 1) and 72 h for HUVECs (fig. 2).

The Dose-Response Effects of Isoflurane Pretreatment

Because of similar results with most parameters in the initial experiment, the remaining experiments were evaluated only in the presence of all three cytokines (TNF- α , IFN- γ , and IL-1 β) and to the time periods at 24 and 72 h. To determine the dose-response effects of isoflurane pretreatment, VSMs and HUVECs were pretreated with 0.75, 1.5, or 2.5% isoflurane for 30 min prior to cytokines.

In VSMs, there was no significant difference between the effects of 0.75, 1.5, and 2.5% isoflurane pretreatment on cell viability and cell survival at either 24 or 72 h. In HUVECs, pretreatment with 0.75% isoflurane had signif-




Fig. 2. The effect of isoflurane (1.5%) delivered for 30 min prior to exposure to cytokines on cell survival (A) and viability (B) after 72 h in human umbilical endothelial cells. Results are for 72 h and are in triplicate. Groups are no cytokines (neg), tumor necrosis factor α (TNF), interferon γ (IFN), interleukin 1β (IL1), or combinations of two or three of these. # Denotes that isoflurane significantly ($P < 0.05$) increased cell survival and cell viability compared to cytokines alone (control). Data are mean \pm SD.

icantly less effect on cell survival and viability compared with 1.5%; however, the effect of 0.75% was greater than no isoflurane pretreatment. The effects of 1.5% and 2.5% isoflurane were equal at both 24 and 72 h (fig. 3, survival only).

The Effect of Timing of the Isoflurane Treatment Relative to the Cytokine Exposure

To determine whether the timing of 1.5% isoflurane administration either prior to or immediately after cytokines was critical, the time between the end of the isoflurane pretreatment and the cytokine exposure was extended to 1, 2, and 4 h in VSMs. It was also determined whether isoflurane treatment was effective if administered at the same time or when initiated 30 min after cytokine exposure.

In VSMs, cell survival and viability were the same whether 1.5% isoflurane pretreatment was for 30 min immediately prior to the cytokines or whether the time between the end of the 30 min of isoflurane pretreat-

ment and the initiation of cytokine exposure was extended to 1 h. After a 2-h delay, the protective effect of isoflurane was less than after a 1-h delay but still significantly greater than no pretreatment. After a 4-h delay, the isoflurane pretreatment had no protective effect on cell survival or viability.

When exposure to isoflurane and the cytokines were initiated simultaneously, the effects on cell survival and viability were the same as with the 30-min isoflurane treatment prior to cytokine exposure. In contrast, isoflurane did not affect cell survival or viability when it was initiated 30 min after cytokine exposure. Results for these timed experiments were the same for 24 and 72 h (fig. 4, 72-h survival only).

The Effect of Halothane versus Isoflurane Pretreatment on Cytokines and H₂O₂-induced Cell Death

To determine whether halothane has similar effects as isoflurane, VSMs were exposed to 1.5% isoflurane or 1.2% halothane for 30 min prior to cytokine exposure. To determine whether isoflurane protection and halothane protection were similar in H₂O₂-induced cell death, pretreatment for 30 min was evaluated in VSMs containing H₂O₂ (1 mM).

In cytokines, 1.2% halothane significantly increased VSM cell survival and increased cell viability to the same extent as did isoflurane at 24 and 72 h (data not shown). H₂O₂ significantly decreased cell survival and viability. In H₂O₂, 1.5% isoflurane and 1.2% halothane had the same protective effect on cell survival and viability at 24 and 72 h (fig. 5, 72-h survival only).

The Effect of Glibenclamide and Staurosporine on Isoflurane Pretreatment

We investigated whether glibenclamide, which can inhibit K_{ATP} channels, and staurosporine, which can inhibit PKCs, alter the effects of anesthetic pretreatment in VSMs and HUVECs. Pretreatment experiments with 1.5% isoflurane were performed in the presence of glibenclamide (10 μ M) or staurosporine (4 nM) in cultures exposed to all three cytokines or H₂O₂ (1 mM).

In VSMs and HUVECs, glibenclamide and staurosporine had no effect on cell survival or viability in cells not exposed to cytokines or H₂O₂ in the presence or absence of isoflurane. Conversely, the pretreatment effects of isoflurane in VSM exposed to cytokines were attenuated by glibenclamide and staurosporine at 24 and 72 h. In HUVECs, glibenclamide and staurosporine completely abolished the pretreatment effect of isoflurane at 24 and 72 h (fig. 6, 72 h only). Likewise, the effects of isoflurane and halothane on cell survival and viability were abolished with glibenclamide and staurosporine in H₂O₂-induced cell death at both 24 and 72 h (fig. 5, 72-h survival only).




Fig. 3. The dose response of isoflurane delivered for 30 min prior to exposure to cytokines on cell survival in vascular smooth muscle cells (VSM) and human umbilical vein endothelial cells (HUVEC). Results are for 24 and 72 h. # Denotes that isoflurane significantly ($P < 0.05$) increased cell survival compared to cytokines alone (control). * Denotes that 0.75% significantly ($P < 0.05$) increased cell survival compared to cytokines alone but less than 1.5%. Data are mean \pm SD.

DNA Fragmentation Studies

A TUNEL assay was used to evaluate the effects of 1.5% isoflurane pretreatment on DNA fragmentation in cytokine experiments. VSMs and HUVECs were evaluated after 72 h. Cytokines increased TUNEL staining by $78 \pm 1\%$ in HUVECs and $65 \pm 1\%$ in SMCs after 72 h. Isoflurane pretreatment almost completely abolished staining so that it only measured $8 \pm 1\%$ in SMCs and $7 \pm 1\%$ in HUVECs.

DNA fragmentation assay was measured every 2 h up to 8 h to determine whether there were differences between cytokines and H_2O_2 . In VSMs, cytokines increased lysate DNA fragmentation by 2.5-fold after 8 h but did not alter supernatant DNA fragmentation compared to controls, which is suggestive of apoptosis. In contrast, H_2O_2 increased supernatant DNA fragmentation by 3.8-fold after 8 h but did not alter lysate DNA fragmentation, which is suggestive of necrosis.





Fig. 4. The effect of timing of isoflurane (1.5%) delivered for 30 min (horizontal line) on cell survival in vascular smooth muscle cells exposed to cytokines for 72 h. Isoflurane was completed 4, 2, or 1 h or immediately prior to cytokines, administered simultaneously with cytokines, or initiated 30 min after cytokines. # Denotes that isoflurane significantly ($P < 0.05$) increased cell survival compared to cytokines alone. * Denotes isoflurane significantly ($P < 0.05$) increased cell survival compared to no isoflurane but less than when completed 1 h prior to cytokines. Cell survival of the groups in which isoflurane was administered 4 h prior or 30 min after cytokine exposure were not different from controls (not shown). Data are mean \pm SD.

Discussion

Previous *in vivo* studies indicated that anesthetic pretreatment protects the myocardium from ischemia-reperfusion injury, and *in vitro* studies indicated that continuous volatile anesthetic administration attenuates apoptosis and cell death.²¹⁻²³ Thus, we hypothesized that isoflurane pretreatment would protect VSMs and HUVECs from cytokine-induced cell death and increase cell viability. In this study, we showed that 30 min of pretreatment with 1.5% isoflurane increased cell survival and cell viability due to cytokine-induced injury for at least 96 h in VSMs and 72 h in HUVECs. Pretreatment with isoflurane and halothane also increased VSM and HUVEC cell survival and viability for at least 72 h after exposure to H_2O_2 .

The protective effects of isoflurane pretreatment from cytokine-induced cell death were evaluated at three different concentrations ranging from 0.75 to 2.5%. There was no significant difference in the protective effect among all three concentrations in VSMs; however, in HUVECs 1.5% and 2.5% provided greater protection than 0.75%. Lui *et al.*²⁷ demonstrated that pretreatment with concentrations as low as 0.2% isoflurane were sufficient

Fig. 5. The effect of isoflurane (1.5%) and halothane (1.2%) delivered for 30 min prior to exposure to hydrogen peroxide (H_2O_2) or no exposure to H_2O_2 (neg) on vascular smooth muscle cell survival after 72 h. # Denotes that isoflurane and halothane significantly ($P < 0.05$) increased cell survival compared to H_2O_2 without anesthetic. Staurosporine or glibenclamide abolished the effect of isoflurane. Data are mean \pm SD.




to protect cultured chick myocytes from ischemia-reperfusion injury. While we did not evaluate isoflurane concentrations below 0.75%, it is possible that similarly low concentrations may be protective in our model with VSMs. On the other hand, the decreased effect of 0.75% in HUVECs indicates that low concentrations may not be protective in all cell types. Our study indicates that concentrations above 1.5% may not offer additional protective effects. In fact, Johnson *et al.*²² reported that supraclinical concentration of isoflurane (4%) and halothane (2.7%) afforded less protection than clinical concentration during H_2O_2 -induced cell death. Clinically relevant concentrations of volatile anesthetics also are protective in animal models of myocardial ischemia-

reperfusion injury,²⁴ suggesting a close correlation between *in vivo* and *in vitro* results. Possible clinical relevance is further supported by evidence that clinical concentrations of isoflurane may decrease myocardial damage in humans undergoing coronary artery bypass surgery.²¹

Protection of VSMs by isoflurane (1.5%) and halothane (1.2%) pretreatment in cytokine- and H_2O_2 -induced injury were the same. Similarly, Zaugg *et al.*²³ showed that continuous administration of halothane and isoflurane inhibited norepinephrine-induced apoptosis to the same extent. In rat hearts, halothane and isoflurane have also been shown to have similar protective effects on myocardial injury.³⁰ However, other studies have shown that

Fig. 6. The effects of staurosporine and glibenclamide on 30 min of isoflurane (1.5%) pretreatment in cytokine-exposed cells or cells not exposed to cytokines (neg). Studies are in vascular smooth muscle cell and human umbilical endothelial cell culture after 72 h. # Denotes that isoflurane significantly ($P < 0.05$) increased survival. * Denotes that staurosporine and glibenclamide significantly ($P < 0.05$) attenuated the effects of isoflurane in vascular smooth muscle cells (VSMs). Staurosporine and glibenclamide abolished the effects of isoflurane in human umbilical vein endothelial cells (HUVECs) exposed to cytokines. Staurosporine and glibenclamide had no effect in control cultures. Data are mean \pm SD.



the effects of volatile anesthetics are not similar. Johnson *et al.*²² indicated that halothane was more protective than isoflurane in H₂O₂-induced cell death. The opposite result was demonstrated by Roscoe *et al.*,³¹ who indicated that isoflurane but not halothane induced protection of cultured human cardiomyocytes. Whether or not there are significant differences in protection from isoflurane and halothane are undoubtedly model and injury dependent; however, our study indicates that pretreatment with each increases cell survival and viability to the same extent in the presence of cytokines and H₂O₂ in VSMs and HUVECs.

The timing of the isoflurane administration was critical for increasing cell survival and viability in VSMs. We administered isoflurane either immediately prior to cytokines; 1, 2, and 4 h prior to cytokines; simultaneously with cytokines; or 30 min after cytokines. As long as the 30 min of isoflurane was completed within 1 h prior to the administration of cytokines or administered at the same time as the cytokines, the maximal protective effects were observed. The protective effect decreases when the time between the completion of isoflurane administration and cytokine exposure is extended to 2 h and is abolished at 4 h. When isoflurane treatment is initiated 30 min after the cytokine exposure, there is also no protective effect. These results indicate several important findings. First, isoflurane must be present at the time of or prior to the initiating events of cytokine-induced cell death. In other words, isoflurane is effective in inhibiting the initiation of the cytokine cascade but is ineffective in halting the cascade once initiated. Second, the mechanisms by which isoflurane prevents the initiation of the cytokine-induced cell death are lost after 2–4 h. Previous studies have not evaluated this time period; however, in dog hearts, anesthetics may be delivered as early as 30 min prior to ischemia.²⁶ The apparent 2- to 4-h window that we demonstrated is particularly interesting because once the cytokine cascade is inhibited, the increase in cell survival and viability lasts at least 96 h. Because the effects of isoflurane appear to be lost after 2–4 h, it is likely that this increase in long-term survival is explained by inhibition of the initiation of the cytokine cascade. The observation that simultaneous administration of isoflurane with cytokine exposure also was effective indicates that the protective mechanisms have a very rapid onset.

Previous *in vivo* studies have suggested that stimulation of K_{ATP} channels or PKC is the protective mechanism for IPC and anesthetic preconditioning against ischemic injury.³² Blockade of isoflurane and halothane protection by glibenclamide and staurosporine argues for involvement of similar mechanisms against cytokine- and H₂O₂-induced cell death in cultured VSMs and HUVECs. Similar protective effects of volatile anesthetics and associated mechanisms in our cell culture studies compared to *in vivo* studies may not be surprising since

ischemic-reperfusion injury is partly mediated by cytokine and H₂O₂ release.³ Importantly, because of the 4-h window, the effects of isoflurane and halothane on PKC and/or K_{ATP} channels appear to be transient. The precise mechanisms by which PKC and/or K_{ATP} channels may alter cell death are unclear, although the mechanisms appear to be effective against different cytokines alone and in combination and against H₂O₂. Because isoflurane must be present at the initiation of cytokines, factors which are responsible for delayed protection, such as elevated heat shock proteins, are unlikely to be involved in our study.^{12,13} Whether these conclusions are also true for IPC or anesthetic preconditioning *in vivo* remains to be studied.

It is well accepted that K_{ATP} channels are involved in the mechanistic pathway of both ischemic and anesthetic preconditioning in *in vivo* models of myocardial protection.^{24,26} Volatile anesthetics offer myocardial protection in coronary ligation models, an effect which is inhibited by blocking the K_{ATP} channels.²⁰ Similarly, K_{ATP} channels have been shown to be involved in protection of coronary endothelium during ischemia-reperfusion *in vivo*.³³ Initial studies focused on the role of sarcolemmal K_{ATP} channel activity in IPC, but since the isolation of a mitochondrial K_{ATP} channel, studies have shown that the mitochondrial channel is crucial to the protection elicited by IPC.^{18,19,34} Ockaili *et al.*¹⁷ reported that direct stimulation of the mitochondrial K_{ATP} channel produced protection from ischemic injury in rats while blockade of this channel eliminated the protective effect. We evaluated only the effects of nonspecific K_{ATP} channel blockade with glibenclamide, and, although it is likely that mitochondrial K_{ATP} are more importantly involved, further studies will be required to determine whether the same applies to isoflurane and halothane protection during cytokine- and H₂O₂-induced cell death.

Activation of PKC appears to be a critical step in preconditioning because the phenomenon can be induced by PKC activators or blocked by PKC inhibitors. Wang *et al.*¹⁰ have shown that stimulation of PKC provides protection against Ca²⁺ overload injury in isolated rat hearts. Zhao *et al.*³⁵ demonstrated that transduced, basally active PKC- δ conferred protection from simulated ischemic injury in cell culture. Blockade of PKC abolishes the protective effects on various forms of preconditioning, including Ca²⁺ preconditioning and direct activation of mitochondrial K_{ATP} channels in isolated rat hearts.¹⁰ There are numerous PKC isozymes, of which the ϵ isozyme has recently been shown to be critical in IPC.³⁷ Staurosporine is a nonspecific antagonist of a PKC, and further evaluation will be required to determine the role of specific PKC isoenzymes in the VSM and HUVEC cell cultures.

It has been hypothesized that PKC is part of a linear cascade with the mitochondrial K_{ATP} channel as the end

effector. Sato *et al.*¹⁶ demonstrated that increased mitochondrial K_{ATP} channel opening is protective during an ischemic injury and can be elicited through priming the channel with PKC activity or by direct stimulation of the channel. This linear cascade, however, is challenged by the findings both *in vitro* and *in vivo* that blockade of PKC activity with simultaneous direct stimulation of mitochondrial K_{ATP} channels does not lead to any protective effect.^{10,36} This is supported by our study because if isoflurane increases K_{ATP} channel activity, blockade of PKC by staurosporine should not have abolished the protective effect. Based on these findings, it appears that PKC and mitochondrial K_{ATP} channels are involved in a network more complicated than a simple linear cascade. While previous studies have indicated that these mechanisms apply to ischemic-reperfusion injury, our study suggests that these mechanisms also may be involved in cellular protection from cytokines and H_2O_2 .

The observation in our study that isoflurane and halothane inhibit cell death in both cytokine- and H_2O_2 -induced injury may be important. Evaluation of DNA fragments in lysate *versus* supernatant suggests that cytokines induce apoptosis while H_2O_2 causes necrosis. Differentiation of apoptosis and necrosis is complex and is not conclusive by DNA fragmentation assays alone; however, our finding is consistent with other studies that have used additional methodologies.^{4,23} These results are also consistent with other studies that indicate that volatile anesthetics inhibit both apoptosis and necrosis.^{22,23} The role of PKC in inhibiting both cytokine-induced apoptosis and H_2O_2 -induced necrosis is supported by a study by Lui *et al.*,³⁷ who showed that IPC activation of PKC ϵ attenuates both apoptosis and necrosis in cardiomyocytes. Furthermore, evidence of the role of isoflurane in inhibition of apoptosis and/or necrosis is suggested by our results indicating staining of DNA fragmentation by TUNEL assay is greatly attenuated in VSMs and HUVECs.

In conclusion, this study demonstrates that 30 min of pretreatment with 1.5% isoflurane decreases cell death due to cytokine-induced injury for at least 96 h in VSMs and 72 h in HUVECs. The timing of isoflurane administration is critical as isoflurane inhibits the initiation of the cytokine cascade but does not appear to halt it. These initial inhibitor studies suggest involvement of PKC and K_{ATP} channels in isoflurane and halothane protection against both cytokine- and H_2O_2 -induced cell death of VSMs and HUVECs.

The authors thank Tom G. Obrig, Ph.D. (Professor of Internal Medicine and Immunology, University of Virginia Health System, Charlottesville, Virginia), and Julianne J. Sando, Ph.D. (Professor of Anesthesiology), for their gift of cells, their help, and their advice.

References

- Moncada S, Higgs: The L-arginine-nitric oxide pathway. *N Eng J Med* 1993; 329:2002-12
- Rubino A, Yellon D: Ischemic preconditioning of the vasculature an overlooked phenomenon for protecting the heart. *Trends Pharmacol Sci* 2000; 21: 225-30
- Dimmeler S, Hermann C, Zeiher AM: Apoptosis of endothelial cells: Contribution of the pathophysiology of atherosclerosis. *Eur Cytokine Netw* 1998; 9:697-8
- Geng YJ, Wu Q, Matuszyski M, Hansson GK, Libby P: Apoptosis of vascular smooth muscle cells induced by *in vitro* stimulation with interferon- γ , tumor necrosis factor- α , and interleukin-1 β . *Arterioscler Thromb Vasc Biol* 1996; 16: 19-27
- Kvietys PR, Granger DN: Endothelial cell monolayers as a tool for studying microvascular pathology. *Am J Physiol* 1997; 273:G1189-99
- Walsh K, Smith RC, Hyo-Soo K: Vascular cell apoptosis in remodeling, restenosis, and plaque rupture. *Circ Res* 2000; 87:184-8
- Li D, Yang B, Metha JL: Tumor Necrosis Factor- α enhances hypoxia-reoxygenation-mediated apoptosis in cultured human coronary artery endothelial cells: Critical role of protein kinase C. *Cardiovasc Res* 1999; 42:805-13
- Murry CE, Jennings RB, Reimling KA: Preconditioning with ischemia: A delay of lethal cell injury. *Circulation* 1986; 74:1124-36
- Speechly-Dick ME, Grover GJ, Yellon DM: Does ischemic preconditioning in the human involve protein kinase and the ATP-dependent K^+ -channel? Studies of contractile function after stimulated ischemia in an atrial *in vitro* model. *Circ Res* 1995; 77:1030-5
- Wang Y, Ashraf M: Role of protein kinase C in mitochondrial KATP channel, mediated protection against Ca^{2+} overload injury in rat myocardium. *Circ Res* 1994; 84:1156-65
- Miyawaki H, Astraf M: Ca^{2+} as a mediator of ischemic preconditioning. *Cir Res* 1997; 80:790-9
- Hoag JB, Qian Y-Z, Nayeem MA, D'Angelo M, Kukreja RC: ATP-sensitive potassium channel mediates delayed ischemic protection by heat stress in rabbit heart. *Am J Physiol* 1997; 42:H861-68
- Pell TJ, Yellon DM, Goodwin RW, Baxter GF: Myocardial ischemic tolerance following heat stress is abolished by ATP-sensitive potassium channel blockade. *Cardiovasc Drugs Ther* 1997; 11:679-86
- Cleveland JC Jr, Meldrum DR, Rowland RT, Banerjee A, Harken AH: Adenosine preconditioning of human myocardium is dependent upon the ATP-sensitive K^+ channel. *J Mol Cell Cardiol* 1997; 29:175-82
- Liang BT, Gross GJ: Direct preconditioning of cardiac myocytes via opioid receptors and KATP channels. *Circ Res* 1998; 83:110-4
- Sato T, O'Rourke B, Marban E: Modulation ATP-dependent K^+ channels by protein kinase C. *Cir Res* 1998; 83:110-4
- Ockaili R, Emani VR, Okubo S, Brown M, Krottapalli K, Kukreja RC: Opening of mitochondrial K_{ATP} channel induces early and delayed cardioprotective effect: Role of nitric oxide. *Am J Physiol* 1999; 277:H2425-34
- Liu Y, Sato T, O'Rourke B, Marban E: Mitochondrial ATP-dependent potassium channels: Novel effectors of cardioprotection. *Circulation* 1998; 97:2463-9
- Baines CP, Lui GS, Birincioglu M, Critz D, Cohen MV, Downey JM: Ischemic preconditioning depends on interaction between mitochondrial K_{ATP} channels and actin cytoskeleton. *Am J Physiol* 1999; 276:H1361-8
- Novalija E, Fujita S, Kampine JP, Stowe DF: Sevoflurane mimics ischemic preconditioning effects on coronary flow and nitric oxide release in isolated hearts. *ANESTHESIOLOGY* 1999; 91:701-12
- Belhomme D, Peynet J, Louzy M, Launay J-M, Kitakaze M, Menasché P: Evidence for preconditioning by isoflurane in coronary artery bypass graft surgery. *Circulation* 1999; 100(suppl II):II340-4
- Johnson ME, Sill JC, Uhl CB, Halsey TJ, Gores GJ: Effect of volatile anesthetics on hydrogen peroxide-induced injury in aortic and pulmonary arterial endothelial cells. *ANESTHESIOLOGY* 1996; 84:103-16
- Zaugg M, Jamali NZ, Luchinetti E, Shafiq SA, Siddiqui MAQ: Norepinephrine-induced apoptosis is inhibited in adult rat ventricular myocytes exposed to volatile anesthetics. *ANESTHESIOLOGY* 2000; 93:209-13
- Crystal GJ, Gurevicius J, Salem MR, Zhou X: Role of adenosine triphosphate-sensitive potassium channels in coronary vasodilatation by halothane, isoflurane, and enflurane. *ANESTHESIOLOGY* 1997; 86:448-58
- Kersten JR, Schemeling TJ, Pagel PS, Gross, GJ, Warltier DC: Mechanism of myocardial protection by isoflurane: Role of adenosine triphosphate-regulated potassium (K_{ATP}) channels. *ANESTHESIOLOGY* 1996; 85:794-807
- Kersten JR, Schemeling TJ, Pagel PS, Gross, GJ, Warltier DC: Isoflurane mimics ischemic preconditioning via activation of K_{ATP} channels: Reduction of myocardial infarct size with an acute memory phase. *ANESTHESIOLOGY* 1997; 87: 361-70
- Lui H, Zhu X, Mcpherson B, Shao Z, Li C, Yao Z: Isoflurane blocks ischemia-reperfusion induced apoptosis via protein kinase C in myocytes. *Anesth Analg* 2001; 92:S1-363
- Korzeniewski C, Calewood DM: An enzyme-release assay for natural cytotoxicity. *J Immunol Methods* 1983; 64:313-20
- Wang J, Keiser J: Molecular characterization of rabbit CPP32 and its function in vascular smooth muscle cell apoptosis. *Am J Physiol* 1998; 274: H1132-40
- Boutros A, Wang J, Capuano C: Isoflurane and halothane increase adenosine triphosphate preservation, but do not provide additive recovery of function after ischemia, in preconditioned rat hearts. *ANESTHESIOLOGY* 1997; 86:109-17

31. Roscoe AK, Christensen JD, Lynch C III: Isoflurane, but not halothane, induces protection of human myocardium *via* adenosine A1 receptors and adenosine triphosphate-sensitive potassium channels. *ANESTHESIOLOGY* 2000; 92: 1692-701
32. Wang Y, Kyoji H, Ashraf M: Activation of mitochondrial ATP-sensitive K⁺ channel for cardiac protection against ischemic injury is dependent on protein kinase C activity. *Circ Res* 1999; 85:731-41
33. Ismaeil MS, Tkachenko I, Gamperl AK, Kickey RF, Cason CA: Mechanisms of isoflurane-induced preconditioning in rabbits. *ANESTHESIOLOGY* 1999; 90:812-21
34. Pomerantz BJ, Robinson TN, Heimbach JK, Calkins CM, Miller SA, Banerjee A, Harken AH: Selective mitochondrial KATP channel opening controls human myocardial preconditioning: Too much of a good thing? *Surgery* 2000; 128: 368-73
35. Zhao J, Renner O, Wightman L, Sugden P, Stewart L, Miller A, Latchman D, Marber M: The expression of constitutively active isoforms of protein kinase C to investigate preconditioning. *J Biol Chem* 1998; 273:23072-9
36. Gaudette GR, Krukenkamp IB, Saltman AE, Horimoto H, Levitsky S: Preconditioning with PKC and ATP-sensitive potassium channels: A codependent relationship. *Ann Thorac Surg* 2000; 70:602-8
37. Lui H, McPherson BC, Yao Z: Preconditioning attenuates apoptosis and necrosis: Role of protein kinase C ϵ and δ isoforms. *Am J Physiol* 2001; 281: H404-10