

Isoflurane Decreases ATP Sensitivity of Guinea Pig Cardiac Sarcolemmal K_{ATP} Channel at Reduced Intracellular pH

Anna Stadnicka, Ph.D.,* Zeljko J. Bosnjak, Ph.D.†

Background: Volatile anesthetics can protect the myocardium against ischemic injury by opening the adenosine triphosphate (ATP)-sensitive potassium (K_{ATP}) channels. However, direct evidence for anesthetic-channel interaction is still limited, and little is known about the role K_{ATP} channel modulators play in this effect. Because pH is one of the regulators of K_{ATP} channels, the authors tested the hypothesis that intracellular pH (pHi) modulates the direct interaction of isoflurane with the cardiac K_{ATP} channel.

Methods: The effects of isoflurane on sarcolemmal K_{ATP} channels were investigated at pHi 7.4 and pHi 6.8 in excised inside-out membrane patches from ventricular myocytes of guinea pig hearts.

Results: At pHi 7.4, intracellular ATP (1–1,000 μ M) inhibited K_{ATP} channels and decreased channel open probability (P_o) in a concentration-dependent manner with an IC_{50} of $8 \pm 1.5 \mu$ M, and isoflurane (0.5 mM) either had no effect or decreased channel activity. Lowering pHi from 7.4 to 6.8 enhanced channel opening by increasing P_o and reduced channel sensitivity to ATP, with IC_{50} shifting from 8 ± 1.2 to $45 \pm 5.6 \mu$ M. When applied to the channels activated at pHi 6.8, isoflurane (0.5 mM) increased P_o and further reduced ATP sensitivity, shifting IC_{50} to $110 \pm 10.0 \mu$ M.

Conclusions: Changes in pHi appear to modulate isoflurane interaction with the cardiac K_{ATP} channel. At pHi 6.8, which itself facilitates channel opening, isoflurane enhances channel activity by increasing P_o and reduces sensitivity to inhibition by ATP without changing the unitary amplitude of single channel current or the conductance. These results support the hypothesis of direct isoflurane- K_{ATP} channel interaction that may play a role in cardioprotection by volatile anesthetics.

THE adenosine triphosphate (ATP)-sensitive potassium (K_{ATP}) channels are thought to play an important role in anesthetic preconditioning, a protection afforded by volatile anesthetics against ischemia and reperfusion injury.^{1–6} Although cellular mechanisms of this protection remain the focus of many investigations, direct evidence for interaction of volatile anesthetics with the K_{ATP} channel is still limited.^{7,8} Such an interaction is clinically important because volatile anesthetics may mimic or enhance the protective mechanism of K_{ATP} channel opening.

A characteristic property of K_{ATP} channels is their sensitivity to inhibition by physiologic concentrations of

intracellular ATP ([ATP]_i) that is controlled by a number of cytosolic factors including nucleotide diphosphates (ADP), phospholipids such as phosphatidylinositol 4,5-bisphosphate (PIP₂), and intracellular protons. Intracellular pH (pHi) modulates the activity of native K_{ATP} channels in pancreatic β cells,⁹ skeletal muscle,^{10,11} heart,^{12–16} and cloned K_{ATP} channels in *Xenopus* oocyte expression system.^{17–19} The acidic species of K_{ATP} channel modulators have been implicated in the mechanism of pH-dependent regulation of sensitivity to ATP.²⁰ An interaction of intracellular protons with ATP in regulating channel activity has been also suggested recently.²¹ High sensitivity to activation by intracellular protons implies an important role for K_{ATP} channels in regulation of cellular excitability during various metabolic stresses that often are accompanied by a decrease in pHi. However, whether pHi is a factor in volatile anesthetic- K_{ATP} channel interaction has not been established.

We have recently reported that during whole cell or cell-attached patch clamp conditions isoflurane potentiates the cardiac K_{ATP} channel current (I_{KATP}) and increases open probability (P_o) of channels previously activated by an uncoupler of oxidative phosphorylation, 2,4-dinitrophenol, and the K_{ATP} channel opener, pinacidil.^{8,22} Potentiation, however, did not occur in the inside-out patches where at pHi 7.4 isoflurane either had no effect or decreased channel activity.^{7,8} Differential on-cell *versus* cell-free effects of isoflurane suggested that other intracellular factors might be involved in anesthetic potentiation.

In the present study, we tested the hypothesis that pHi is one of the endogenous factors modulating isoflurane interaction with the cardiac K_{ATP} channel. Because intracellular acidosis, an effect characteristic of early ischemia, may also occur perioperatively before or during administration of the volatile anesthetic agents, we investigated whether decreasing pHi alters interaction of isoflurane with the K_{ATP} channel regarding modulation of channel P_o and sensitivity to inhibition by [ATP]_i.

Materials and Methods

Cell Isolation

After approval by the Institutional Animal Use and Care Committee of the Medical College of Wisconsin, single ventricular myocytes were isolated from guinea pig hearts by enzymatic dissociation with collagenase Type II, (Gibco/Invitrogen, Life Technologies, Grand Island, NY) and protease Type XIV, (Sigma, St Louis, MO) as

* Assistant Professor of Anesthesiology, † Professor of Anesthesiology and Physiology.

Received from the Department of Anesthesiology, the Medical College of Wisconsin, Milwaukee, Wisconsin. Submitted for publication June 27, 2002. Accepted for publication September 23, 2002. Supported in part by the grant HL 34708-14 (Z. J. Bosnjak) from the National Institutes of Health, Bethesda, Maryland.

Address reprint requests to Dr. Stadnicka: Department of Anesthesiology, Medical College of Wisconsin, 8701 Watertown Plank Rd., Milwaukee, Wisconsin 53226. Address electronic mail to: astadnic@mcw.edu. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

reported previously.²³ Ventricular myocytes were stored in modified Tyrode solution, and only calcium-tolerant, rod-shaped cells with distinct cross-striations were used for experiments within 8 h after isolation.

Solutions

The modified Tyrode solution contained NaCl, 132 mM; KCl, 4.8 mM; $MgCl_2$, 1.2 mM; $CaCl_2$, 1 mM; HEPES, 10 mM; and glucose, 5 mM; at pH adjusted to 7.4 with NaOH.

For single channel recordings in the inside-out patch clamp configuration, the bath solution facing the intracellular side of membrane patches contained KCl, 140 mM; $MgCl_2$, 0.5 mM; EGTA, 1 mM; HEPES, 10 mM; and variable 0–1 mM K_2 -ATP, at pH 7.4 or pH 6.8 adjusted with KOH or HCl. The pipette solution facing the extracellular side of membrane patches contained KCl, 140 mM; $MgCl_2$, 0.5 mM; $CaCl_2$, 0.5 mM; and HEPES, 10 mM at pH 7.4 adjusted with KOH. All chemicals were purchased from Sigma (St. Louis, MO).

Isoflurane (Baxter Healthcare, Deerfield, IL) was delivered to the recording chamber in the bath solution. Anesthetic solution was prepared by adding a measured aliquot of isoflurane to a known volume of bath solution and dispersing it by sonication. This solution was then transferred into a gas-tight glass syringe reservoir to be delivered to the recording chamber by a gravity-fed perfusion system. Isoflurane was used at a clinically relevant concentration of 0.53 ± 0.04 mM ($n = 94$) equivalent to 1.06 vol% at 20–23°C, or 0.9 minimum alveolar concentration (MAC) in guinea pigs and humans. This concentration was chosen because at 1.0 MAC, isoflurane has been shown to protect the myocardium *in vivo* and *in vitro*^{3,6} and to enhance activity of whole cell I_{KATP} and single K_{ATP} channels.^{8,22} Isoflurane concentrations were determined by the headspace analysis method using a Shimadzu GC 8A flame ionization detection gas chromatograph (Shimadzu, Kyoto, Japan) from aliquots of the bath solution sampled directly from the recording chamber.

Single Channel Recordings and Analysis

Ventricular cells were placed in a RC-16 recording chamber (Warner, Hamden, CT) on the stage of an inverted IMT-2 microscope (Olympus, Tokyo, Japan). Single K_{ATP} channel activity was monitored in the inside-out configuration of the patch clamp technique at 20–23°C. Patch pipettes were pulled from borosilicate glass tubing (Garner Glass, Claremont, CA) with a horizontal PC-84 micropipette puller (Sutter Instruments, Novato, CA). The tips were heat-polished with an MF-83 microforge (Narishige, Tokyo, Japan). Pipettes had resistances of 7–12 M Ω when filled with the extracellular solution. After gigaseal formation, the inside-out patches were excised by rapidly pulling the pipette away from the cell. In this configuration, the intracellular side of the mem-

brane patch was directly exposed to the intracellular bath solution. Channel activity was recorded using a List EPC-7 amplifier (ALA Scientific Instruments, Westbury, NY) interfaced to a personal computer through a Digiata 1200B (Axon Instruments, Foster City, CA). Data were acquired using pClamp8 software (Axon Instruments, Foster City, CA). The current signal was filtered at 500 Hz through an 8-pole Bessel low-pass filter and sampled at 1 kHz. Single channel data were analyzed with pClamp8 software (Axon Instruments, Foster City, CA) and Origin6 software (OriginLab, Northampton, MA).

At symmetric 140 mM K^+ concentration, unitary outward current through single K_{ATP} channels was monitored at the transmembrane patch potential of +40 mV. The 60-s recordings were made at each experimental step. The K_{ATP} channels were identified by the single channel conductance, sensitivity to inhibition by [ATP]_i, and blockade by glibenclamide (1 μ M). A 50% threshold criterion was used for detecting the open state. Amplitude of single channel current was determined from the all-points amplitude histograms constructed from data segments of 60-s duration. Channel P_o was determined from the ratios of the area under the peaks in the all-points amplitude histograms fitted with a Gaussian function. The number (N) of channels in each patch was estimated during brief exposure to ATP-free internal solution (0 ATP) at the end of the experiments. P_o was calculated using the equation $P_o = [1 - (P_c)^{1/n}]$ where P_c is the channel closed state probability. For measurements of ATP sensitivity, P_o of each patch was normalized to P_o determined at 0 ATP to control for variations in P_o among patches. The sensitivity to ATP (1, 10, 50, 100, and 1,000 μ M) was determined at pHi 7.4 and pHi 6.8 in the absence or presence of isoflurane. The experimental protocols were completed within 10–12 min after patch excision. To minimize channel rundown, we used the Ca^{2+} -free and low Mg^{2+} intracellular solution and exposed each patch to 0 ATP only at the end of experimental protocols because even a brief exposure to ATP-free solution immediately after patch excision could accelerate rundown. Therefore, the number of channels in patches could have been underestimated in our study. Channel rundown occurred more frequently at pHi 7.4. Decreasing pHi to 6.8 tended to stabilize the channels and slow rundown. Recordings from patches exhibiting a significant rundown were excluded from analysis. For measurement of ATP sensitivity, the relationship between [ATP]_i and P_o was fitted by Hill equation:

$$\begin{aligned} \text{Normalized } P_o &= P_o/P_{o_{\max}} \\ &= 1/(1 + ([ATP]_i/IC_{50})^{nH}) \end{aligned}$$

where P_o is channel open probability at any test [ATP]_i; $P_{o_{\max}}$ is open probability at 0 [ATP]_i; IC_{50} is ATP concentration for half-maximal effect; and nH is Hill coefficient.




Fig. 1. Summary data for adenosine triphosphate (ATP) sensitivity of single K_{ATP} channels at intracellular pH (pHi) 7.4. Symbols and error bars are means \pm SEM. Each data point is a mean from four inside-out patches. Normalized mean open probability (Po) data obtained in control (open symbols) and in the presence of 0.5 mM isoflurane (closed symbols) are plotted against intracellular ATP ([ATP]i) concentrations. Solid line is a Hill fit (see Methods) to mean control data. The IC_{50} for ATP inhibition and the Hill coefficient are reported in Results. At pHi 7.4, isoflurane decreased Po independently of [ATP]i. The dotted line connects mean isoflurane data points.

Statistical Analysis

Data are presented as mean \pm SEM. Comparisons were made using paired or unpaired Student *t* test. Multiple group means were compared by analysis of variance with a Student-Newman-Keuls test. Differences with a two-tailed $P < 0.05$ were accepted as significant.

Results

The effects of isoflurane on the outward current through K_{ATP} channels were investigated at pHi 7.4 and pHi 6.8 in the inside-out patches from guinea pig ventricular myocytes at a symmetric 140 mM K^+ concentration and the patch potential of +40 mV.

ATP Sensitivity of Single K_{ATP} Channels and Isoflurane Effects at pHi 7.4

Membrane patches were excised into the intracellular solution containing 0.2 mM ATP. Multiple channel openings that appeared on patch excision decreased within 30–40 s, and thereafter only the activity of spontaneously operative channels was recorded. To assess ATP dependence of isoflurane effects, we first evaluated ATP sensitivity of the channel during control conditions at pHi 7.4. The following protocol was carried out at each tested [ATP]i: control at pHi 7.4, isoflurane at pHi 7.4, washout of isoflurane at pHi 7.4, and 0 ATP at pHi 7.4. Only one concentration of ATP was tested per patch. ATPi inhibited channel activity and decreased Po in a

concentration-dependent manner. Figure 1 shows summary data for [ATP]i-normalized Po relationship at pHi 7.4 in the control and during application of 0.5 mM isoflurane. Each data point is a mean from four patches. During control conditions, increasing [ATP]i caused a concentration-dependent decrease in Po. Fitting mean data to the Hill equation yielded an IC_{50} for ATP inhibition of $8 \pm 1.5 \mu\text{M}$ and a Hill coefficient (nH) of 0.6 ± 0.05 . When applied to the internal side of patches at pHi 7.4, isoflurane decreased Po at [ATP]i less than 50 μM but had no marked effect on Po at [ATP]i greater than 50 μM . Figure 2 shows the sample traces of single K_{ATP} channel activity recorded at pHi 7.4 in the control and during application of 0.5 mM isoflurane at 50 μM [ATP]i. In a patch containing five channels, isoflurane decreased channel activity in a reversible manner. As shown in figure 3, at 100 μM [ATP]i channel activity was much lower and little affected by isoflurane. The unitary amplitudes of 2.2 ± 0.1 pA (control) and 2.1 ± 0.1 pA (isoflurane) and the conductance were not altered by the anesthetic at pHi 7.4.

Isoflurane Effects on ATP Sensitivity at pHi 6.8

To test whether anesthetic- K_{ATP} channel interaction is modulated by pHi, the effects of isoflurane were examined by decreasing pHi from 7.4 to 6.8. The following protocol was carried out at each tested [ATP]i: control at pHi 7.4, control at pHi 6.8, isoflurane at pHi 6.8, washout of isoflurane at pHi 6.8, and 0 ATP at pHi 6.8. This protocol also included a control baseline at pHi 7.4 because decreasing pHi itself is known to enhance opening of the cardiac K_{ATP} channels.^{12–14,16} Only one concentration of ATP was tested per patch. Figure 4 shows recordings at 100 μM [ATP]i from a patch containing three active channels. Infrequent at pHi 7.4, channel opening increased markedly when decreasing pHi to 6.8. Application of 0.5 mM isoflurane at pHi 6.8 enhanced channel activity and further increased Po. Isoflurane effects were reversible during washout. Figure 5 shows summary data for [ATP]i-Po relationship obtained during the above condition where each patch was sequentially exposed to the internal solution at pHi 7.4 and pHi 6.8 and to 0.5 mM isoflurane at pHi 6.8. Each data point is a mean from six patches. At pHi 7.4, fitting the [ATP]i-Po relationship to the Hill equation yielded an IC_{50} of $8 \pm 1.2 \mu\text{M}$ and nH of 0.6 ± 0.04 . Decreasing pHi from 7.4 to 6.8 caused a rightward shift of the curve with IC_{50} of $45 \pm 5.6 \mu\text{M}$ and nH of 0.8 ± 0.1 . The IC_{50} value was approximately fivefold greater than that at pHi 7.4, and both values were different from each other at $P < 0.05$. When applied at pHi 6.8, isoflurane further decreased ATP sensitivity, and the curve shifted further to the right, yielding an IC_{50} of $110 \pm 10.0 \mu\text{M}$ and nH of 1.05 ± 0.12 . The IC_{50} value was more than twofold greater than that at pHi 6.8 alone, and the values were significantly different at $P < 0.05$. Neither decreasing




Fig. 2. Effect of isoflurane on single K_{ATP} channels at intracellular pH (pHi) 7.4. (*Upper*) Current traces recorded from an inside-out patch in the presence of $50 \mu\text{M}$ intracellular ATP ([ATP]_i) at patch potential of +40 mV. The patch contained five channels, as determined at 0 ATP. Dashed lines denote the closed state (C). Upward deflection indicates channel opening. Isoflurane (0.5 mM) applied in the intracellular solution reversibly inhibited channel activity and decreased open probability (P_o) (*Lower*) All-points amplitude histograms from the recordings above demonstrate that isoflurane did not affect the amplitude of unitary current.

pHi to 6.8 nor application of isoflurane at pHi 6.8 altered the amplitude of unitary current, which was $2.2 \pm 0.1 \text{ pA}$ at pHi 7.4, $2.3 \pm 0.1 \text{ pA}$ at pHi 6.8, and $2.3 \pm 0.1 \text{ pA}$ at pHi 6.8 with isoflurane. Single channel conductance remained in a range of 55–57 pS.

Discussion

Results from this study provide direct evidence for isoflurane inhibition of ATP sensitivity of cardiac K_{ATP} channels at reduced pHi. In the inside-out patches at pHi 6.8, isoflurane potentiates single K_{ATP} channels by increasing P_o and reduces channel sensitivity to ATP without affecting amplitude of unitary outward current and conductance.

Isoflurane alone did not activate whole cell $I_{K_{ATP}}$ in human atrial⁶ and guinea pig ventricular myocytes,²² however, isoflurane potentiated the whole cell $I_{K_{ATP}}$ preactivated by pinacidil or 2,4-dinitrophenol.²² In guinea pig ventricular myocytes, isoflurane increased P_o of K_{ATP} channels in the cell-attached but not in the inside-out membrane patches.⁸ By contrast, isoflurane decreased the activity of single K_{ATP} channels in the inside-out patches from rabbit ventricular myocytes.⁷ The pHi in all of these studies was kept at 7.4.

In the present study, in the inside-out patches and at pHi 7.4, isoflurane inhibited single K_{ATP} channel activity and decreased P_o at [ATP]_i less than $50 \mu\text{M}$ in a concentration-independent manner but had no effect on channel activity at [ATP]_i greater than $50 \mu\text{M}$. This finding confirms the results of Fujimoto *et al.*⁸, who reported lack of isoflurane effects on K_{ATP} channel activity in the inside-out patches at $300 \mu\text{M}$ [ATP]_i. We also found that isoflurane does not decrease ATP sensitivity of the channel at pHi 7.4. This observation differs from that of Han *et al.*⁷, who reported a decrease in ATP sensitivity after exposure of inside-out patches to isoflurane at pHi 7.4. Reasons for this discrepancy are not clear, but taking aside species (rabbit *vs.* guinea pig) differences, they could be related to differences in the experimental design. First, Han *et al.*⁷ examined the effects of isoflurane at the membrane potential of -70 mV and thus investigated the inward current through K_{ATP} channels, whereas our studies focused on the unitary outward current at the membrane potential of 0 mV (Fujimoto *et al.*⁸) and +40 mV (present study). This raises the question whether anesthetics may differentially modulate the inward *versus* outward conductance of the K_{ATP} channel. Second, our measurements were taken during the exposure to isoflurane, whereas Han *et al.*⁷ measured




Fig. 3. Effect of isoflurane on K_{ATP} channel activity at intracellular pH (pHi) 7.4 in the presence of $100 \mu\text{M}$ intracellular ATP ([ATP]i). (*Upper*) Shown are recordings from a patch containing two active channels as determined at 0 ATP. (C) denotes channel closed state, and upward deflection indicates the open state. At $100 \mu\text{M}$ [ATP]i, isoflurane (0.5 mM) had negligible effect on channel activity. Open probability (P_o) values are shown at each experimental step. (*Lower*) All-points amplitude histograms from recordings above show no changes in unitary current amplitude with isoflurane.

ATP sensitivity before and after application of isoflurane, as shown in their figure 1 (middle and lower) and figure 3A and B. Third, from their figures 1, 2A, and 3A it appears that ATP was absent during the patch exposure to isoflurane. Regardless of these differences, both studies have demonstrated that a volatile anesthetic, isoflurane, may directly interact with the sarcolemmal K_{ATP} channel.

Our results suggest that pHi may be one of the factors that modulate isoflurane-channel interaction. The K_{ATP} channels are sensitive to changes in pHi, and a decrease in pHi is a potent stimulus for their activation by the mechanism that involves a decrease in sensitivity to inhibition by ATP. The optimal effect occurs at pHi 6.8 to 6.5, and further decrease or increase in pHi leads to channel inhibition.^{12,15,17-19,21} For instance, intracellular acidification to pHi less than 6.5 inhibits cardiac K_{ATP} channels by inducing multiple subconductance levels.¹⁵ As anticipated, in our study decreasing pHi from 7.4 to 6.8 increased channel opening. This effect was the result of increased P_o and reduced ATP sensitivity, as reflected by the rightward shift in the [ATP]i - P_o curve with IC_{50} increasing from $8 \mu\text{M}$ to $45 \mu\text{M}$. Decreasing pHi to 6.8 did not affect the amplitude of unitary outward current, as also reported by others.¹² The IC_{50} values for ATP inhibition obtained by us at near physiologic and mild aci-

dotic pHi are in the range of previously reported concentrations.^{12,14} However, these values are not identical, and our Hill coefficients are lower. This is not surprising because the experimental conditions vary among studies, and it is well known that many factors may alter ATP sensitivity of K_{ATP} channels. These include the variations in experimental protocols and ionic conditions, the presence and concentration of monovalent and divalent ions (Ca^{2+} , Mg^{2+}) and glucose, differences in the range of pHi under study, outward or inward channel conductance under study, and species differences. In addition, pHi sensitivity of K_{ATP} channels may be modified by ATP^{21} and Mg^{2+} ions.

Our study demonstrated that mild intracellular acidosis modulates direct interaction of isoflurane with the K_{ATP} channel. At pHi 6.8, isoflurane potentiates channel activity by decreasing ATP sensitivity and shifting IC_{50} for ATP inhibition from $45 \mu\text{M}$ to $110 \mu\text{M}$. The mechanism by which intracellular acidosis modulates isoflurane- K_{ATP} channel interaction is unknown, and we can only be speculative on this point. It has been established that pH sensing is an inherent property of Kir6.2 subunits of the K_{ATP} channel.¹⁷ Three separate domains in the Kir6.2 protein—the N terminus, C terminus, and M2 domain—are involved in pH regulation, and the proton-sensing amino acid residues responsible for modulation of chan-




Fig. 4. Effect of decreasing intracellular pH (pHi) from 7.4 to 6.8 on K_{ATP} channel activity in the absence and presence of isoflurane. (Upper) Sixty-second recordings of channel activity at 100 μM intracellular ATP ([ATP]i) from an inside-out patch that was exposed sequentially to intracellular solution at pHi 7.4, at pHi 6.8, isoflurane at pHi 6.8, washout of isoflurane at pHi 6.8, and 0 ATP at pHi 6.8. Decreasing pHi to 6.8 increased channel activity, and during these conditions, isoflurane further enhanced channel activity. Open probability (Po) values determined at each step of the experimental protocol are shown below traces. Dashed lines denote the closed state (C). Upward deflection indicates channel opening. (Lower) All-points amplitude histograms from recordings above. The amplitude of unitary outward current was not changed when decreasing pHi or during application of isoflurane at pHi 6.8.

nel activity were identified in these domains.^{18,19} Intracellular protons appear to increase the activity of K_{ATP} channel by specifically binding to histidine (His-175) on the C-terminus of the Kir6.2 subunit,^{18,19} and this site is




Fig. 5. Summary data for intracellular ATP ([ATP]i)-normalized open probability (Po) relationship obtained at intracellular pH (pHi) 7.4, pHi 6.8, and during isoflurane application at pHi 6.8. Each data point is a mean ± SEM from six patches. Solid lines are Hill fits to the normalized Po data. Decreasing pHi from 7.4 to 6.8 shifted the curve to the right, suggesting a decrease in ATP sensitivity. At pHi 6.8, isoflurane caused further rightward shift in [ATP]i-Po relationship, suggesting further decrease in ATP sensitivity. IC₅₀ and Hill coefficient values are reported in Results.

independent of the ATP-binding site, lysine (K185).^{17,21} Although independent, these sites appear to interact with each other, and allosteric modulation of the cloned K_{ATP} channels by ATP and H⁺ has been recently demonstrated.²¹ Whether the allosteric modulation of channel activity by intracellular protons and ATP may play a role in isoflurane potentiation is not known. Because during ionic conditions of our study at pHi 6.8 isoflurane increased channel activity by reducing ATP sensitivity but did not affect unitary current amplitude or conductance, the channel pore is probably not targeted by the anesthetic. However, there is still a possibility of anesthetic interaction with the C terminus of the Kir6.2 subunit harboring the proton and ATP-binding sites. Furthermore, we cannot exclude a possibility of anesthetic interaction with the SUR2A subunit, modulating channel gating.^{24,25} Previous findings from our laboratory and results of this study suggest that isoflurane may enhance opening of the K_{ATP} channel previously activated or modified by the action of intracellular channel regulators, and that pHi may be one of them.

The functional significance of the mitochondrial *ver-sus* sarcolemmal K_{ATP} channels for cardioprotection remains controversial. Recent evidence suggests a role for mitochondrial K_{ATP} channels in the initiation of cardioprotection,²⁶ but sarcolemmal K_{ATP} channels have also been indicated in the protection afforded by ischemic

preconditioning.²⁷ Activation of the K_{ATP} channels is thought to be crucial for anesthetic preconditioning. However, the precise mechanism by which the enhancement of K_{ATP} channel activity by volatile anesthetics protects the myocardium is not yet established. Cardiac K_{ATP} channels are closed at physiologic [ATP]_i. During pathophysiologic conditions, such as ischemia, K_{ATP} channels activate during the first few minutes of the ischemic insult,²⁸ long before any significant decrease in [ATP]_i.²⁹ This suggested that other intracellular factors must be involved, and modulation of ATP sensitivity by factors targeting specifically the SUR subunit or the Kir6.x subunit has been demonstrated for ADP,^{24,25} PIP₂,^{30,31} and pHi.^{17,21} A transient decrease in pHi that occurs during ischemia³²⁻³⁴ and accompanies other metabolic stresses may promote opening of sarcolemmal K_{ATP} channels by decreasing sensitivity to ATP, thus allowing isoflurane interaction with the channel, leading to further enhancement of channel activity. Recent studies implicated an important role of the adenylate kinase and creatine kinase-mediated phosphotransfer in communicating mitochondria-generated signals to the sarcolemmal K_{ATP} channels.^{35,36} Whether volatile anesthetics alter the signal transfer to the cell subsarcolemmal compartment and K_{ATP} channels is an open question.

Our results support a possible role of volatile anesthetics in ischemia because of early ischemic acidosis.³² However, as demonstrated in the animal models and in humans, volatile anesthetics precondition the myocardium independently of ischemia.^{3,6,37-39} In clinical settings, the pHi dependence of K_{ATP} channel potentiation by isoflurane would likely play a protective role perioperatively when various metabolic stresses may produce a transient decrease in pHi.

In conclusion, this study provides evidence for pHi-dependent modulation of direct isoflurane interaction with the cardiac sarcolemmal K_{ATP} channel in guinea pig ventricular myocytes. Although at near physiologic pHi isoflurane has no effect nor inhibits K_{ATP} channel, at reduced pHi of 6.8, isoflurane increases channel Po and decreases sensitivity to inhibition by ATP as reflected by the rightward shift of the [ATP]_i-Po relationship.

References

1. Kersten JR, Schmeling TJ, Hettrick DR, Pagel PS, Gross GA, Warltier DC: Mechanism of myocardial protection by isoflurane. Role of adenosine triphosphate-regulated potassium (K_{ATP}) channels. *ANESTHESIOLOGY* 1996; 85:794-807
2. Kersten JR, Lowe D, Hettrick DA, Pagel PS, Gross GJ, Warltier DC: Glyburide, a KATP channel antagonist, attenuates the cardioprotective effects of isoflurane in stunned myocardium. *Anesth Analg* 1996; 83:27-33
3. Kersten JR, Schmeling TJ, Pagel PS, Gross GJ, Warltier DC: Isoflurane mimics ischemic preconditioning *via* activation of K_{ATP} channels: reduction of myocardial infarct size with an acute memory phase. *ANESTHESIOLOGY* 1997; 87:361-70
4. Nakayama M, Fujita S, Kanaya N, Tsuchida H, Namiki A: Blockade of ATP-sensitive K⁺ channels abolishes the anti-ischemic effects of isoflurane in dog hearts. *Acta Anaesthesiol Scand* 1997; 41:531-5
5. Toller WG, Gross ER, Kersten JR, Pagel PS, Gross GJ, Warltier DC: Sarcolemmal and mitochondrial adenosine triphosphate-dependent potassium channels: Mechanism of desflurane-induced cardioprotection. *ANESTHESIOLOGY* 2000; 92:1731-9
6. Rosco AK, Christensen JD, Lynch III C: Isoflurane but not halothane, induces protection of human myocardium *via* adenosine A₁ receptors and adenosine triphosphate-sensitive potassium channels. *ANESTHESIOLOGY* 2000; 92:1692-701
7. Han J, Kim E, Ho WK, Earm YE: Effects of volatile anesthetic isoflurane on ATP-sensitive K⁺ channels in rabbit ventricular myocytes. *Biochem Biophys Res Commun* 1996; 229:852-6
8. Fujimoto K, Bosnjak ZJ, Kwok WM: Isoflurane-induced facilitation of the cardiac sarcolemmal K_{ATP} channel. *ANESTHESIOLOGY* 2002; 97:57-65
9. Fan Z, Tokuyama Y, Makielski JC: Modulation of ATP-sensitive K⁺ channels by internal acidification in insulin-secreting cells. *Am J Physiol* 1994; 267:C1036-44
10. Davies NW: Modulation of ATP-sensitive K⁺ channels in skeletal muscle by intracellular protons. *Nature* 1990; 343:375-7
11. Davies NW, Standen NB, Stadnfield PR: The effects of intracellular pH on ATP-dependent potassium channels of frog skeletal muscle. *J Physiol (Lond)* 1992; 445:549-68
12. Koyano T, Kakei M, Nakashima H, Yoshinaga M, Matsuoka T, Tanaka H: ATP-regulated K⁺ channels are modulated by intracellular H⁺ in guinea-pig ventricular cells. *J Physiol (Lond)* 1993; 463:747-66
13. Cuevas J, Bassett AL, Cameron JS, Furukawa T, Myerburg RJ, Kimura S: Effect of H⁺ on ATP-regulated K⁺ channels in feline ventricular myocytes. *Am J Physiol* 1991; 261:H755-61
14. Fan Z, Makielski JC: Intracellular H⁺ and Ca²⁺ modulation of trypsin-modified ATP-sensitive K⁺ channels in rabbit ventricular myocytes. *Circ Res* 1993; 72:715-22
15. Fan Z, Furukawa T, Sawanobori T, Makielski JC, Hiraoka M: Cytoplasmic acidosis induces multiple conductance states in ATP-sensitive potassium channels of cardiac myocytes. *J Membrane Biol* 1993; 136:169-79
16. Lederer WJ, Nichols CG: Nucleotide modulation of the activity of rat heart ATP-sensitive K⁺ channels in isolated membrane patches. *J Physiol (Lond)* 1989; 419:193-211
17. Xu H, Cui N, Yang Z, Wu J, Giwa LR, Abdulkadir L, Sharma P, Jiang C: Direct activation of cloned K_{ATP} channels by intracellular acidosis. *J Biol Chem* 2001a; 276:12898-902
18. Piao H, Cui N, Xu H, Mao J, Rojas A, Wang R, Abdulkadir L, Li L, Wu J, Jiang C: Requirement of multiple protein domains and residues for gating K_{ATP} channels by intracellular pH. *J Biol Chem* 2001; 276:36673-80
19. Xu H, Wu J, Cui N, Abdulkadir L, Wang R, Mao J, Giwa LR, Chanchevalap S, Jiang C: Distinct histidine residues control the acid-induced activation and inhibition of the cloned K_{ATP} channel. *J Biol Chem* 2001; 276:38690-6
20. Alekseev AE, Gomez LA, Aleksandrova LA, Brady PA, Terzic A: Opening of cardiac sarcolemmal KATP channels by dinitrophenol separate from metabolic inhibition. *J Membrane Biol* 1997; 17:203-14
21. Wu J, Cui N, Piao H, Wang Y, Xu H, Mao J, Jiang C: Allosteric modulation of the mouse Kir6.2 channel by intracellular H⁺ and ATP. *J Physiol (Lond)* 2002; 543:495-504
22. Kwok WM, Martinelli AT, Fujimoto K, Suzuki A, Stadnicka A, Bosnjak ZJ: Differential modulation of the cardiac ATP-sensitive potassium channel by isoflurane and halothane. *ANESTHESIOLOGY* 2002; 97:50-6
23. Stadnicka A, Bosnjak ZJ, Kampine JP, Kwok WM: Modulation of cardiac inward rectifier K⁺ current by halothane and isoflurane. *Anesth Analg* 2000; 90:824-33
24. Zingman LV, Alekseev AE, Bienengraeber M, Hodgson D, Karger AB, Dzeja PP, Terzic A: Signaling in channel/enzyme multimers: ATPase transitions in SUR Module Gate ATP-sensitive K⁺ conductance. *Neuron* 2001; 31:233-45
25. Zingman LV, Hodgson DM, Bienengraeber M, Karger AB, Kathmann EC, Alekseev AE, Terzic A: Tandem function of nucleotide binding domains confers competence to sulfonylurea receptor in gating ATP-sensitive K⁺ channels. *J Biol Chem* 2002; 277:14206-10
26. Light PE: Cardiac K_{ATP} channels and ischemic preconditioning: current perspectives. *Can J Cardiol* 1999; 15:1123-30
27. Suzuki M, Sasaki N, Miki T, Sakamoto N, Ohmoto-Sekine Y, Tamagawa M, Seino S, Marban E, Nakaya H: Role of sarcolemmal K_{ATP} channels in cardioprotection against ischemia/reperfusion injury in mice. *J Clin Invest* 2002; 109:509-16
28. Nichols CG, Lederer WJ: Adenosine triphosphate-sensitive potassium channels in the cardiovascular system. *Am J Physiol* 1991; 261:H1675-86
29. Deutsch N, Weiss JN: ATP-sensitive K⁺ channel modification by metabolic inhibition in isolated guinea-pig ventricular myocytes. *J Physiol (Lond)* 1993; 465:163-79
30. Krauter T, Ruppertsberg JP, Baukowitz T: Phospholipids as modulators of K_{ATP} channels: Distinct mechanisms for control of sensitivity to sulphonylureas, K⁺ channel openers and ATP. *Mol Pharmacol* 2002; 59:1086-92
31. Fan Z, Makielski JC: Phosphoinositides decrease ATP sensitivity of the cardiac ATP-sensitive K⁺ channels. A molecular probe for the mechanism of ATP-sensitive inhibition. *J Gen Physiol* 1999; 114:251-69

32. Eisner DA, Nichols CG, O'Neill SC, Smith GL, Valdeolmillos M: The effects of metabolic inhibition on intracellular calcium and pH in isolated rat ventricular cells. *J Physiol (Lond)* 1989; 411:393-418
33. Orchard CH, Kentish JC: Effects of changes in pH on the contractile function of cardiac muscle. *Am J Physiol* 1990; C967-81
34. Steenbergen C, Perlman ME, London RE, Murphy E: Mechanism of preconditioning. Ionic alterations. *Circ Res* 1993; 72:112-25
35. Carrasco AJ, Dzeja PP, Alekseev AE, Pucar D, Zingman LV, Abraham MR, Hodgson D, Bienengraeber M, Puceat M, Janssen E, Wieringa B, Terzic A: Adenylate kinase phosphotransfer communicates cellular energetic signals to ATP-sensitive potassium channels. *Proc Natl Acad Sci U S A* 2001; 98:7623-8
36. Abraham MR, Selivanov VA, Hodgson DM, Pucar D, Zingman LV, Wieringa B, Dzeja PP, Alekseev AE, Terzic A: Coupling of cell energetics with membrane metabolic sensing. Integrative signaling through creatine kinase phosphotransfer disrupted by M-CK gene knock-out. *J Biol Chem* 2002; 277:24427-34
37. Cason BA, Gamperl AK, Slocum RE, Hickey RF: Anesthetic-induced preconditioning: previous administration of isoflurane decreases myocardial infarct size in rabbits. *ANESTHESIOLOGY* 1997; 87:1182-90
38. Belhomme D, Peynet J, Louzy M, Launay JM, Kitakaze M, Menasche P: Evidence for preconditioning by isoflurane in coronary artery bypass graft surgery. *Circulation* 1999; 100(suppl II):340-4
39. Hanouz JL, Yvon A, Massetti M, Lepage O, Babatasi G, Khayat A, Bricard H, Gerard JL: Mechanisms of desflurane-induced preconditioning in isolated human right atria in vitro. *ANESTHESIOLOGY* 2002; 97:33-41