

ERRATUM

The article by Meier *et al.* entitled "Hyperoxic Ventilation Reduces 6-h Mortality at the Critical Hemoglobin Concentration" was published in the January issue of the Journal (ANESTHESIOLOGY 2004; 100:70-6) with errors in table 1, table 2, and figure 6. The unit for heart rate in table 1 and table 2 should have read 1/min, and the units for base excess and arterial lactate concentration in table 1, table 2, and figure 6 should have read mmol/l. The authors regret these errors.

ERRATUM

The article by Xu *et al.* entitled "Halothane and Isoflurane Augment Depolarization-induced Cytosolic Ca^{2+} Transients and Attenuate Carbachol-stimulated Ca^{2+} Transients" was published in the June 2000 issue of the Journal (ANESTHESIOLOGY 2000; 92:1746-56) with an error in the labeling of figure 4. The corrected figure and legend appear below. The authors regret this error.




Fig. 4. (A-E) Representative traces of $[Ca^{2+}]_{cyt}$ -changes with sequential stimulation using 100 mM KCl followed by 1 mM carbachol (CAB). (A) Control. (B-F) Cells were pretreated for 5 min with 10 mM caffeine (CAF). (B), 1 μ M thapsigargin (TG) (C), 10 mM CAF and 1 μ M TG (D), 10 mM CAF and 1 μ M CAB (E), or with 1 μ M TG and 1 μ M CAB (F). (G, H) Statistical summary. Values presented are mean \pm SEM ($n = 2-19$). The significance of the differences among different groups was tested using the Student-Neumann-Keuls and labeled as follows: * $P < 0.05$ to the control group; \$ $P < 0.05$ to the TG group.