

# Effects of Volatile Anesthetics on Store-operated $\text{Ca}^{2+}$ Influx in Airway Smooth Muscle

Christina M. Pabelick, M.D.,\* Binnaz Ay, M.D.,† Yedatore S. Prakash, M.D., Ph.D.,‡ Gary C. Sieck, Ph.D.§

**Background:** In airway smooth muscle (ASM), volatile anesthetics deplete sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$  stores by increasing  $\text{Ca}^{2+}$  "leak." Accordingly, SR replenishment becomes dependent on  $\text{Ca}^{2+}$  influx. Depletion of SR  $\text{Ca}^{2+}$  stores triggers  $\text{Ca}^{2+}$  influx via specific plasma membrane channels, store-operated  $\text{Ca}^{2+}$  channels (SOCC). We hypothesized that anesthetics inhibit SOCC triggered by increased SR  $\text{Ca}^{2+}$  "leak," preventing SR replenishment and enhancing ASM relaxation.

**Methods:** In porcine ASM cells, SR  $\text{Ca}^{2+}$  was depleted by cyclopiazonic acid or caffeine in 0 extracellular  $\text{Ca}^{2+}$ , nifedipine and KCl (preventing  $\text{Ca}^{2+}$  influx through L-type and SOCC channels). Extracellular  $\text{Ca}^{2+}$  was rapidly introduced to selectively activate SOCC. After SOCC activation, SR was replenished and the protocol repeated in the presence of 1 or 2 minimum alveolar concentration halothane, isoflurane, or sevoflurane. In other cells, characteristics of SOCC and interactions between acetylcholine (ACh) and volatile anesthetics were examined.

**Results:** Cyclopiazonic acid produced slow SR leak, whereas the caffeine response was transient in ASM cells. Reintroduction of extracellular  $\text{Ca}^{2+}$  rapidly increased  $[\text{Ca}^{2+}]_i$ . This influx was insensitive to nifedipine, SKF-96365, and KBR-7943, inhibited by  $\text{Ni}^{2+}$  and blockade of inositol 1,4,5-triphosphate-induced SR  $\text{Ca}^{2+}$  release, and enhanced by ACh. Preexposure to 1 or 2 minimum alveolar concentration halothane completely inhibited  $\text{Ca}^{2+}$  influx when extracellular  $\text{Ca}^{2+}$  was reintroduced, whereas isoflurane and sevoflurane produced less inhibition. Only halothane and isoflurane inhibited ACh-induced augmentation of  $\text{Ca}^{2+}$  influx.

**Conclusion:** Volatile anesthetics inhibit a  $\text{Ni}^{2+}/\text{La}^{3+}$ -sensitive store-operated  $\text{Ca}^{2+}$  influx mechanism in porcine ASM cells, which likely helps maintain anesthetic-induced bronchodilation.

VOLATILE anesthetic-induced bronchodilation involves a reduction in intracellular  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ) that is key to development and maintenance of force in airway smooth muscle (ASM) cells.<sup>1-5</sup> These effects of anesthetics are partly attributable to an inhibition of  $\text{Ca}^{2+}$  influx via L-type  $\text{Ca}^{2+}$  channels.<sup>5-7</sup> We<sup>1,2</sup> and others<sup>8</sup> have shown that anesthetics also deplete  $\text{Ca}^{2+}$  stores of the sarcoplasmic reticulum (SR) by increasing  $\text{Ca}^{2+}$  "leakage." Such anesthetic-induced SR  $\text{Ca}^{2+}$  leak occurs via both inositol 1,4,5-triphosphate or ryanodine receptor (RyR) channels.<sup>2</sup>

$[\text{Ca}^{2+}]_i$  regulation in ASM is mediated by both extracellular  $\text{Ca}^{2+}$  influx and SR  $\text{Ca}^{2+}$  release. Both inositol 1,4,5-trisphosphate receptor<sup>9</sup> and RyR<sup>10</sup> channels play important roles in ASM  $\text{Ca}^{2+}$  regulation.  $\text{Ca}^{2+}$  influx occurs through both voltage-gated<sup>11</sup> and receptor-gated channels.<sup>12</sup> There is now considerable evidence from a variety of cell types that controlled  $\text{Ca}^{2+}$  influx through store-operated  $\text{Ca}^{2+}$  channels (SOCC; also termed capacitative  $\text{Ca}^{2+}$  entry) occurs in response to SR  $\text{Ca}^{2+}$  depletion, thus allowing for replenishment of intracellular  $\text{Ca}^{2+}$  stores.<sup>13-16</sup> Such influx does not occur through either L-type or receptor operated channels.<sup>13</sup>

There is now considerable evidence for SOCC in vascular<sup>13,17-20</sup> and other smooth muscle types.<sup>21</sup> Previous studies suggested that SOCC-mediated  $\text{Ca}^{2+}$  influx is restricted to the replenishment of inositol 1,4,5-trisphosphate-sensitive  $\text{Ca}^{2+}$  stores.<sup>13,22</sup> Recent studies in tissues other than ASM have suggested that  $\text{Ca}^{2+}$  release through RyR channels may also trigger SOCC.<sup>21,23</sup> The role of SOCC in ASM had been suggested by experiments in guinea pigs based on comparisons of contractions induced by agonists versus inhibitors of the SR  $\text{Ca}^{2+}$  adenosine triphosphatase.<sup>24</sup> Furthermore, SR  $\text{Ca}^{2+}$  adenosine triphosphatase inhibitors have been shown to increase  $[\text{Ca}^{2+}]_i$  in human bronchioles and bovine ASM, suggesting the involvement of SOCC.<sup>25</sup> However, characterization of SOCC-mediated  $\text{Ca}^{2+}$  influx in ASM is relatively lacking.

Volatile anesthetic-induced SR  $\text{Ca}^{2+}$  leak (and eventual depletion) may be expected to trigger SOCC-mediated  $\text{Ca}^{2+}$  influx in ASM cells. However, we hypothesize that volatile anesthetics will also inhibit such influx, thus accentuating the state of SR depletion.

## Materials and Methods

### Cell Preparation

Details of the ASM cell preparation technique have been previously described.<sup>10,26</sup> Porcine tracheae were obtained from a local abattoir. The smooth muscle layer was excised, minced in Hanks' balanced salt solution (HBSS) with 10 mM HEPES (pH 7.4; Invitrogen, Carlsbad, CA), and then incubated first for 2 h with 20 U/ml papain and 2000 U/ml DNase (Worthington Biochemical Corp., Lakewood, NJ) and subsequently for 1 h at 37°C with 1 mg/ml type intravenous collagenase (Worthington). The sample was triturated, centrifuged, and resuspended in minimum essential medium with 10% fetal calf serum. Isolated cells were plated on collagen-coated glass coverslips, and incubated for 1 h in 5%  $\text{CO}_2$  at 37°C.

\* Assistant Professor, † Research Fellow, ‡ Associate Professor, Department of Anesthesiology, § Professor, Department of Physiology and Biomedical Engineering, Mayo Clinic College of Medicine, and † Department of Anesthesiology, Marmara University, Istanbul, Turkey.

Received from the Department of Anesthesiology, Mayo Clinic College of Medicine, Rochester, Minnesota. Submitted for publication November 25, 2003. Accepted for publication March 29, 2004. Supported by grant GM56686 (GCS) from the National Institutes of Health, Bethesda, Maryland, the Mayo Foundation, Rochester, Minnesota (YSP, CMP) and Marmara University, Istanbul, Turkey (BA). Presented in part for a poster discussion at the American Society of Anesthesiologists Annual Meeting, San Francisco, California, October 11-15, 2003.

Address reprint requests to Dr. Prakash: Associate Professor of Anesthesiology: 4-184 W Jos SMH, Mayo Clinic, Rochester, Minnesota 55905. Address electronic mail to: prakash.js@mayo.edu. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

### Real-Time Confocal Imaging

Details of the confocal imaging technique for ASM cells have also been previously published.<sup>10,26</sup> Briefly, cells attached to coverslips were loaded with 5  $\mu\text{M}$  fluo-3 AM (Molecular Probes, Eugene, OR) for 30–45 min at 37°C in HBSS, washed, mounted on an open slide chamber (RC-25F, Warner Instruments, Hamden, CT), and perfused with HBSS at room temperature. An Odyssey XL real-time confocal system (Noran Instruments, Middleton, WI) with an Ar-Kr laser (488 nm line) and a Nikon 40 $\times$ /1.3 oil-immersion objective lens was used to visualize ASM cells. Using manufacturer-provided software, regions of interest encompassing single cells were outlined for measurements of  $[\text{Ca}^{2+}]_i$ . Empirical calibration of fluo-3 fluorescence concentrations for  $[\text{Ca}^{2+}]_i$  based on exposure to known  $\text{Ca}^{2+}$  concentrations was performed as described previously.<sup>26</sup>

### Administration of Volatile Anesthetics

As previously described,<sup>2</sup> calibrated online vaporizers were used to add halothane (Wyeth-Ayerst Laboratories, Philadelphia, PA), isoflurane, and sevoflurane (Abbott Laboratories, Deerfield, IL) to the aerating gas mixture. Aqueous anesthetic concentrations equivalent to 1 and 2 adult porcine minimum alveolar concentration (MAC) at room temperature (25°C) were determined for halothane and isoflurane by gas chromatography and electron capture detector (Hewlett-Packard 5880A; Hewlett-Packard, Sunnyvale, CA) and for sevoflurane by a flame ionization detector. Concentrations were halothane 1 MAC  $0.35 \pm 0.09$  mm, 2 MAC  $0.53 \pm 0.10$  mm, isoflurane 1 MAC  $0.4 \pm 0.09$  mm, 2 MAC  $0.6 \pm 0.11$  mm, and sevoflurane 1 MAC  $0.50 \pm 0.10$  mm, 2 MAC  $0.68 \pm 0.10$  mm.

### Store-Operated $\text{Ca}^{2+}$ Influx

Baseline  $\text{Ca}^{2+}$  concentrations were measured while ASM cells were perfused with HBSS. Extracellular  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_o$ ) was then removed by exposure to zero- $\text{Ca}^{2+}$  HBSS (5 mM EGTA). In the continued absence of  $[\text{Ca}^{2+}]_o$ , cells were also exposed to 1  $\mu\text{M}$  nifedipine and 10 mM KCl to ensure that L-type  $\text{Ca}^{2+}$  channels were not activated during the protocol. Cells were then rapidly exposed to 1  $\mu\text{M}$  cyclopiazonic acid (CPA) in zero- $\text{Ca}^{2+}$  HBSS. This technique has been frequently used to induce passive SR depletion by inhibiting SR  $\text{Ca}^{2+}$  adenosine triphosphatase with continued SR  $\text{Ca}^{2+}$  leak, presumably from both inositol 1,4,5-trisphosphate- and ryanodine-sensitive SR stores.<sup>13,16</sup> Continued SR  $\text{Ca}^{2+}$  leak resulted in increased  $[\text{Ca}^{2+}]_i$  concentrations that typically reached a plateau concentration or started to trend down (reflecting increasing  $\text{Ca}^{2+}$  efflux *via* the plasma membrane). At this point, 2.5 mM  $[\text{Ca}^{2+}]_o$  was rapidly reintroduced (in the continued presence of CPA). The observed  $[\text{Ca}^{2+}]_i$  response was then characterized using various pharmacological manipulations.

In separate sets of ASM cells, we performed additional


studies using caffeine for selective depletion of ryanodine-sensitive  $\text{Ca}^{2+}$  stores and the assessment of any interactions between  $\text{Ca}^{2+}$  release *via* RyR channels and SOCC. The protocol involved replacing CPA in the above experiments with 5 mM caffeine. As with the CPA protocols, pharmacological manipulations were used to characterize SOCC.

### Effect of Volatile Anesthetics on Store-Operated $\text{Ca}^{2+}$ Influx

SOCC in ASM cells was first established by performing a control protocol using CPA or caffeine in the presence of nifedipine and KCl (as above). Once SOCC-mediated  $\text{Ca}^{2+}$  influx was observed, the cells were washed for 15–20 min with HBSS to remove CPA (or caffeine) and to replenish SR  $\text{Ca}^{2+}$  stores.  $[\text{Ca}^{2+}]_o$  was then removed and the cells exposed to either CPA or caffeine. Once a  $[\text{Ca}^{2+}]_i$  response was observed (slow development of plateau with CPA or a transient with caffeine), the cells were exposed for 1 min to 1 or 2 MAC halothane, isoflurane, or sevoflurane in the continued presence of CPA or caffeine. This technique ensured that the anesthetics did not influence the SR  $\text{Ca}^{2+}$  release process itself, but were present in sufficient concentration before activation of SOCC. In the continued presence of CPA (or caffeine), and anesthetic,  $[\text{Ca}^{2+}]_o$  was rapidly reintroduced and the  $[\text{Ca}^{2+}]_i$  response was recorded. In control experiments, the SOCC protocol was performed twice without anesthetic, with an intervening wash in HBSS.

### Interactions Between Acetylcholine and Volatile Anesthetic Effects on Store-Operated $\text{Ca}^{2+}$ Influx

As above, SOCC in ASM cells was first established by performing a control protocol using CPA or caffeine in the presence of nifedipine and KCl. Cells were then washed in HBSS.  $[\text{Ca}^{2+}]_o$  was removed, nifedipine and KCl were added, and SR  $\text{Ca}^{2+}$  release was induced by CPA or caffeine. In the continued presence of CPA or caffeine, the cells were exposed to 10 nM, 100 nM, or 1  $\mu\text{M}$  acetylcholine (ACh) for 2 min. This ensured activation of muscarinic receptors occurred after SR  $\text{Ca}^{2+}$  release had already occurred. Activation of muscarinic receptors occurred with both protocols; however, in the case of CPA, SR  $\text{Ca}^{2+}$  depletion is passive, whereas for caffeine, complete release through RyR channels occurs, although inositol 1,4,5-trisphosphate-induced release is dependent on ACh concentration. In the continued presence of CPA (or caffeine) and ACh,  $[\text{Ca}^{2+}]_o$  was rapidly reintroduced to activated SOCC. Cells were then washed for a second time, and  $[\text{Ca}^{2+}]_o$  removed (with addition of nifedipine and KCl). After CPA or caffeine exposure, ACh was added as above and after 1 min, cells were exposed to 1 or 2 MAC halothane, isoflurane, or sevoflurane for an additional 1 min. This ensured that muscarinic activation occurred before anesthetic exposure and that confounding effects of anesthetic interactions with



**Fig. 1.**  $\text{Ca}^{2+}$  influx *via* store operated  $\text{Ca}^{2+}$  channels in porcine airway smooth muscle cells. The sarcoplasmic reticulum was depleted either by cyclopiazonic acid or caffeine in the absence of extracellular  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_o$ ). Subsequently,  $[\text{Ca}^{2+}]_o$  was reintroduced in the continued presence of cyclopiazonic acid or caffeine. HBSS = Hanks' balanced salt solution.

the receptor itself were avoided.  $[\text{Ca}^{2+}]_o$  was then rapidly reintroduced. In control protocols, after a control SOCC activation, the ACh protocol was performed twice, without anesthetic.

#### Statistical Analysis

For each ASM cell, comparisons before and after exposure to a drug were made using paired Student *t* test. Repeated measures analysis of variance was used for multiple comparisons with *post hoc* Bonferroni and Sheffé *F* tests. All experimental protocols were not performed in the same cells. Results were replicated in at least 3–5 cells obtained from each of five animals (paired comparisons within cells, independent testing across cells). Statistical significance was tested at  $P < 0.05$ . Values are reported as mean  $\pm$  SD.

## Results

#### Establishment of Store-Operated $\text{Ca}^{2+}$ Influx

At the beginning of experiments, baseline  $[\text{Ca}^{2+}]_i$  concentrations of ASM cells ranged from 70 to 100 nM ( $78 \pm 10$  nM;  $n = 128$ ). Neither removal of  $[\text{Ca}^{2+}]_o$  nor addition of nifedipine and KCl significantly affected  $[\text{Ca}^{2+}]_i$  concentrations. In the absence of  $[\text{Ca}^{2+}]_o$ , 1  $\mu\text{M}$  CPA resulted in increased  $[\text{Ca}^{2+}]_i$  concentrations that reached a plateau (228 to 518 nM;  $n = 65$ ). Subsequent rapid reintroduction of  $[\text{Ca}^{2+}]_o$  resulted in a further, sustained increase of  $[\text{Ca}^{2+}]_i$  (fig. 1A; 340 to 660 nM;  $P < 0.05$  when compared to the first plateau). In contrast to CPA, exposure to 5 mM caffeine in the absence of  $[\text{Ca}^{2+}]_o$ ,

**Table 1.** Characteristics of SOCC-Mediated  $\text{Ca}^{2+}$  Influx in ASM

Drug	$\text{Ca}^{2+}$ Influx (% peak CPA response)	$\text{Ca}^{2+}$ Influx (% peak caffeine response)
Control	$39.4 \pm 8.3$	$25.1 \pm 4.9$
1 $\mu\text{M}$ $\text{Ni}^{2+}$	$13.3 \pm 2.6^*$	$9.7 \pm 1.2^*$
1 $\mu\text{M}$ $\text{La}^{3+}$	$8.6 \pm 2.6^*$	$7.2 \pm 2.0^*$
10 $\mu\text{M}$ SKF96365	$9.9 \pm 5.7^*$	$6.3 \pm 3.4^*$
10 $\mu\text{M}$ KBR7943	$38.6 \pm 8.0$	$23.7 \pm 4.8$
10 nM ACh	$38.5 \pm 10.2$	$26.0 \pm 5.7$
100 nM ACh	$51.2 \pm 6.9^*$	$31.1 \pm 3.2^*$
1 $\mu\text{M}$ ACh	$65.2 \pm 5.2^*$	$38.6 \pm 3.5^*$

Values are means  $\pm$  SE.

\* Significant difference from control.

ACh = acetylcholine; CPA = cyclopiazonic acid; SOCC = store-operated  $\text{Ca}^{2+}$  channel.

resulted in a transient  $[\text{Ca}^{2+}]_i$  response (fig. 1B;  $n = 63$ ). Reintroduction of  $[\text{Ca}^{2+}]_o$  resulted in a relatively sustained increase in  $[\text{Ca}^{2+}]_i$  ranging from 110 to 386 nM (compared to a peak caffeine response ranging from 455 to 890 nM, the influx response was  $\sim 25\%$  of the caffeine peak). The influx observed with reintroduction of  $[\text{Ca}^{2+}]_o$  after caffeine exposure was significantly smaller than that after CPA exposure ( $P < 0.05$ ).

#### Characterization of Store-Operated $\text{Ca}^{2+}$ Influx

It has been previously reported that SOCC-mediated  $\text{Ca}^{2+}$  influx to be inhibited by  $\text{Ni}^{2+}$  and  $\text{La}^{3+}$ , at least in tissues other than ASM. After control activation of SOCC with CPA ( $n = 10$ ) or caffeine ( $n = 10$ ), and HBSS wash, ASM cells were exposed to nifedipine, KCl and 1  $\mu\text{M}$   $\text{NiCl}_2$  or  $\text{LaCl}_3$  in HBSS.  $[\text{Ca}^{2+}]_o$  was then removed and, in the continued presence of nifedipine, KCl and  $\text{Ni}^{2+}$  (or  $\text{La}^{3+}$ ), cells were reexposed to CPA or caffeine, with subsequent reintroduction of  $[\text{Ca}^{2+}]_o$ . In the presence of either 1  $\mu\text{M}$   $\text{Ni}^{2+}$  or  $\text{La}^{3+}$ , reintroduction of  $[\text{Ca}^{2+}]_o$  did not significantly increase  $[\text{Ca}^{2+}]_i$  (Table 1;  $P < 0.05$  compared to control) indicating that SOCC-mediated  $\text{Ca}^{2+}$  influx in ASM cells is sensitive to both ions.

To determine whether the observed  $\text{Ca}^{2+}$  influx was mediated by mechanisms other than SOCC (because L-type  $\text{Ca}^{2+}$  channels were already inhibited by nifedipine and clamped membrane potential), inhibitors of store-operated  $\text{Ca}^{2+}$  influx (SKF-96365; 10  $\mu\text{M}$ ) and  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchange (KBR-7943; 10  $\mu\text{M}$ ) were tested. Although the bidirectional plasma membrane  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchanger may not play a major role in most smooth muscle types, expression of this protein has been found in ASM.<sup>27</sup> When operating in the "reverse" or influx mode, the exchanger can bring  $\text{Ca}^{2+}$  intracellularly, in exchange for  $\text{Na}^+$ . After control SOCC activation with CPA ( $n = 11$ ) or caffeine ( $n = 11$ ) and HBSS wash, ASM cells were exposed to nifedipine, KCl, and 10  $\mu\text{M}$  SKF-96365 or KBR7943 in HBSS. The SOCC protocol was then repeated in the continued presence of these agents. In 90% of cells, SKF-96365 inhibited SOCC-mediated




Fig. 2. Effect of volatile anesthetics on store operated Ca<sup>2+</sup> channel (SOCC)-mediated Ca<sup>2+</sup> influx. After a control verification of SOCC-mediated Ca<sup>2+</sup> influx (A; caffeine protocol shown), airway smooth muscle (ASM) cells were washed (dotted lines) and reexposed to cyclopiazonic acid (CPA) or caffeine. Cells were then exposed to 1 or 2 minimum alveolar concentration (MAC) volatile anesthetic. Pre-exposure to 1 or 2 MAC halothane or isoflurane resulted in significant reduction of Ca<sup>2+</sup> influx on reintroduction of [Ca<sup>2+</sup>]<sub>o</sub> after SR Ca<sup>2+</sup> depletion by CPA (B) or caffeine (C). In contrast, only 2 MAC sevoflurane decreased influx in either protocol. Halothane and isoflurane were comparable in their inhibitory effect. \* = significant difference from control; # = significant difference between 1 and 2 MAC; % = significant difference from halothane ( $P < 0.05$ ); 0 MAC = control response.

Ca<sup>2+</sup> influx with both CPA and caffeine protocols (Table 1;  $P < 0.05$  compared with controls). In contrast, the presence of KBR-7943 had no effect on the [Ca<sup>2+</sup>]<sub>i</sub> response upon reintroduction of [Ca<sup>2+</sup>]<sub>o</sub> (Table 1), demonstrating that Na<sup>+</sup>/Ca<sup>2+</sup> exchange did not substantially contribute to the observed Ca<sup>2+</sup> influx.

#### Effect of ACh on Store-Operated Ca<sup>2+</sup> Influx

Compared with controls where no ACh was present, preexposure of ASM cells to either 100 nM or 1 μM ACh significant increased Ca<sup>2+</sup> influx with reintroduction of [Ca<sup>2+</sup>]<sub>o</sub> for both CPA and caffeine protocols ( $P < 0.05$ ; Table 1). In contrast, in other controls where the ACh protocol was repeated, there was ~5–7% decrease in the Ca<sup>2+</sup> influx upon repetition (not shown).

#### Effect of Volatile Anesthetics on Store-Operated Ca<sup>2+</sup> Influx

In ASM cells where SOCC-mediated Ca<sup>2+</sup> influx was first established, repetition of the protocol (CPA or caffeine) resulted in a 5–8% decrease in the observed Ca<sup>2+</sup> influx (rundown control). In contrast, exposure to halothane, isoflurane, or sevoflurane all resulted in significant decreases in the extent of Ca<sup>2+</sup> influx after reintroduction of [Ca<sup>2+</sup>]<sub>o</sub> ( $P < 0.05$  compared to control for each anesthetic at 1 or 2 MAC, both same cell and rundown control; fig. 2). For halothane and isoflurane, the effects at 1 MAC were significantly less than at 2 MAC, whereas for sevoflurane, the effects were comparable between 1 and 2 MAC. Among the three anesthetics, sevoflurane caused the least inhibition. At 2 MAC, halothane and isoflurane produced comparable inhibition (fig. 2). Be-




Fig. 3. Interaction between ACh and volatile anesthetic effects on store operated  $Ca^{2+}$  channel (SOCC)-mediated  $Ca^{2+}$  influx. Preexposure to  $1 \mu M$  ACh introduced after cyclopiazonic acid (CPA) or caffeine exposure was complete significantly enhanced SOCC-mediated  $Ca^{2+}$  influx. However, 1 or 2 minimum alveolar concentration (MAC) halothane or isoflurane (but not sevoflurane) significantly blunted ACh enhancement of  $Ca^{2+}$  influx after SR  $Ca^{2+}$  depletion by either CPA (panel A) or caffeine (panel B). Halothane and isoflurane were comparable in their inhibitory effect. \* = significant difference from control; # = significant difference between 1 and 2 MAC; % = significant difference from halothane ( $P < 0.05$ ).

tween the CPA and caffeine protocols, for each volatile anesthetic, the effect on  $Ca^{2+}$  influx was significantly greater for CPA compared to caffeine ( $P < 0.05$ ).

#### Interaction Between ACh and Volatile Anesthetic Effects on Store-Operated $Ca^{2+}$ Influx

In ASM cells in which both SOCC-mediated  $Ca^{2+}$  influx and an ACh-induced augmentation of influx were first established using the CPA or caffeine protocols, exposure to halothane and isoflurane, but not sevoflurane, resulted in significant decreases in the  $Ca^{2+}$  influx after reintroduction of  $[Ca^{2+}]_o$  ( $P < 0.05$  compared to both same cell and ACh rundown control for 1 or 2 MAC; fig. 3). Compared to anesthetic effects on  $Ca^{2+}$  influx in the absence of ACh, anesthetic inhibition of ACh-induced augmentation of influx was significantly greater ( $P < 0.05$ ). For halothane and isoflurane, the effects at 1 MAC were significantly less than at 2 MAC, with halothane causing greater inhibition. For both halothane and isoflu-

rane, the effect on ACh augmentation of  $Ca^{2+}$  influx was significantly greater for the CPA protocol ( $P < 0.05$ ).

## Discussion

Volatile anesthetic effects on ASM have been previously shown to involve inhibition of  $Ca^{2+}$  influx *via* L-type  $Ca^{2+}$  channels<sup>5-7</sup> as well as increased SR  $Ca^{2+}$  leak *via* both inositol 1,4,5-trisphosphate and RyR channels.<sup>1,2,8</sup> Such a leak may be expected to trigger SOCC-mediated  $Ca^{2+}$  influx in ASM cells, thus providing a mechanism for replenishment of depleted SR  $Ca^{2+}$  stores. However, in support of our hypothesis, the results of this study demonstrate that volatile anesthetics also inhibit such store-operated  $Ca^{2+}$  influx, thus accentuating the state of SR depletion. The results further demonstrate that volatile anesthetics, especially halothane and isoflurane, interfere with ACh-induced enhancement of SOCC-mediated  $Ca^{2+}$  influx, potentially maintaining bronchodilation even during agonist stimulation.

#### $[Ca^{2+}]_i$ Regulation in ASM

Regulation of  $[Ca^{2+}]_i$  in ASM is mediated by both extracellular  $Ca^{2+}$  influx and SR  $Ca^{2+}$  release *via* both inositol 1,4,5-trisphosphate receptor<sup>9</sup> and RyR<sup>10</sup> channels. Previous studies have demonstrated that  $Ca^{2+}$  influx in ASM occurs through both voltage-gated<sup>11</sup> and receptor-gated channels.<sup>12,28-30</sup> Agonists such as ACh are known to produce SR  $Ca^{2+}$  release *via* both inositol 1,4,5-trisphosphate and RyR channels.<sup>10</sup> Indeed, it is thought that the initial  $[Ca^{2+}]_i$  response to ACh stimulation is SR  $Ca^{2+}$  release, whereas maintenance of ASM  $[Ca^{2+}]_i$  and thus muscle tone is thought to involve sustained  $Ca^{2+}$  influx. However, we have previously shown that the  $[Ca^{2+}]_i$  response to ASM is oscillatory, mediated by repetitive SR  $Ca^{2+}$  release and reuptake, where initiation of oscillations is dependent on  $Ca^{2+}$  release through inositol 1,4,5-trisphosphate receptor channels, but sustenance of oscillations occurs through  $Ca^{2+}$ -induced  $Ca^{2+}$  release mechanisms *via* RyR channels.<sup>10,26</sup> The amplitude of  $[Ca^{2+}]_i$  oscillations represent SR  $Ca^{2+}$  content, and frequency depends on sensitivity for  $Ca^{2+}$  release *via* RyR channels. These studies also showed that  $Ca^{2+}$  influx serves to maintain and replenish SR  $Ca^{2+}$  because oscillations were not sustained in the absence of extracellular  $Ca^{2+}$  or in the presence of inhibitors of  $Ca^{2+}$  influx.<sup>10,26</sup> Our previous study on SOCC<sup>31</sup> as well as the current report emphasize the importance of this SR- and  $[Ca^{2+}]_o$ -dependent mechanism in maintaining the repletion state of the SR in ASM.

#### Effect of Volatile Anesthetics on $[Ca^{2+}]_i$ in ASM

Volatile anesthetics have been shown to target several  $[Ca^{2+}]_i$  regulatory mechanisms in ASM at both SR and

the plasma membrane. In terms of SR  $\text{Ca}^{2+}$  release, we and others have shown that volatile anesthetics deplete SR  $\text{Ca}^{2+}$  by decreasing SR  $\text{Ca}^{2+}$  content (store) *via* increased “leak” through both inositol 1,4,5-trisphosphate and RyR channels.<sup>2,32-35</sup> We have previously demonstrated that clinically relevant concentrations of halothane decrease both amplitude and frequency of ACh-induced  $[\text{Ca}^{2+}]_i$  oscillations. We interpreted these findings as a decrease in SR  $\text{Ca}^{2+}$  content and reduced sensitivity for  $\text{Ca}^{2+}$  induced  $\text{Ca}^{2+}$  release by volatile anesthetics.<sup>35</sup> These data collectively demonstrate that volatile anesthetics deplete SR  $\text{Ca}^{2+}$  stores, thus preventing an increase of  $[\text{Ca}^{2+}]_i$  in response to agonist stimulation. Anesthetic-induced depletion of SR  $\text{Ca}^{2+}$  then raises the question of whether  $\text{Ca}^{2+}$  influx *via* SOCC (if present) is then activated.

Given the existence of voltage-gated and receptor-operated  $\text{Ca}^{2+}$  influx mechanisms in ASM, there is already considerable evidence in the literature that the decrease in  $[\text{Ca}^{2+}]_i$  by volatile anesthetics involves inhibition of  $\text{Ca}^{2+}$  influx<sup>3,5</sup> through voltage-gated L-type  $\text{Ca}^{2+}$  channels.<sup>5,6</sup> These studies underline the inhibitory effect of volatile anesthetics on plasma membrane  $\text{Ca}^{2+}$  regulatory proteins. Accordingly, the results of the current study demonstrating inhibition of SOCC in the plasma membrane are consistent. Functionally, such inhibition only emphasizes the accentuation of anesthetic effects on  $[\text{Ca}^{2+}]_i$  in ASM because prevention of influx would only maintain the SR in a state of depletion, thus keeping  $[\text{Ca}^{2+}]_i$  low even with agonist stimulation.

#### *SOCC-Mediated $\text{Ca}^{2+}$ Influx and Volatile Anesthetics Effects*

In addition to previously established modes of  $\text{Ca}^{2+}$  influx, *i.e.*, voltage-gated<sup>11</sup> and receptor-gated channels,<sup>12,28-30</sup> in a recent study, we<sup>31</sup> found that  $\text{Ca}^{2+}$  influx in ASM occurs in response to SR  $\text{Ca}^{2+}$  depletion. Such a mechanism for influx in response to the state of SR  $\text{Ca}^{2+}$  stores was initially reported in parotid acinar cells and vascular smooth muscle in which influx was found to be dependent on  $[\text{Ca}^{2+}]_o$  but not L-type or receptor operated channels.<sup>13,36</sup> Other studies have demonstrated that SR  $\text{Ca}^{2+}$  adenosine triphosphatase inhibitors (such as thapsigargin, similar to CPA) increase  $\text{Ca}^{2+}$  influx or smooth muscle tone.<sup>20,37-40</sup> In these studies, whereas  $[\text{Ca}^{2+}]_o$  is necessary for maintenance of muscle tone or  $[\text{Ca}^{2+}]_i$ ,  $\text{Ca}^{2+}$ -channel blockers do not consistently prevent influx, suggesting that both voltage-gated  $\text{Ca}^{2+}$  channels and SOCC contribute to smooth muscle  $\text{Ca}^{2+}$  regulation and tone. Studies to date suggest that SOCC may play a significant role in tonic muscles such as blood vessels.<sup>13,17,18</sup> Only a few studies, including our own, have reported the existence of SOCC in ASM cells.<sup>31,41</sup>

Although the channels that represent SOCC have not been definitively identified, members of the tryptophan

family have been implicated in several studies, and mammalian tryptophan genes have been identified.<sup>15,42</sup> Some of these isoforms have been shown to be involved in SOCC.<sup>15,43</sup> We have previously verified the expression of several tryptophan C isoforms in porcine ASM,<sup>31</sup> although other studies have reported tryptophan C expression in guinea pig ASM.<sup>44</sup> Therefore, it is possible that SOCC in porcine ASM is mediated *via* tryptophan C proteins.

There are scarce data on anesthetic effects on SOCC (or capacitative  $\text{Ca}^{2+}$  influx per some authors). Horibe *et al.*<sup>45</sup> found that propofol inhibits capacitative  $\text{Ca}^{2+}$  entry in pulmonary artery smooth muscle. Tas *et al.*<sup>46</sup> found that in primary human endothelial cells, isoflurane inhibits histamine induced  $\text{Ca}^{2+}$  influx *via* SOCC. However, this group also found that enflurane enhances  $\text{Ca}^{2+}$  influx in rat glioma cells.<sup>47</sup> Pochet *et al.*<sup>48</sup> reported that trichloroethanol, the active metabolite of chloral hydrate, inhibits SOCC-mediated  $\text{Ca}^{2+}$  influx in submandibular acinar cells. The current study is the first to report the effects of anesthetics on SOCC in airway smooth muscle.

Studies from several cell types have now shown that SOCC does not appear to be activated by changes in membrane potential, thus making this mechanism different from L-type  $\text{Ca}^{2+}$  channels.<sup>13</sup> In accordance, in our previous study, we found that changes in membrane potential (induced by altering extracellular  $\text{K}^+$  concentration) did not affect SOCC-mediated  $\text{Ca}^{2+}$  influx in ASM.<sup>31</sup> Furthermore,  $\text{Ca}^{2+}$  influx after SR  $\text{Ca}^{2+}$  depletion by CPA was found to be insensitive to nifedipine, confirming that L-type  $\text{Ca}^{2+}$  channels do not mediate the observed influx. However, the existence of L-type  $\text{Ca}^{2+}$  channels in ASM has already been established,<sup>11,49</sup> as has the inhibitory effect of volatile anesthetics.<sup>5,6</sup> Certainly, volatile anesthetic inhibition of L-type channels would also prevent SR repletion, and blunt the  $[\text{Ca}^{2+}]_i$  response to ACh. Therefore, in the current study, we used nifedipine to block the confounding effect of this mechanism on the observed influx and volatile anesthetic effects. Addition of KCl to the extracellular medium also helped maintain membrane potential at a value where L-type  $\text{Ca}^{2+}$  channels were unlikely to be activated.

Consistent with our previous study, and with studies in other smooth muscle types, we found that SOCC is blocked by  $\text{Ni}^{2+}$ <sup>17,19,21,31,41</sup> and  $\text{La}^{3+}$ .<sup>17,21,31</sup> Although both ions are capable of nonspecifically inhibiting influx *via* several mechanisms, inhibition of influx by  $\mu\text{M}$  concentrations of either ion strongly indicate that the observed influx is mediated *via* SOCC.

Previous studies in tissues other than ASM have found that SOCC-mediated  $\text{Ca}^{2+}$  influx occurs after depletion of inositol 1,4,5-trisphosphate-sensitive SR  $\text{Ca}^{2+}$  stores.<sup>13,22</sup> In our previous study, we found that SOCC in ASM is triggered after SR  $\text{Ca}^{2+}$  release *via* inositol 1,4,5-trisphosphate receptor channels.<sup>31</sup> It has been previ-

ously demonstrated that volatile anesthetics markedly increase inositol 1,4,5-trisphosphate concentrations in ASM cells, even in the absence of agonist stimulation<sup>2</sup> (but see Yamakage *et al.*<sup>8</sup>). Accordingly, anesthetic exposure should enhance SOCC-mediated  $\text{Ca}^{2+}$  influx. However, we<sup>31</sup> and others<sup>14</sup> have found that inositol 1,4,5-trisphosphate by itself does not trigger SOCC-mediated  $\text{Ca}^{2+}$  influx even in the presence of agonist stimulation, and it is SR  $\text{Ca}^{2+}$  release *via* inositol 1,4,5-trisphosphate receptor channels that is important. In this regard, previous studies have demonstrated that volatile anesthetics enhance SR  $\text{Ca}^{2+}$  leak *via* inositol 1,4,5-trisphosphate receptor channels.<sup>2,32-35</sup> This should lead to enhanced SOCC-mediated  $\text{Ca}^{2+}$  influx in the presence of anesthetics. However, the current study demonstrates that despite this potential enhancement, volatile anesthetics can inhibit  $\text{Ca}^{2+}$  influx.

In addition to triggering of SOCC after release *via* inositol 1,4,5-trisphosphate receptor channels, there are now data in skeletal muscle<sup>50,51</sup> and in ASM<sup>31</sup> demonstrating SOCC activation in response to depletion of caffeine-sensitive SR  $\text{Ca}^{2+}$  stores and subsequent introduction of  $[\text{Ca}^{2+}]_o$ . Again, it has been previously demonstrated that volatile anesthetics increase SR  $\text{Ca}^{2+}$  leak *via* RyR channels,<sup>2,52,53</sup> suggesting that in the presence of anesthetics, SOCC-mediated influx may be enhanced. However, as with inositol 1,4,5-trisphosphate receptor channels, the data from the current study (*i.e.*, the caffeine protocol), demonstrates that volatile anesthetics overwhelm any potential enhancement of influx resulting from depletion of caffeine-sensitive  $\text{Ca}^{2+}$  stores.

In ASM, exposure to ACh appears to enhance SOCC-mediated influx. However, it is not entirely clear whether agonist stimulation modulates SOCC because influx is confounded by the presence of mechanisms for  $\text{Ca}^{2+}$  influx that are neither voltage-gated nor mediated by SOCC, *i.e.*, G-protein coupled receptor-operated mechanisms.<sup>41,54,55</sup> Whether such a mechanism is involved in ASM is also not clear.<sup>41</sup> In our previous study,<sup>31</sup> we found that selective activation of the non-SOCC component by preventing SR  $\text{Ca}^{2+}$  depletion resulted in significantly less enhancement of  $\text{Ca}^{2+}$  influx by ACh compared with when SR depletion was present, suggesting the existence of both SOCC and noncapacitative  $\text{Ca}^{2+}$  influx in ASM. Regardless, the data at least suggests that ACh does modulate SOCC in ASM.

Muscarinic modulation of SOCC has the potential for at least partial inhibition of anesthetic blockade of SOCC; however, it is possible that anesthetic effects on SOCC more likely help maintain bronchodilation, even during agonist stimulation. Furthermore, volatile anesthetics also inhibit muscarinic receptor activation.<sup>56,57</sup> Accordingly, even in the presence of ACh, volatile anesthetics likely maintain their inhibition of SOCC.

### Significance of SOCC and Volatile Anesthetic Effects

Increased  $[\text{Ca}^{2+}]_i$  is key to force generation in ASM. SOCC-mediated  $\text{Ca}^{2+}$  influx has been shown to produce ASM contraction.<sup>25</sup> Accordingly, anesthetic inhibition of SOCC has the potential to prevent SR  $\text{Ca}^{2+}$  repletion and agonist-induced bronchoconstriction. Volatile anesthetics are known to affect other mechanisms that regulate both  $[\text{Ca}^{2+}]_i$  (*e.g.*, SR and voltage-gated  $\text{Ca}^{2+}$  channels) and force (*e.g.*,  $\text{Ca}^{2+}$  sensitivity of force generation<sup>58</sup>), albeit with different relative potencies (*e.g.*, sevoflurane has a much smaller effect on influx, but is more potent than isoflurane in decreasing  $\text{Ca}^{2+}$  sensitivity). Nonetheless, anesthetic inhibition of SOCC may significantly contribute to the overall bronchodilatory effect of these agents by decreasing overall  $[\text{Ca}^{2+}]_i$  availability and thus help to maintain bronchodilation.

The authors thank Larry Hunter, M.S. (Associate in Research, Department of Anesthesiology, Mayo Clinic, Rochester, Minnesota) and Thomas Keller (Technician, Department of Anesthesiology, Mayo Clinic, Rochester, Minnesota) for their superb technical assistance.

### References

1. Pabelick CM, Prakash YS, Kannan MS, Jones KA, Warner DO, Sieck GC: Effect of halothane on intracellular calcium oscillations in porcine tracheal smooth muscle cells. *Am J Physiol* 1999; 276:L81-9
2. Pabelick CM, Prakash YS, Kannan MS, Sieck GC: Effect of halothane on sarcoplasmic reticulum  $\text{Ca}^{2+}$  release channels in porcine airway smooth muscle. *ANESTHESIOLOGY* 2001; 95:207-15
3. Jones KA, Housmans PR, Warner DO, Lorenz RR, Rehder K: Halothane alters cytosolic calcium transient in tracheal smooth muscle. *Am J Physiol* 1993; 265:L80-6
4. Yamakage M: Direct inhibitory mechanisms of halothane on canine tracheal smooth muscle contraction. *ANESTHESIOLOGY* 1992; 77:546-53
5. Yamakage M, Kohro S, Kawamata T, Namiki A: Inhibitory effects of four inhaled anesthetics on canine tracheal smooth muscle contraction and intracellular  $\text{Ca}^{2+}$  concentration. *Anesth Analg* 1993; 77:67-72
6. Yamakage M, Hirshman CA, Croxton TL: Volatile anesthetics inhibit voltage-dependent  $\text{Ca}^{2+}$  channels in porcine tracheal smooth muscle cells. *Am J Physiol* 1995; 268:L187-91
7. Warner DO, Jones KA, Lorenz RR, Pabelick CM: Muscarinic receptor stimulation modulates the effect of halothane on  $\text{Mn}^{2+}$  influx in airway smooth muscle. *Am J Physiol* 1997; 273:C868-73
8. Yamakage M, Kohro S, Matsuzaki T, Tsuchida H, Namiki A: Role of intracellular  $\text{Ca}^{2+}$  stores in the inhibitory effect of halothane on airway smooth muscle contraction. *ANESTHESIOLOGY* 1998; 89:165-73
9. Coburn RF, Baron CB: Coupling mechanisms in airway smooth muscle. *Am J Physiol* 1990; 258:L119-33
10. Kannan MS, Prakash YS, Brenner T, Mickelson JR, Sieck GC: Role of ryanodine receptor channels in  $\text{Ca}^{2+}$  oscillations of porcine tracheal smooth muscle. *Am J Physiol (Lung Cell Mol Physiol)* 1997; 272:L659-64
11. Worley JF, Kotlikoff MI: Dihydropyridine-sensitive single calcium channels in airway smooth muscle cells. *Am J Physiol* 1990; 259:L468-80
12. Murray RK, Kotlikoff MI: Receptor-activated calcium influx in human airway smooth muscle cells. *J Physiol* 1991; 435:123-44
13. Parekh AB, Penner R: Store depletion and calcium influx. *Physiol Rev* 1997; 77:902-30
14. Putney JW, McKay RR: Capacitative calcium entry channels. *BioEssays* 1999; 21:38-46
15. Putney JW: TRP, inositol 1,4,5-trisphosphate receptors, and capacitative calcium entry. *Proc Natl Acad Sci U S A* 1999; 96:14669-71
16. Putney JW, Broad LM, Braun FJ, Lievreumont JP, Bird GS: Mechanisms of capacitative calcium entry. *J Cell Sci* 2001; 114:2223-9
17. McDaniel SS, Platoshyn O, Wang J, Yu Y, Sweeney M, Krick S, Rubin IJ, Yuan JX: Capacitative  $\text{Ca}^{2+}$  entry in agonist-induced pulmonary vasoconstriction. *Am J Physiol Lung Cell Mol Physiol* 2001; 280:L870-80
18. Gardner JD, Benoit JN: Effects of capacitative calcium entry on agonist-induced calcium transients in A7r5 vascular smooth muscle cells. *J Biomed Sci* 2000; 7:304-10
19. Ng LC, Gurney AM: Store-operated channels mediate  $\text{Ca}^{2+}$  influx and contraction in rat pulmonary artery. *Circ Res* 2001; 89:923-9
20. Albert AP, Large WA: Activation of store-operated channels by noradren-

- aline via protein kinase C in rabbit portal vein myocytes. *J Physiol* 2002; 544: 113-25
21. Wayman CP, Gibson A, McFadzean I: Depletion of either ryanodine- or  $IP_3$ -sensitive calcium stores activates capacitative calcium entry in mouse aortic smooth muscle cells. *Pflugers Arch* 1998; 435:231-9
  22. Hoth M, Penner R: Calcium release-activated calcium current in rat mast cells. *J Physiol* 1993; 465:359-86
  23. Fellner SK, Arendshorst WJ: Ryanodine receptor and capacitative  $Ca^{2+}$  entry in fresh preglomerular vascular smooth muscle cells. *Kidney Int* 2000; 58:1686-94
  24. Liu X, Farley JM: Depletion and refilling of acetylcholine- and caffeine-sensitive  $Ca^{++}$  stores in tracheal myocytes. *J Pharmacol Exp Ther* 1996; 277: 789-95
  25. Sweeney M, McDaniel SS, Platoshyn O, Zhang S, Yu Y, Lapp BR, Zhao Y, Thistlethwaite PA, Yuan JX: Role of capacitative  $Ca^{2+}$  entry in bronchial contraction and remodeling. *J Appl Physiol* 2002; 92:1594-602
  26. Prakash YS, Kannan MS, Sieck GC: Regulation of intracellular calcium oscillations in porcine tracheal smooth muscle cells. *Am J Physiol Cell Physiol* 1997; 272:C966-75
  27. Pitt A, Knox AJ: Molecular characterization of the human airway smooth muscle  $Na^+/Ca^{2+}$  exchanger. *Am J Respir Cell Mol Biol* 1996; 15:726-30
  28. Ito Y, Takagi K, Tomita T: Relaxant actions of isoprenaline on guinea-pig isolated tracheal smooth muscle. *Br J Pharmacol* 1995; 116:2738-42
  29. Matyas S, Pucovsky V, Bauer V: Involvement of different  $Ca^{2+}$  sources in changes of responsiveness of guinea-pig trachea to repeated administration of histamine and acetylcholine. *Gen Physiol Biophys* 1995; 14:51-60
  30. al-Hassani MH, Garcia JG, Gunst SJ: Differences in  $Ca^{2+}$  mobilization by muscarinic agonists in tracheal smooth muscle. *Am J Physiol* 1993; 264:L53-9
  31. Ay B, Prakash YS, Pabelick CM, Sieck GC: Store-operated  $Ca^{2+}$  entry in porcine airway smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 2004; 286:L909-17
  32. Hossain MD, Evers AS: Volatile anesthetic-induced efflux of calcium from  $IP_3$ -gated stores in clonal (GH3) pituitary cells. *ANESTHESIOLOGY* 1994; 80:1379-89
  33. Fehr DM, Larach DR, Zangari KA, Schuler HG: Halothane constricts bovine pulmonary arteries by release of intracellular calcium. *J Pharmacol Exp Ther* 1996; 277:706-13
  34. Su JY, Zhang CC: Intracellular mechanisms of halothane's effect on isolated aortic strips of the rabbit. *ANESTHESIOLOGY* 1989; 71:409-17
  35. Pabelick CM, Prakash YS, Kannan MS, Jones KA, Warner DO, Sieck GC: Effect of halothane on intracellular calcium oscillations in porcine tracheal smooth muscle cells. *Am J Physiol* 1999; 276:L81-9
  36. Xuan YT, Wang OL, Wharton AR: Thapsigargin stimulates  $Ca^{2+}$  entry in vascular smooth muscle cells: Nicardipine-sensitive and -insensitive pathways. *Am J Physiol* 1992; 262:C1258-65
  37. Ohta T, Kawai K, Ito S, Nakazato Y:  $Ca^{2+}$  entry activated by emptying of intracellular  $Ca^{2+}$  stores in ileal smooth muscle of the rat. *Br J Pharmacol* 1995; 114:1165-70
  38. Gibson A, McFadzean I, Wallace P, Wayman CP: Capacitative  $Ca^{2+}$  entry and the regulation of smooth muscle tone. *Trends Pharmacol Sci* 1998; 19:266-9
  39. Flemming R, Cheong A, Dedman AM, Beech DJ: Discrete store-operated calcium influx into an intracellular compartment in rabbit arteriolar smooth muscle. *J Physiol* 2002; 543:455-64
  40. Dalrymple A, Slater DM, Beech D, Poston L, Tribe RM: Molecular identification and localization of Trp homologues, putative calcium channels, in pregnant human uterus. *Mol Hum Reprod* 2002; 8:946-51
  41. Ito S, Kume H, Yamaki K, Katoh H, Honjo H, Kodama I, Hideharu H: Regulation of capacitative and noncapacitative receptor-operated  $Ca^{2+}$  entry by rho-kinase in tracheal smooth muscle. *Am J Respir Cell Mol Biol* 2002; 26:491-8
  42. Minke B, Cook B: TRP channel proteins and signal transduction. *Physiol Rev* 2002; 82:429-72
  43. Xu SZ, Beech DJ: TrpC1 is a membrane-spanning subunit of store-operated  $Ca^{2+}$  channels in native vascular smooth muscle cells. *Circ Res* 2001; 88:84-7
  44. Ong HL, Brereton HM, Harland ML, Barritt GJ: Evidence for the expression of transient receptor potential proteins in guinea pig airway smooth muscle cells. *Respirology* 2003; 8:23-32
  45. Horibe M, Kondo I, Damron DS, Murray PA: Propofol attenuates capacitative calcium entry in pulmonary artery smooth muscle cells. *ANESTHESIOLOGY* 2001; 95:681-8
  46. Tas PW, Stobetael C, Roewer N: The volatile anesthetic isoflurane inhibits the histamine-induced  $Ca^{2+}$  influx in primary human endothelial cells. *Anesth Analg* 2003; 97:430-5
  47. Tas PW, Roewer N: The volatile anesthetic enflurane activates capacitative  $Ca^{2+}$  channels in rat glioma C6 cells. *Toxicol Lett* 1998; 100-101:265-9
  48. Pochet S, Keskiner N, Fernandez M, Marino A, Chaib N, Dehay JP, Metioui M: Multiple effects of trichloroethanol on calcium handling in rat submandibular acinar cells. *Br J Pharmacol* 2002; 136:568-80
  49. Shieh CC, Petrini MF, Dwyer TM, Farley JM: Calcium mobilization and muscle contraction induced by acetylcholine in swine trachealis. *J Biomed Sci* 1995; 2:272-82
  50. McKenzie S, Marley PD: Caffeine stimulates  $Ca^{2+}$  entry through store-operated channels to activate tyrosine hydroxylase in bovine chromaffin cells. *Eur J Neurosci* 2002; 15:1485-92
  51. Shin DW, Pan Z, Kim E, Lee J, Bhat M, Parness J, Kim do H, J: A retrograde signal from calsequestrin for the regulation of store-operated  $Ca^{2+}$  entry in skeletal muscle. *J Biol Chem* 2003; 278:3286-92
  52. Lynch C, Frazer MJ: Anesthetic alteration of ryanodine binding by cardiac calcium release channels. *Biochim Biophys Acta* 1994; 1194:109-17
  53. Blanck TJ, Peterson CV, Barody B, Tegazzin V, Lou J: Halothane, enflurane, and isoflurane stimulate calcium leakage from rabbit sarcoplasmic reticulum. *ANESTHESIOLOGY* 1992; 76:813-21
  54. Barritt GJ: Receptor-activated  $Ca^{2+}$  inflow in animal cells: a variety of pathways tailored to meet different intracellular  $Ca^{2+}$  signaling requirements. *Biochem J* 1999; 337:153-69
  55. McFadzean I, Gibson A: The developing relationship between receptor-operated and store-operated calcium channels in smooth muscle. *Br J Pharmacol* 2002; 135:1-13
  56. Kai T, Jones KA, Warner DO: Halothane attenuates calcium sensitization in airway smooth muscle by inhibiting G-proteins. *ANESTHESIOLOGY* 1998; 89:1543-52
  57. Yoshimura H, Jones KA, Perkins WJ, Warner DO: Dual effects of hexanol and halothane on the regulation of calcium sensitivity in airway smooth muscle. *ANESTHESIOLOGY* 2003; 98:871-80
  58. Kai T, Bremerich DH, Jones KA, Warner DO: Drug-specific effects of volatile anesthetics on  $Ca^{2+}$  sensitization in airway smooth muscle. *Anesth Analg* 1998; 87:425-9