

Additive Effects of Sevoflurane and Propofol on γ -Aminobutyric Acid Receptor Function

Luke E. Sebel, B.S.,* James E. Richardson, M.D.,† Sunil P. Singh, B.S.,‡ Shannon V. Bell, B.S.,§ Andrew Jenkins, Ph.D.#

Background: Previous studies have shown that propofol and sevoflurane enhance the function of γ -aminobutyric acid type A (GABA_A) receptors. However, it is not known whether these two drugs modulate the same molecular pathways. In addition, little is known about receptor function in the presence of both propofol and sevoflurane. The aim of this study was to better understand the interactions of propofol and sevoflurane with the GABA_A receptor.

Methods: Wild-type α_1 , β_2 , γ_2 s GABA_A receptor subunit complementary DNAs were transfected into human embryonic kidney cells grown on glass coverslips using a calcium phosphate transfection method. After transfection (36–72 h), cells were whole cell patch clamped and exposed to combinations of the following: 0.3–1,000 μ M γ -aminobutyric acid (GABA), 0–10 μ M propofol, and 0–1,650 μ M sevoflurane. Chemicals were delivered to the cells using two 10-channel infusion pumps and a rapid solution exchanger.

Results: Both propofol and sevoflurane alone enhanced the amplitude of GABA_A receptor responses to submaximal concentrations of GABA in a dose-dependent manner. The enhancement was underpinned by an increase in the apparent affinity of the receptor for GABA. Coapplication of both anesthetics further enhanced the apparent affinity of the receptor for GABA.

Conclusions: Response surface modeling of the potentiation of GABA responses (0.3–1,000 μ M) by sevoflurane and propofol revealed that the two anesthetics modulated receptor function in an additive manner. These results are consistent with recent mutagenesis studies, suggesting that these two drugs have separate binding sites and converging pathways of action on the GABA_A receptor.

MORE commonly than not, anesthesiologists use a combination of more than one drug to achieve a desired anesthetic state. With up to a dozen drugs often being used, the potential for drug interactions is huge. Therefore, understanding the interactions of general anesthetics is a critical part of understanding anesthesia practice.

The aim of this study and an accompanying clinical study is to better understand the interactions of propofol and sevoflurane.

Anesthetic interactions fall into three categories: The drugs can be additive, antagonistic (subadditive), or synergistic (supra-additive). Additive interactions occur when the effect from two drugs in combination equals the effect of either alone in an amount equal to the sum of the two drugs, after normalizing the concentrations to the intrinsic potency of each drug. Subadditive interactions occur when the effect of the combination is less than either alone, and synergism occurs when the effect is greater than either alone.¹ Additive interactions often occur when two drugs act *via* a similar mechanism, whereas synergism occurs when the two drugs act *via* different mechanisms.² Anesthesiologists have become experts at manipulating multidrug regimens. By making use of synergistic drug combinations to achieve a desired end point in an anesthetic-sparing manner, it is the hope of anesthesiologists to reduce patient recovery time and to increase patient safety.

Propofol and sevoflurane have both been shown to enhance the function of the γ -aminobutyric acid type A (GABA_A) receptor in neurons and in recombinant systems.^{3–6} GABA_A receptors mediate the fast inhibition of neuronal excitability by γ -aminobutyric acid (GABA), the most common inhibitory neurotransmitter in the central nervous system. GABA_A receptors are ligand-gated ion channels constructed from five subunits. The $\alpha_1\beta_2\gamma_2$ receptor subtype is the most prevalent in synapses of the adult mammalian central nervous system, accounting for approximately 40% of the total complement of GABA_A receptors.^{7,8} The function of neuronal GABA_A receptors is modulated by many general anesthetics⁹ at concentrations consistent with their clinical use.¹⁰

Recent site-directed mutagenesis studies have provided compelling evidence that many general anesthetics interact with two distinct binding sites on the GABA_A receptor. Mutation of a conserved serine (Ser270) in the second transmembrane domain of the α subunit is sufficient to block receptor enhancement by halogenated ether and alkane anesthetics such as isoflurane, sevoflurane, halothane, and chloroform.^{11–14} Conversely, mutation of a conserved asparagine (Asn265) in the second transmembrane domain or a conserved methionine (Met286) in the third transmembrane domain of the β subunit abolishes receptor modulation by propofol and etomidate.^{15,16} It is not yet known how drug interactions at these loci result in an enhancement of receptor func-

This article is accompanied by an Editorial View. Please see: Hemmings HC Jr: Do general anesthetics add up? ANESTHESIOLOGY 2006; 104:1120–2.

* Undergraduate Student, Department of Physics, Emory College, Atlanta, Georgia. † Resident Anesthesiologist, § Lead Research Associate, # Assistant Professor, Department of Anesthesiology, Emory University School of Medicine. ‡ Medical Student, New York University School of Medicine, New York, New York.

Received from the Department of Anesthesiology, Emory University School of Medicine, Atlanta, Georgia. Submitted for publication July 27, 2005. Accepted for publication November 30, 2005. Support was provided solely from institutional and/or departmental sources. Presented in part at the Annual Meeting of the American Society of Anesthesiologists, Las Vegas, Nevada, October 26, 2004.

Address correspondence to Dr. Jenkins: Department of Anesthesiology, Emory University School of Medicine, 1462 Clifton Road Northeast, Suite 420, Atlanta, Georgia 30322. andrew.jenkins@emoryhealthcare.org. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

tion. The aim of this study was to determine whether receptor modulation by combinations of both anesthetics occurred in an additive, synergistic, or antagonistic manner.

Materials and Methods

γ -Aminobutyric acid type A receptor complementary DNAs (cDNAs; gift from Neil Harrison, Ph.D., Professor of Pharmacology and Director, C.V. Starr Laboratory of Molecular Neuropharmacology, Department of Anesthesiology, Weill Cornell Medical College, New York, New York) were expressed *via* the vector pCIS2 in human embryonic kidney (HEK) 293 cells (American Type Culture Collection, Manassas, VA) as previously described.¹⁷ HEK-293 cells were cultured on poly-D-lysine-treated coverslips in a solution containing Eagle minimum essential medium supplemented with 5% fetal bovine serum (Hyclone, Logan, UT), L-glutamine (0.292 μ g/ml), penicillin G sodium (100 U/ml), and streptomycin sulfate (100 μ g/ml). For the transient expression of GABA_A receptors, cells were transfected using the CaPO₄ precipitation technique.^{18,19} The GABA_A receptor cDNAs and adeno-associated virus-green fluorescent protein cDNA (gift from H. Trent Spencer, Ph.D., Assistant Professor, Department of Pediatrics, Division of Hematology/Oncology and Bone Marrow Transplantation, Emory University School of Medicine, Atlanta, Georgia) were precipitated for 30 min at room temperature in a 160- μ l solution containing 125 mM CaCl₂, 140 mM NaCl, 750 μ M Na₂HPO₄, and 2.5 μ g of each cDNA. After 30 min, the mixture was added to cells grown on coverslips. The cDNA was in contact with the HEK cells for 24 h in an atmosphere containing 3% CO₂ (37°C) before being removed and replaced with fresh culture medium in an atmosphere of 5% CO₂ (37°C).

Coverslips of transfected cells were transferred 48–72 h after cDNA removal to a recording chamber and perfused continuously with extracellular solution (145 mM NaCl, 3 mM KCl, 1.5 mM CaCl₂, 1 mM MgCl₂, 6 mM D-glucose, 10 mM HEPES/NaOH adjusted to pH 7.4). Whole cell patch clamp recordings from fluorescing HEK-293 cells (voltage clamped at –60 mV) were made using a Multiclamp 700A amplifier (Axon Instruments, Foster City, CA) as described previously.²⁰ The resistance of the patch pipette was 4–6 M Ω when filled with intracellular solution (145 mM *N*-methyl-D-glucamine hydrochloride, 5 mM dipotassium ATP, 1.1 mM EGTA, 2 mM MgCl₂, 5 mM HEPES/KOH, 0.1 mM CaCl₂ adjusted to pH 7.2). In addition to the continuous bath perfusion with extracellular medium, solutions including GABA, general anesthetics, or both were applied rapidly to the cell using a motor-driven solution exchange device (Rapid Solution Changer RSC-160; Molecular Kinetics, Indianapolis, IN). Solutions were exchanged within approxi-

mately 50 ms. Laminar flow out of the rapid solution changer head was achieved by driving all solutions at identical flow rates (1.00 ml/min) *via* a multichannel infusion pump (KD Scientific, Holliston, MA). The solution changer was driven by protocols in the acquisition program pCLAMP 9 (Axon Instruments).

For GABA concentration-response measurements, cells were superfused with extracellular saline before application of one of eight GABA concentrations for 2 s, followed by a return to saline for at least 8 s before any subsequent GABA application. Below 100 μ M GABA, the responses did not desensitize; at and above 100 μ M, the amplitude of the responses declined by 10–15% in the continued presence of the agonist. Responses were low-pass-filtered (100 Hz; –3 dB, four-pole Bessel) and digitized with a 1322A interface (Axon Instruments) using pCLAMP 9 and stored for off-line analysis. Because intracellular and extracellular solutions contained equal chloride concentrations (145 mM), the chloride equilibrium potential was around 0 mV. All experiments were performed at room temperature (21°–24°C).

Stock solutions of GABA and propofol were diluted in extracellular solutions shortly before use. Sevoflurane solutions were prepared by injection of liquid anesthetic with a gas-tight syringe (Hamilton, Reno, NV) into intravenous drip bags containing defined volumes of extracellular solutions (100 ml) and used for up to 4 h. Clinically relevant concentrations of general anesthetics were used throughout the study; the aqueous concentration for 1 minimum alveolar concentration (MAC) sevoflurane was taken to be 330 μ M,⁵ and the anesthetic EC₅₀ (AC₅₀) concentration for propofol was taken to be 2 μ M.²¹ Losses of general anesthetics in this perfusion system have been measured using gas chromatography and typically represent only 5–10% of the initial total drug concentration.²² Sevoflurane was obtained from Abbott Laboratories (North Chicago, IL), and propofol (2,6 di-isopropylphenol) was obtained from Sigma (St. Louis, MO).

For each GABA exposure, the peak current amplitudes were measured and the GABA concentration-response data for each cell with and without general anesthetic were extracted from the raw data using our own software package. The analysis software was written to calculate nonlinear dose response curve parameters using Visual Basic macros within Microsoft Excel (Microsoft Corp., Redmond, WA) to facilitate efficient data organization. Dose-response parameters were optimized using GRG2, a version of the Generalized Reduction Gradient algorithm included in Microsoft Excel.²³ Using iterative processing and extensively automated file handling, we were able to process multiple data streams simultaneously.

The current peaks were fitted to a Hill equation of the form $I = I_{\max} * [GABA]^{nH} / ([GABA]^{nH} + EC_{50}^{nH})$, where I is the peak of each current, I_{\max} is the maximum

whole-cell current amplitude, [GABA] is the GABA concentration, EC_{50} is the GABA concentration eliciting a half-maximal current, and n_H is the Hill coefficient.

Concentration-response relations were recorded in the absence and presence of general anesthetic in the same cell. This enabled us to determine the control and anesthetic-modulated GABA EC_{50} s, defined as EG_{50} and EG_{50}' respectively. The fractional effect on EC_{50} was defined as $(1 - EG_{50}'/EG_{50})$.

Concentration-effect relations were calculated for propofol and sevoflurane. The fractional effects of anesthetic alone on GABA EC_{50} were fitted to a Hill equation of the form $E = [\text{anesthetic}]^{\text{slope}} / ([\text{anesthetic}]^{\text{slope}} + C_{50}^{\text{slope}})$, where E is the fractional effect of the anesthetic on GABA EC_{50} . For sinistral concentration-response shifts ($0 < E \leq 1$), [anesthetic] is the concentration of propofol or sevoflurane, C_{50} is the anesthetic concentration eliciting a half maximal effect, and slope is the Hill coefficient for the anesthetic concentration-effect relation. Statistical significance was assessed using a one-way analysis of variance with Dunnett test for multiple comparisons. Data are presented as mean \pm SEM.

The response surface for the modulation of GABA_A receptor function by propofol and sevoflurane was determined using the method of Minto *et al.*²⁴ Briefly, anesthetic concentrations were normalized to the C_{50} of each drug:

$$U_A = \frac{[A]}{C_{50,A}} \quad (1)$$

$$U_B = \frac{[B]}{C_{50,B}} \quad (2)$$

We used U_A and U_B to define a new variable θ , the drug ratio of A and B.

$$\theta = \frac{U_B}{U_A + U_B} \quad (3)$$

In the absence of sevoflurane (propofol alone), $U_A = 0$ and hence $\theta = 1$. Conversely, in the absence of propofol (sevoflurane alone), $U_B = 0$ and hence $\theta = 0$. When equal quantities of the drugs are present, $\theta = 0.5$.

Substituting these terms into the Hill equation, we obtain the following function:

$$E = \frac{\left(\frac{U_A + U_B}{U_{50}(\theta)}\right)^{\gamma(\theta)}}{1 + \left(\frac{U_A + U_B}{U_{50}(\theta)}\right)^{\gamma(\theta)}} \quad (4)$$

$U_{50}(\theta)$ defines the potency of the drug combination relative to the potency of either drug alone, and $\gamma(\theta)$ is the sigmoidicity of the response surface. When $U_{50}(\theta) > 1$, the interaction is synergistic. When $U_{50}(\theta) < 1$, the

interaction is subadditive or antagonistic. When $U_{50}(\theta) = 1$, the interaction is additive.

It has been demonstrated²⁴ that most isoboles have a simple inward or outward curvature and that they can be well approximated by a simple second order polynomial of the form

$$U_{50}(\theta) = 1 - \beta_{2U}\theta + \beta_{2U}\theta^2 \quad (5)$$

and

$$\gamma(\theta) = 1 - \beta_{2\gamma}\theta + \beta_{2\gamma}\theta^2 \quad (6)$$

These functions were fitted to the concentration-effect data using the Gauss-Newton nonlinear least-squares method (Statistics Toolbox function "nlinfit"; MATLAB, Natick, MA). The results were verified using the NONMEM program (Globomax, Hanover, MD).

Results

After transfection with adeno-associated virus-green fluorescent protein and GABA_A receptor α_1 , β_2 , and γ_{2S} and cDNAs, HEK-293, more than 90% of the cells were found to fluoresce, indicating that successful transfection conditions had occurred. Fluorescing cells were whole cell voltage clamped at -60 mV and superfused with extracellular saline. Application of GABA at eight different concentrations (0.3-1,000 μM) to the cells under these conditions elicited inward chloride currents in a concentration-dependent manner. Addition of 2 μM propofol to the extracellular medium resulted in an increase in the amplitudes of currents activated by 1-30 μM GABA, a small decrease in peak currents activated by 100-1,000 μM GABA, and an increase in the baseline noise of the recording. A representative whole cell voltage clamp recording showing the effect of 2 μM propofol on GABA_A receptor function is shown in figure 1. These effects were fully reversible. Propofol, 2 μM , reduced the GABA EC_{50} by a factor of 0.59 ± 0.05 , a fractional effect of 0.41 ± 0.05 (table 1). These experiments were repeated using 0.2-10 μM propofol. Propofol was found to reduce the GABA EC_{50} in a concentration-dependent manner. The C_{50} for the effect of propofol on the GABA EC_{50} was $C_{50,P} = 2.1 \pm 0.1 \mu\text{M}$, and the slope was 1.02 ± 0.05 .

Next, we repeated these experiments using sevoflurane. Addition of 330 μM sevoflurane to the extracellular saline also resulted in an increase in the amplitudes of currents activated by 1-30 μM GABA and an increase in the baseline noise of the recording (see representative trace in fig. 2). However, in contrast to the experiments with propofol, no reduction in the amplitudes of currents activated by 100-1,000 μM GABA was observed. These effects were also fully reversible. Sevoflurane, 330 μM , reduced the GABA EC_{50} by a factor of 0.46 ± 0.06 , a fractional effect of 0.54 ± 0.06 (table 2). These exper-




Fig. 1. Propofol, 2 μM , enhances currents activated by low ($< 30 \mu\text{M}$) concentrations of γ -aminobutyric acid (GABA). Whole cell responses to 0.3, 1, 3, 10, 30, 100, 300, and 1,000 μM GABA in the absence (first eight responses) and presence (last eight responses) of 2 μM propofol. Whole cell recordings were made from HEK-293 cells voltage clamped at -60 mV expressing $\alpha_1\beta_2\gamma_{2s}$ GABA_A receptor subunits. The filled bar above the current traces denotes the period of propofol application, and the open bars denote the period of GABA application. The application of 2 μM propofol shifted the EC_{50} for GABA from 4.8 to 2.6 μM , a fractional shift of 0.54.




Fig. 2. Sevoflurane, 330 μM , enhances currents activated by low ($< 100 \mu\text{M}$) concentrations of γ -aminobutyric acid (GABA). Whole cell responses to 0.3, 1, 3, 10, 30, 100, 300, and 1,000 μM GABA in the absence (first eight responses) and presence (last eight responses) of 330 μM sevoflurane. Whole cell recordings were made from HEK-293 cells voltage clamped at -60 mV expressing $\alpha_1\beta_2\gamma_{2s}$ GABA_A receptor subunits. The filled bar above the current traces denotes the period of sevoflurane application, and the open bars denote the period of GABA application. The application of 330 μM sevoflurane shifted the EC_{50} for GABA from 35.8 to 20.1 μM , a fractional shift of 0.56.

iments were repeated using 33–1,650 μM sevoflurane. Sevoflurane also reduced the GABA EC_{50} in a concentration-dependent manner. The C_{50} for the effect of sevoflurane on the GABA EC_{50} was $\text{C}_{50,S} = 306 \pm 15 \mu\text{M}$, and the slope was 1.00 ± 0.04 .

Finally, we measured the effect of combinations of propofol and sevoflurane on GABA receptor activation. The fractional effects of combinations of propofol and sevoflurane on GABA EC_{50} are shown in table 3. Figure 3 shows a representative recording of the combined effect of 1 μM propofol and 165 μM sevoflurane on currents activated by 0.3–1,000 μM GABA. As was observed with sevoflurane and propofol applied alone, the drug combination resulted in an increase in the amplitudes of currents activated by 1–30 μM GABA, a small decrease in peak currents activated by 100–1,000 μM GABA, and an increase in the baseline noise of the recording. These effects were fully reversible. A combination of 1 μM propofol with 165 μM sevoflurane resulted in a fractional effect of 0.52 ± 0.05 (table 3). The data shown in table 3 was used to construct a response

surface model for the modulation of GABA_A receptor function by propofol and sevoflurane (fig. 4).

Discussion

We found that when propofol and sevoflurane were applied alone and together at sufficient concentrations, they increased the amplitudes of submaximal GABA responses. Furthermore, they significantly increased the apparent affinity of GABA for human $\alpha_1\beta_2\gamma_{2s}$ GABA_A receptors heterologously expressed in HEK-293 cells, in 29 of the 33 combinations of anesthetics tested ($P < 0.05$).

Using the response surface modeling method,²⁴ we used these results to define the response surface for the actions of propofol and sevoflurane on the fractional effect on GABA EC_{50} . Using the methods described, we fitted the functions $U_{50}(\theta)$ and $\gamma(\theta)$ for $0 < \theta < 1$ (fig. 5). First, we noticed that the interaction surface for drug combinations retained the sigmoidicity of the two anesthetics applied alone. Second, we noticed that both β_{2U}

Table 1. Fractional Effect of 0–10 μM Propofol on GABA EC_{50}

	[Propofol], μM						
	0	0.2	0.4	1	2	4	10
Fractional effect on GABA EC_{50}	0 ± 0.01	-0.03 ± 0.05	0.06 ± 0.13	0.35 ± 0.02	0.41 ± 0.05	0.5 ± 0.06	0.84 ± 0.01

Values are mean \pm SEM and are determinations of at least five concentration–response shifts from at least three cells, as determined by the fractional change in the effective γ -aminobutyric acid concentration for 50% of maximal activation (GABA EC_{50}). Propofol concentrations are reported as molar concentrations. These concentrations equate to 0, 0.1, 0.2, 0.5, 1, 2, and 5 times the anesthetic EC_{50} of propofol.

Table 2. Fractional Effect of 0–1.65 mm Sevoflurane on GABA EC₅₀

Fractional effect on GABA EC ₅₀	[Sevoflurane], μm						
	0	33	66	165	330	660	1,650
	0 \pm 0.01	0.08 \pm 0.01	0.11 \pm 0.11	0.36 \pm 0.01	0.54 \pm 0.06	0.71 \pm 0.07	0.81 \pm 0.02

Values are mean \pm SEM and are determinations of at least five concentration–response shifts from at least three cells, as determined by the fractional change in the effective γ -aminobutyric acid concentration for 50% of maximal activation (GABA EC₅₀). Sevoflurane concentrations are reported as molar concentrations. These concentrations equate to 0, 0.1, 0.2, 0.5, 1, 2, and 5 times sevoflurane minimum alveolar concentration.

and $\beta_{2\gamma}$ were both close to zero. A NONMEM verification revealed that $\beta_{2U} = 0.25 \pm 0.49$ and $b_{2g} = 0.16 \pm 0.47$ and were not significantly different from zero. Because neither of the fitted functions significantly deviated from unity, we concluded that neither synergism nor antagonism was occurring and that combinations of propofol and sevoflurane resulted in a purely additive effect on receptor function.

The primary aim of this study and its companion article²⁵ was to determine whether propofol and sevoflurane had additive or synergistic effects in humans and on human receptors. In humans, synergy with anesthetic drugs is common. Opioids have been shown to reduce general anesthetic requirement (MAC) but are without any intrinsic anesthetic potency when administered alone. Therefore, the interactions between opioids and general anesthetics can be considered to be highly synergistic. A similar argument can be made for the synergistic actions of midazolam and thiopental.¹ Propofol and sevoflurane, however, are both general anesthetics in their own right, and we might therefore expect there to be less synergism between them. In the companion study to this report, Harris *et al.*²⁵ have shown that in humans, there is indeed no synergism between propofol and sevoflurane for immobility and loss of consciousness. Instead, the two anesthetics were shown to be additive. In this study, we have also shown that these two general anesthetics are additive in their actions on GABA_A receptors, the most common fast inhibitory neu-

rotransmitter receptor in both the brain and the spinal cord, sites thought to be critical for loss of consciousness and immobility, respectively.

The two drugs used in this study were selected for three important reasons. First, although both sevoflurane and propofol can be used alone for both induction and maintenance of anesthesia, these two anesthetics are commonly used together in the clinical setting. Second, these two anesthetics are chemically very different; sevoflurane is a small fluorinated ether, whereas propofol is a large phenol, and it is therefore unlikely that both compounds are capable of making the same array of bonds within a common anesthetic binding site. Finally, both propofol and sevoflurane exist as a single optical isomer. Conducting an additivity study with a pair of anesthetics, one of which exists as a racemic mixture of two or more stereoisomers, would have required an initial investigation into the synergistic, additive, or antagonistic activity of each of the isomers with respect to the other, before any consideration of the second drug could be undertaken. By using propofol and sevoflurane, we removed this requirement from our study.

We hypothesized that it would be unlikely for propofol and sevoflurane to compete for the same binding within the GABA_A receptor because of the large differences in their molecular structure. This hypothesis is strongly supported by the results of site directed mutagenesis experiments that are consistent with the hypothesis that inhaled general anesthetics mediate their effects within

Table 3. Fractional Effect of Propofol and Sevoflurane Combinations on GABA EC₅₀

Fractional Effect on GABA EC ₅₀	[Propofol], μm						
	0	0.2	0.4	1	2	4	10
[Sevoflurane], μm							
0	0.00 \pm 0.01	−0.03 \pm 0.05	0.06 \pm 0.13	0.35 \pm 0.02	0.41 \pm 0.05	0.50 \pm 0.06	0.84 \pm 0.01
33	0.08 \pm 0.01	0.08 \pm 0.01	0.22 \pm 0.05				
66	0.11 \pm 0.11	0.25 \pm 0.14	0.31 \pm 0.05				
165	0.36 \pm 0.01			0.52 \pm 0.05	0.52 \pm 0.08	0.67 \pm 0.02	0.79 \pm 0.02
330	0.54 \pm 0.06			0.60 \pm 0.01	0.58 \pm 0.03	0.79 \pm 0.06	0.90 \pm 0.06
660	0.71 \pm 0.07			0.61 \pm 0.04	0.69 \pm 0.02	0.79 \pm 0.04	0.90 \pm 0.05
1,650	0.81 \pm 0.02			0.84 \pm 0.03	0.90 \pm 0.04	0.89 \pm 0.01	0.91 \pm 0.04

Values are mean \pm SEM and are determinations of at least five concentration–response shifts from at least three cells, as determined by the fractional change in the effective γ -aminobutyric acid concentration for 50% of maximal activation (GABA EC₅₀). Propofol concentrations are reported as molar concentrations. These concentrations equate to 0, 0.1, 0.2, 0.5, 1, 2, and 5 times the anesthetic EC₅₀ of propofol. Sevoflurane concentrations are reported as molar concentrations. These concentrations equate to 0, 0.1, 0.2, 0.5, 1, 2, and 5 times sevoflurane minimum alveolar concentration.




Fig. 3. When applied together, 1 μM propofol and 165 μM sevoflurane enhance currents activated by low ($< 30 \mu\text{M}$) concentrations of γ -aminobutyric acid (GABA). Whole cell responses to 0.3, 1, 3, 10, 30, 100, 300, and 1,000 μM GABA in the absence (first eight responses) and presence (last eight responses) of 1 μM propofol and 165 μM sevoflurane. Whole cell recordings were made from HEK-293 cells voltage clamped at -60 mV expressing $\alpha_1\beta_2\gamma_{2s}$ GABA_A receptor subunits. The *filled bar* above the current traces denotes the period of propofol and sevoflurane application, and the *open bars* denote the period of GABA application. The application of 1 μM propofol and 165 μM sevoflurane together shifted the EC_{50} for GABA from 25.3 to 15.4 μM , a fractional shift of 0.61.

an anesthetic binding cavity located within the GABA_A α subunit,^{11–13,26} whereas propofol interacts directly with a genetically related cavity located within the β subunit.^{15,27,28}

Although there are many data for the separate site hypothesis for propofol and sevoflurane, the same cannot be said of volatile anesthetics that have been shown




Fig. 4. Concentration–response surface for the effect of 0–10 μM propofol and 0–1.65 mM sevoflurane on the fractional change in the effective γ -aminobutyric acid concentration for 50% of maximal activation (GABA EC_{50}) for $\alpha_1\beta_2\gamma_{2s}$ GABA_A receptors expressed in HEK-293 cells. The *horizontal axes* represent anesthetic concentrations normalized to their EC_{50} s for GABA EC_{50} modulation (U_A : sevoflurane; U_B : propofol). The *vertical axis* represents the fractional effect on GABA EC_{50} . The surface was fitted to the GABA EC_{50} modulation data in table 3 using a least squares algorithm written in Matlab.

to interact with the same cavity in GABA_A receptors.¹² However, it is unlikely that different anesthetics interact with this cavity in an identical manner. Rather than having simply antagonistic or additive actions, it is tempting to speculate whether a combination of chloroform and halothane, for example, could produce a better anesthetic combination for modulating GABA_A receptor function than an ether alone.

In addition to having different binding sites, these two drugs have been reported to modulate receptor function *via* overlapping but distinctly different mechanisms. Bai *et al.*³ concluded that propofol slowed the desensitization and deactivation of GABA_A receptors. This finding was supported by O’Shea *et al.*,²⁹ who proposed that propofol acted by modulating receptor gating, not ligand binding. Sevoflurane, however, has been shown to increase the apparent affinity of GABA for its receptor³⁰ and to block receptor function at high anesthetic concentrations, possibly *via* the same mechanism as observed with isoflurane.³¹

A recent study³² compared the effects propofol and another halogenated inhaled anesthetic, halothane, on GABA receptor single-channel currents and concluded that propofol acted by shortening slow closed times, whereas halothane prolonged the slow open times. It is important to note, however, that the effect of both drugs was to ultimately increase the open probability of the receptor’s ion channel.

Under equilibrium conditions, such as those presented here, the anesthetic-mediated increase in amplitudes of currents activated by submaximal GABA concentrations is thought to be underpinned by the stabilization of the open state of the receptor.²⁸ In synapses, where non-equilibrium conditions exist and the shape of the postsynaptic current is dictated by the intrinsic stochastic properties of the synaptic receptors, general anesthetics have been shown to prolong the decay of the postsynaptic current also by stabilizing the open state of the receptor.³

Currently, the overwhelming weight of experimental evidence suggests that propofol^{27,28} and sevoflurane^{5,6,14} alter the function of the nervous system by enhancing GABA_A receptor function. Notwithstanding this evidence, there are many novel proteins currently under investigation that may also play an important role in generating, or contributing to, the anesthetized state.^{33–38} In a recent study,³⁹ propofol and sevoflurane were both shown to depress the activity of ventral horn interneurons in cultured spinal cord slices. These two anesthetics yielded different effects on the patterns of action potential firing, which led the authors to suggest that in the spinal cord, propofol only acted *via* modulation of GABA_A receptors. The actions of sevoflurane were more complex and most likely result from interactions with glycine receptors, GABA_A receptors, and a third, unspecified component.




Fig. 5. Propofol and sevoflurane are additive in their ability to decrease the fractional change in the effective γ -aminobutyric acid concentration for 50% of maximal activation (GABA EC_{50}). $U_{50}(\theta)$ and $\gamma(\theta)$ were determined using the methods described and are plotted in the range $0 < \theta \leq 1$. Both $U_{50}(\theta)$ and $\gamma(\theta)$ did not significantly deviate from 1, indicating that there is no measurable synergism or antagonism between propofol and sevoflurane in the modulation of GABA receptor EC_{50} .

In this study, we examined the combined effect of clinically relevant concentrations of sevoflurane and propofol on the function of GABA_A receptors containing only the α_1 , β_2 , and γ_{2s} receptors. Although this receptor combination accounts for approximately 40% of the GABA_A receptors in the central nervous system, there are several other important tissue-specific combinations that have different kinetics and different general anesthetic sensitivities. It would be interesting if future studies investigated the effects of these two drugs on other subunit combinations (e.g., α_2 , β_3 , and γ_2) using the techniques described here and also electrophysiologic methods with higher temporal resolutions, e.g., single channel recording.

In conclusion, the data presented in this study and the results in the companion article²⁵ show that propofol and sevoflurane modulate GABA_A receptor function and generate the anesthetized state (immobility and loss of consciousness) in an additive manner. Neither of the two studies detected a significant degree of synergism or antagonism between propofol or sevoflurane in the three assays. Therefore, it seems that the most likely explanation for the results described here is that propofol and sevoflurane enhance GABA_A receptor function, in both the brain and the spinal cord. The two drugs do this by interacting at distinctly different binding sites within the same protein. However, the binding of anesthetic to either of the modulatory sites converges on a single effect: the enhancement of GABA_A receptor gating, resulting in an increase in the open probability of the integral ion channel. The resulting alteration in coincidence detection and synchrony in neuronal networks containing these receptors^{40,41} is likely to be fundamental in the generation of the anesthetized state.

The authors thank Steven Shafer, M.D. (Professor of Anesthesia, Stanford University, Stanford, California), for the NONMEM verification; Peter Sebel, M.B., B.S., Ph.D., M.B.A. (Professor of Anesthesiology, Emory University School of Medicine and Adjunct Professor of Psychology, Emory College of Emory University, Atlanta, Georgia), Jay Johansen, M.D., Ph.D. (Associate Professor, Department of Anesthesiology, Emory University School of Medicine, Grady Memorial Hospital, Atlanta, Georgia), Meagan Ward, A.B. (Neuroscience Graduate Student, Department of Physiology and Graduate School of Arts and Sciences, Emory University, Atlanta, Georgia), Adam Hall, Ph.D. (Assistant Professor, Department of Biologic Sciences, Smith College, Northampton, Massachusetts), and Nick Franks, Ph.D. (Professor of Biophysics and Anesthetics, Biophysics Section, The Blackett Laboratory, Imperial College London, United Kingdom), for helpful discussions.

References

1. Tverskoy M, Fleishman G, Bradley EL Jr, Kissin I: Midazolam-thiopental anesthetic interaction in patients. *Anesth Analg* 1988; 67:342-5
2. Evers A, Maze M: Common pharmacodynamic drug interactions in anesthetic practice. *Anesthetic Pharmacology: Physiologic Principles and Clinical Practice: A Companion to Miller's Anesthesia*. New York, Churchill Livingstone, 2004, pp 91-102
3. Bai D, Pennefather PS, MacDonald JF, Orser BA: The general anesthetic propofol slows deactivation and desensitization of GABA_A receptors. *J Neurosci* 1999; 19:10635-46
4. Adodra S, Hales TG: Potentiation, activation and blockade of GABA_A receptors of clonal murine hypothalamic GT1-7 neurones by propofol. *Br J Pharmacol* 1995; 115:953-60
5. Jenkins A, Franks NP, Lieb WR: Effects of temperature and volatile anesthetics on GABA_A receptors. *ANESTHESIOLOGY* 1999; 90:484-91
6. Hapfelmeier G, Schneck H, Kochs E: Sevoflurane potentiates and blocks GABA-induced currents through recombinant $\alpha(1)\beta(2)\gamma(2)$ GABA_A receptors: Implications for an enhanced GABAergic transmission. *Eur J Anaesthesiol* 2001; 18:377-83
7. McKernan RM, Whiting PJ: Which GABA_A-receptor subtypes really occur in the brain? *Trends Neurosci* 1996; 19:139-43
8. Fritschy JM, Mohler H: GABA_A-receptor heterogeneity in the adult rat brain: Differential regional and cellular distribution of seven major subunits. *J Comp Neurol* 1995; 359:154-94
9. Krasowski MD, Harrison NL: General anaesthetic actions on ligand-gated ion channels. *Cell Mol Life Sci* 1999; 55:1278-303
10. Franks NP, Lieb WR: Molecular and cellular mechanisms of general anaesthesia. *Nature* 1994; 367:607-14
11. Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, Harrison RA, Harrison NL: Sites of alcohol and volatile anaesthetic action on GABA_A and glycine receptors. *Nature* 1997; 389:385-9
12. Jenkins A, Greenblatt EP, Faulkner HJ, Bertaccini E, Light A, Lin A, Andreassen A, Viner A, Trudell JR, Harrison NL: Evidence for a common binding cavity for three general anesthetics within the GABA_A receptor. *J Neurosci* 2001; 21:RC136
13. Koltchine VV, Finn SE, Jenkins A, Nikolaeva N, Lin A, Harrison NL: Agonist gating and isoflurane potentiation in the human γ -aminobutyric acid type A receptor determined by the volume of a second transmembrane domain residue. *Mol Pharmacol* 1999; 56:1087-93
14. Nishikawa K, Harrison NL: The actions of sevoflurane and desflurane on the γ -aminobutyric acid receptor type A: Effects of TM2 mutations in the α and β subunits. *ANESTHESIOLOGY* 2003; 99:678-84
15. Krasowski MD, Koltchine VV, Rick CE, Ye Q, Finn SE, Harrison NL: Propofol and other intravenous anesthetics have sites of action on the γ -aminobutyric acid type A receptor distinct from that for isoflurane. *Mol Pharmacol* 1998; 53:530-8
16. Pistis M, Belelli D, McGurk K, Peters JA, Lambert JJ: Complementary regulation of anaesthetic activation of human $(\alpha 6\beta 3\gamma 2L)$ and *Drosophila* (RDL) GABA receptors by a single amino acid residue. *J Physiol* 1999; 515(pt 1):3-18
17. Jenkins A, Andreassen A, Trudell JR, Harrison NL: Tryptophan scanning mutagenesis in TM4 of the GABA_A receptor alpha 1 subunit: Implications for modulation by inhaled anesthetics and ion channel structure. *Neuropharmacology* 2002; 43:669-78
18. Chen C, Okayama H: High-efficiency transformation of mammalian cells by plasmid DNA. *Mol Cell Biol* 1987; 7:2745-52
19. Pritchett DB, Sontheimer H, Gorman CM, Kettenmann H, Seeburg PH, Schofield PR: Transient expression shows ligand gating and allosteric potentiation of GABA_A receptor subunits. *Science* 1988; 242:1306-8
20. Koltchine VV, Ye Q, Finn SE, Harrison NL: Chimeric GABA_A/glycine receptors: Expression and barbiturate pharmacology. *Neuropharmacology* 1996; 35:1445-56
21. Krasowski MD, Jenkins A, Flood P, Kung AY, Hopfinger AJ, Harrison NL: General anesthetic potencies of a series of propofol analogs correlate with

- potency for potentiation of γ -aminobutyric acid (GABA) current at the GABA_A receptor but not with lipid solubility. *J Pharmacol Exp Ther* 2001; 297:338-51
22. Krasowski MD, Harrison NL: The actions of ether, alcohol and alkane general anaesthetics on GABA_A and glycine receptors and the effects of TM2 and TM3 mutations. *Br J Pharmacol* 2000; 129:731-43
23. Lasdon LS, Fox RL, Ratner MW: Nonlinear optimization using the generalized reduced gradient method. *RAIRO Theoretical Informatics and Applications* 1974; 3:73-104
24. Minto CF, Schnider TW, Short TG, Gregg KM, Gentilini A, Shafer SL: Response surface model for anesthetic drug interactions. *ANESTHESIOLOGY* 2000; 92:1603-16
25. Harris RS, Lazar O, Johansen JW, Sebel PS: The interaction of propofol and sevoflurane on loss of consciousness and movement to skin incision during general anesthesia. *ANESTHESIOLOGY* 2006; 104:1170-5
26. Nishikawa K, Jenkins A, Paraskevakis I, Harrison NL: Volatile anesthetic actions on the GABA_A receptors: Contrasting effects of α 1(S270) and β 2(N265) point mutations. *Neuropharmacology* 2002; 42:337-45
27. Krasowski MD, Nishikawa K, Nikolaeva N, Lin A, Harrison NL: Methionine 286 in transmembrane domain 3 of the GABA_A receptor β subunit controls a binding cavity for propofol and other alkylphenol general anesthetics. *Neuropharmacology* 2001; 41:952-64
28. Jurd R, Arras M, Lambert S, Drexler B, Siegwart R, Crestani F, Zaugg M, Vogt KE, Ledermann B, Antkowiak B, Rudolph U: General anesthetic actions *in vivo* strongly attenuated by a point mutation in the GABA_A receptor β 3 subunit. *FASEB J* 2003; 17:250-2
29. O'Shea SM, Wong LC, Harrison NL: Propofol increases agonist efficacy at the GABA_A receptor. *Brain Res* 2000; 852:344-8
30. Haseneder R, Rammes G, Zieglgansberger W, Kochs E, Hapfelmeier G: GABA_A receptor activation and open-channel block by volatile anaesthetics: A new principle of receptor modulation? *Eur J Pharmacol* 2002; 451:43-50
31. Neumahr S, Hapfelmeier G, Scheller M, Schneck H, Franke C, Kochs E: Dual action of isoflurane on the γ -aminobutyric acid (GABA)-mediated currents through recombinant α (1) β (2) γ (2L)-GABA_A-receptor channels. *Anesth Analg* 2000; 90:1184-90
32. Kitamura A, Sato R, Marszalec W, Yeh JZ, Ogawa R, Narahashi T: Halothane and propofol modulation of γ -aminobutyric acid_A receptor single-channel currents. *Anesth Analg* 2004; 99:409-15
33. Franks NP, Dickinson R, de Sousa SLM, Hall AC, Lieb WR: How does xenon produce anaesthesia? (letter). *Nature* 1998; 396:324
34. Franks NP, Honore E: The TREK K2P channels and their role in general anaesthesia and neuroprotection. *Trends Pharmacol Sci* 2004; 25:601-8
35. Nelson LE, Lu J, Guo T, Saper CB, Franks NP, Maze M: The α ₂-adrenoceptor agonist dexmedetomidine converges on an endogenous sleep-promoting pathway to exert its sedative effects. *ANESTHESIOLOGY* 2003; 98:428-36
36. Violet JM, Downie DL, Nakisa RC, Lieb WR, Franks NP: Differential sensitivities of mammalian neuronal and muscle nicotinic acetylcholine receptors to general anesthetics. *ANESTHESIOLOGY* 1997; 86:866-874
37. Flood P, Sonner JM, Gong D, Coates KM: Isoflurane hyperalgesia is modulated by nicotinic inhibition. *ANESTHESIOLOGY* 2002; 97:192-8
38. Downie DL, Hall AC, Lieb WR, Franks NP: Effects of inhalational general anaesthetics on native glycine receptors in rat medullary neurones and recombinant glycine receptors in *Xenopus* oocytes. *Br J Pharmacol* 1996; 118:493-502
39. Grasshoff C, Antkowiak B: Propofol and sevoflurane depress spinal neurons *in vitro* via different molecular targets. *ANESTHESIOLOGY* 2004; 101:1167-76
40. Faulkner HJ, Traub RD, Whittington MA: Disruption of synchronous gamma oscillations in the rat hippocampal slice: A common mechanism of anaesthetic drug action. *Br J Pharmacol* 1998; 125:483-92
41. Dickinson R, Awaiz S, Whittington MA, Lieb WR, Franks NP: The effects of general anaesthetics on carbachol-evoked gamma oscillations in the rat hippocampus *in vitro*. *Neuropharmacology* 2003; 44:864-72