

When Is a Single-injection Nerve Block Not Really a Single Injection?

To the Editor:—I read, with interest, the case report describing a brachial plexopathy after an ultrasound-guided interscalene block in a patient with multiple sclerosis¹ and the accompanying editorial² and would like to make an observation not mentioned in either.

Interscalene blocks have been performed using either mechanical paresthesia or electrical nerve stimulation, for decades, with success rates reported to be 94–99%.^{3–5} In both of these techniques, the entire dose of local anesthetic is injected upon eliciting the initial desired response. These true single-injection techniques occur at the first nerve root, and likely the most superficial one, encountered. Perlas *et al.* used real-time ultrasound to quantify the sensitivity of both paresthesia and motor nerve stimulation techniques. A 22-gauge insulated needle was in the axilla of 103 patients, and after visualizing direct needle-nerve contact, the patients were asked whether they felt any paresthesia. The nerve stimulator was then turned on, and a motor response was sought at 0.5 mA or less. The authors concluded that there are a significant number of false-negative responses (direct needle-nerve contact *not* resulting in paresthesia or motor response) with these traditional methods of localization.⁶ This study showed that direct ultrasound visualization does not prevent intimate needle-nerve contact. Although Koff *et al.*¹ note that their needle “was not seen to penetrate the epineurium by [their] ultrasound image” after the first injection at C5, one must wonder how that initial volume of injection altered the ability to discern the needle-nerve relation of the three subsequent injections/maneuvers used to complete the block.

One of the many questions that needs to be addressed, as we continue to promote the benefits of ultrasound for peripheral nerve blocks, is whether there are any advantages to repositioning a needle

multiple times to be able to visualize local anesthetic spread around each of the nerve roots, because our historic success rates imply that this occurs adequately, with the initial injection. That is, does this practice of diving for individual and deeper nerve roots actually increase the risk to patients? The enemy of very good may prove to be better.

Meg A. Rosenblatt, M.D., The Mount Sinai School of Medicine, New York, New York. meg.rosenblatt@mountsinai.org

References

1. Koff MD, Cohen JA, McIntyre JJ, Carr CF, Sites BD: Severe brachial plexopathy after an ultrasound-guided single-injection nerve block for total shoulder arthroplasty in a patient with multiple sclerosis. *ANESTHESIOLOGY* 2008; 108:325–8
2. Hebl JR: Ultrasound-guided regional anesthesia and the prevention of neurologic injury. *ANESTHESIOLOGY* 2008; 108:186–8
3. Liguori GA, Zayas VM, YaDeau JT, Kahn RL, Paroli L, Buschiazio V, Wu A: Nerve localization techniques for interscalene brachial plexus blockade: A prospective, randomized comparison of mechanical paresthesia *versus* electrical stimulation. *Anesth Analg* 2006; 103:761–7
4. Bishop JY, Sprague M, Gelber J, Krol M, Rosenblatt MA, Gladstone JN, Flatow EL: Interscalene regional anesthesia for arthroscopic shoulder surgery: A safe and effective technique. *J Shoulder Elbow Surg* 2006; 15:567–70
5. Borgat A, Ekatothramis G, Kalbarer F, Benz C: Acute and non-acute complications associated with interscalene block and shoulder surgery: A prospective study. *ANESTHESIOLOGY* 2001; 95:875–80
6. Perlas A, Niaz A, McCartney C, Chan V, Xu D, Abbas S: The sensitivity of motor response to nerve stimulation and paresthesia for nerve localization as evaluated by ultrasound. *Reg Anesth Pain Med* 2006; 31:445–50

(Accepted for publication June 3, 2008.)

Ultrasound-guided Regional Anesthesia: Why Can't We All Just Stay Away from the Nerve?

To the Editor:—Hebl's response¹ to the case report by Koff *et al.*² highlights the quagmire that anesthesiologists have been stuck in since the invention of the nerve stimulator. This quagmire is “How close can I get to the nerve?” Because mechanical trauma and the risk of intraneural injection are two risk factors we have control over, shouldn't we be asking instead “How can I stay away from the nerve?”

Great regional anesthesiologists such as Winnie, Beck, and Dalens have published elegant block techniques using a detailed understanding of anatomy and fascial planes that do not require immediate proximity of the needle to the nerve.^{3–5} Today, we can use high-resolution ultrasound to visualize nerves, blood vessels, and fascial planes. Why not use this new technology to position the needle accurately in a fascial plane containing the nerve rather than as the “visual equivalent of a nerve stimulator”? Small wonder that those still trying to “position the tip of the needle next to the nerve and get a donut sign” have not demonstrated any outcome differences using ultrasound techniques.

At the University of Utah (Salt Lake City, Utah), we have adopted the philosophy correctly stated by Marhofer *et al.*⁶: “Nerves are not blocked by the needle but by the local anesthetic.” The results have been encouraging. All of our techniques for single and continuous nerve block placements are performed by injection into fascial planes containing the nerve and *not* by attempting to place the needle in close proximity to the nerve. We have performed more than 6,000 blocks using only ultrasound guidance, including more than 3,800 continuous catheters. A recent prospective study of 200 single and continuous

interscalene blocks performed here using only ultrasound guidance showed a success rate of 99%, with only 1% of patients having mild, transient sensory deficits.⁷ This is a considerable improvement over existing data for nerve stimulator techniques.^{8,9} Our published data for 620 outpatients with ultrasound-guided femoral, sciatic, and interscalene catheters also show high success and low complications in comparison with nerve stimulator techniques.¹⁰ As ultrasound gains more widespread application, additional outcome data will follow.

To say that ultrasound will not significantly improve patient safety is shortsighted. Many of the early techniques for ultrasound-guided blocks are still a variation on the nerve stimulator theme of “put the needle as close as possible to the nerve.” As we learn to use ultrasound to stay away from the nerve instead of getting close to it, we may be pleasantly surprised by the results. Dr. Hebl and others suggest that in ultrasound we have not found the “holy grail” of regional anesthesia.^{1,11} In our opinion, it could be the “holy grail”; we simply must know how to use it.

Jeffrey D. Swenson, M.D.,* Jennifer J. Davis, M.D. *University of Utah, Salt Lake City, Utah. jeff.swenson@hsc.utah.edu

References

1. Hebl JR: Ultrasound-guided regional anesthesia and the prevention of neurologic injury: Fact or fiction? *ANESTHESIOLOGY* 2008; 108:186–8

2. Koff MD, Cohen JA, McIntyre JJ, Carr CF, Sites BD: Severe brachial plexopathy after an ultrasound-guided single-injection nerve block for total shoulder arthroplasty in a patient with multiple sclerosis. *ANESTHESIOLOGY* 2008; 108:325-8
3. Winnie AP: Interscalene brachial plexus block. *Anesth Analg* 1970; 49:455-66
4. Beck GP: Anterior approach to sciatic nerve block. *ANESTHESIOLOGY* 1963; 24:222-4
5. Dalens B, Vanneville G, Tanguy A: Comparison of the fascia iliaca compartment block with the 3-in-1 block in children. *Anesth Analg* 1989; 69:705-13
6. Marhofer P, Greher M, Kapral S: Ultrasound guidance in regional anaesthesia. *Br J Anaesth* 2005; 94:7-17
7. Swenson JD, Davis JJ, Greis PE, Burks RT, Tashjian RZ: Interscalene block for postoperative analgesia using only ultrasound guidance: The outcome in 200 patients. *J Clin Anesth* 2008; (in press)

8. Brull R, McCartney CJ, Chan VW, El-Beheiry H: Neurological complications after regional anesthesia: Contemporary estimates of risk. *Anesth Analg* 2007; 104:965-74
9. Borgeat A, Ekatothramis G, Kalberer F, Benz C: Acute and nonacute complications associated with interscalene block and shoulder surgery: A prospective study. *ANESTHESIOLOGY* 2001; 95:875-80
10. Swenson JD, Bay N, Loose E, Bankhead B, Davis J, Beals TC, Bryan NA, Burks RT, Greis PE: Outpatient management of continuous peripheral nerve catheters placed using ultrasound guidance: An experience in 620 patients. *Anesth Analg* 2006; 103:1436-43
11. Horlocker TT, Wedel DJ: Ultrasound-guided regional anesthesia: In search of the holy grail. *Anesth Analg* 2007; 104:1009-11

(Accepted for publication June 3, 2008.)

Anesthesiology 2008; 109:749

Copyright © 2008, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Can Ultrasound Impact the Risk-Benefit Ratio for Nerve Blocks?

To the Editor:—Dr. Hebl's editorial¹ is thought provoking. The real potential of ultrasound lies in the opportunity to approach nerve blocks in entirely new ways. Regional anesthesia for hand surgery provides an example. Many anesthesiologists consider it axiomatic that one should avoid performing nerve blocks close to the tourniquet. This is partially defensive because "when the site of the block is close to the tourniquet, electromyographic and conduction nerve studies cannot allow such differentiation and by default, the anesthesiologist is usually blamed."² Despite this caveat, the most commonly used regional technique for hand surgery with an upper arm tourniquet is the axillary block. If one subscribes to the theory that serious nerve damage from axillary block is often the result of vascular trauma and a medial brachial fascial compartment syndrome,³ rather than direct needle-nerve trauma, it is even more imperative to avoid axillary block with an upper arm tourniquet.

Ultrasound should permit the near total abandonment of the axillary block. With ultrasound, peripheral nerve blocks in the mid forearm become straightforward.^{4,5} These blocks should replace the axillary block for minor hand surgery (e.g., carpal tunnel, digit fractures).⁶ Before ultrasound, many anesthesiologists avoided supraclavicular and infraclavicular blocks for fear of pneumothorax. With ultrasound, the lung and major vessels are directly visualized, so the proceduralist avoids them as a matter of course. For major hand, wrist, and elbow surgery, ultrasound-guided supraclavicular or infraclavicular blocks should be the first choice whenever an upper arm tourniquet is used.

The gist of Hebl's editorial and Koff *et al.*'s case report⁷ seems to be that this patient had an unfavorable risk-benefit ratio due to preexisting neurologic disease. But, where was the benefit in the first place? For total shoulder replacement, the requirement for significant pain control will far outlast any single-shot nerve block. The "tip-of-the iceberg" reports of permanent diaphragmatic paralysis⁸ after uneventful interscalene blocks offer another reason to demure the use of single-shot interscalene blocks for shoulder surgery. Ultrasound may allow more use of suprascapular blocks, either single shot or continuous.⁹ The ultrasound-guided suprascapular block may not be quite as effective, but it avoids the interscalene's dreaded complications. For major shoulder surgery without evidence of preexisting brachial plexopathy, continuous brachial plexus blocks are still encouraged. The risk of a continuous block can be justified to the patient because it provides prolonged analgesia matched to the requirements of major surgery.

Ultrasound allows the abandonment of some, but not all, traditional practices associated with nerve blocks. Many ultrasonographers no longer aspirate before injection, because ultrasound is a far more sensitive detector of intravascular injection than any aspiration test. The same cannot be said for potential intrafascicular injection. As Dr. Hebl notes, the precise position of the tip and the nerve is not always apparent until one has injected 2-3 ml of local anesthetic. In the unlikely situation where one has actually placed the needle tip into a nerve fascicle, the damage may already be done before anything shows up on ultrasound. Therefore, ultrasound may not obviate the need for injection pressure monitoring during nerve blocks. Injection pressure can be monitored with an inexpensive proprietary device (BSmart; Concert Medical, Norwell, MA) or by assessing the compression of an air bubble above the local anesthetic in the syringe.¹⁰

Only when we thoroughly rethink our approach to surgical nerve blocks will ultrasound fulfill its potential for improved safety.

Richard K. Baumgarten, M.D., Wayne State University, Detroit, Michigan. rkbaumgarten@comcast.net

References

1. Hebl JR: Ultrasound-guided regional anesthesia and the prevention of neurologic injury: Fact or fiction? *ANESTHESIOLOGY* 2008; 108:186-8
2. Chelly JE, Casati A: Perineural infusion of local anesthetics: "More to the review." *ANESTHESIOLOGY* 2007; 106:191
3. Tsao BE, Wilbourn AJ: Infraclavicular brachial plexus injury following axillary regional block. *Muscle Nerve* 2004; 30:44-8
4. Foxall GL, Skinner D, Hardman JG, Bedforth NM: Ultrasound anatomy of the radial nerve in the distal upper arm. *Reg Anesth Pain Med* 2007; 32:217-20
5. McCartney CJ, Xu D, Constantinescu C, Abbas S, Chan VW: Ultrasound examination of peripheral nerves in the forearm. *Reg Anesth Pain Med* 2007; 32:434-9
6. Gebhard RE, Al-Samsam T, Greger J, Khan A, Chelly JE: Distal nerve blocks at the wrist for outpatient carpal tunnel surgery offer intraoperative cardiovascular stability and reduce discharge time. *Anesth Analg* 2002; 95:351-5
7. Koff MD, Cohen JA, McIntyre JJ, Carr CF, Sites BD: Severe brachial plexopathy after an ultrasound-guided single-injection nerve block for total shoulder arthroplasty in a patient with multiple sclerosis. *ANESTHESIOLOGY* 2008; 108:325-8
8. Ediale KR, Myung CR, Neuman GG: Prolonged hemidiaphragmatic paralysis following interscalene brachial plexus block. *J Clin Anesth* 2004; 16:573-5
9. Meier G, Bauereis C, Maurer H: The modified technique of continuous suprascapular nerve block: A safe technique in the treatment of shoulder pain [in German]. *Anaesthesist* 2002; 51:747-53
10. Tsui BC, Li LX, Pillay JJ: Compressed air injection technique to standardize block injection pressures. *Can J Anaesth* 2006; 53:1098-102

(Accepted for publication June 3, 2008.)