

Modulation of Opioid Actions by Nitric Oxide Signaling

Noboru Toda, M.D., Ph.D.,* Shiroh Kishioka, M.D., Ph.D.,† Yoshio Hatano, M.D., Ph.D.,‡ Hiroshi Toda, M.D., Ph.D.§

Nitric oxide (NO) plays pivotal roles in controlling physiological functions, participates in pathophysiological intervention, and is involved in mechanisms underlying beneficial or untoward actions of therapeutic agents. Endogenous nitric oxide is formed by three isoforms of nitric oxide synthase: endothelial, neurogenic and inducible. The former two are constitutively present mainly in the endothelium and nervous system, respectively, and the latter one is induced by lipopolysaccharides or cytokines mainly in mitochondria and glial cells. Constitutively formed nitric oxide modulates the actions of morphine and related analgesics by either enhancing or reducing antinociception. Tolerance to and dependence on morphine or its withdrawal syndrome are likely prevented by nitric oxide synthase inhibition. Information concerning modulation of morphine actions by nitric oxide is undoubtedly useful in establishing new strategies for efficient antinociceptive treatment and for minimizing noxious and unintended reactions.

THE labile molecule NO has been widely recognized to play pivotal roles in the regulation of physiologic functions; in contrast, it also participates in pathophysiological intervention. Nitric oxide (NO) synthesis was first found in vascular endothelial cells,^{1–4} central and peripheral nerve cells and fibers,⁵ and macrophages.⁶ Investigations on the functional role of this inorganic molecule have been extended to other organs and tissues in the whole body. The NO/cyclic guanosine monophosphate (cyclic GMP) signaling pathway contributes to mechanisms underlying the action of therapeutic agents, the most promising through mechanisms of enhancing the action of endogenous NO is phosphodiesterase-5 inhibitors such as sildenafil⁷ and its congeners.⁸ Morphine and other opioid agonists exert analgesia through μ -opioid receptors at spinal and multiple supraspinal sites. δ -Opioid receptor agonists also are potent analgesics in animals and humans. In animals, agonists for

κ -receptors produce analgesia that is mediated at spinal sites. Some literature reports on opioid analgesics research have revealed that nitric oxide is involved in therapeutic actions but can also have untoward effects. Information concerning nitric oxide that undoubtedly plays important roles in modulating analgesic effects of opioids and their side effects would provide us with clues for establishing strategies for appropriate opioid therapy to enhance analgesic actions while minimizing tolerance, dependence, withdrawal syndrome, and other side effects.

Our previous review article⁹ summarized the involvement of nitric oxide in the actions of a variety of anesthetic agents without discussing the opioid analgesics, although these are useful for anesthesia and inevitably important for intolerable pain. The present article describes the involvement of endogenous nitric oxide in the antinociceptive effects of morphine and other related analgesics and the tolerance, dependence, and withdrawal syndrome associated with the use of these drugs.

Synthesis and Actions of NO

Nitric oxide is produced when L-arginine is transformed to L-citrulline through catalysis by nitric oxide synthase (NOS) in the presence of oxygen and cofactors. Ca^{2+} is required for the activation of neuronal NOS (nNOS, NOS I) and endothelial NOS (NOS III) but not inducible NOS (iNOS, NOS II). nNOS, mostly a soluble enzyme, is constitutively expressed in the brain,⁵ peripheral nerves, and kidneys. Endothelial NOS is also constitutively expressed mostly in particulate fractions of the endothelial cell.¹⁰ iNOS is not constitutively expressed but is induced mainly in macrophages in response to bacterial lipopolysaccharide and cytokines. The synthesis of nitric oxide by these NOS isoforms is inhibited by L-arginine analogs, including N^G -monomethyl-L-arginine (L-NMMA),¹¹ N^G -nitro-L-arginine (L-NA),^{12,13} L-NA methyl-ester (L-NAME),¹² and asymmetric dimethylarginine.¹⁴ 7-Nitroindazol (7-NI) is one of the most promising nNOS inhibitors introduced.¹⁵ Aminoguanidine has a long history as a selective iNOS inhibitor.¹⁶ Nitro compounds, including nitroglycerin, sodium nitroprusside, and S-nitroso-N-acetylpenicillamine, are capable of liberating nitric oxide and are called nitric oxide donors.

Endothelial nitric oxide causes vasodilatation, decreased vascular resistance, lowered blood pressure, inhibition of platelet aggregation and adhesion, inhibition

* Professor Emeritus, Shiga University of Medical Science, Shiga, Japan, and Toyama Institute for Cardiovascular Pharmacology Research, Osaka, Japan. † Professor of Pharmacology, Department of Pharmacology, ‡ Professor of Anesthesiology, Department of Anesthesiology, Wakayama Medical University, Wakayama, Japan. § Head of Anesthesiology, Department of Anesthesiology, Kyoto Katsura Hospital, Kyoto, Japan.

Received from the Toyama Institute for Cardiovascular Pharmacology Research, Osaka, Japan. Submitted for publication June 19, 2008. Accepted for publication September 8, 2008. Support was provided solely by institutional and/or departmental sources.

David S. Warner, M.D., served as Handling Editor for this article.

Address correspondence and reprint requests to Dr. Toda: Toyama Institute for Cardiovascular Pharmacology Research, 7-13, 1-Chome, Chuo-ku, Osaka 541-0052, Japan. n.toda.toyama-bldg@orion.ocn.ne.jp. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

of leukocyte adhesion and transmigration, and reduced vascular smooth muscle proliferation. Nitrovasodilators *via* release of nitric oxide activate soluble guanylyl cyclase and produce cyclic GMP from guanosine triphosphate in smooth muscle cells. Methylene blue, oxyhemoglobin, and 1H[1,2,4]oxadiazolo [4,3-a]quinoxalin-1-one¹⁷ inhibit guanylyl cyclase activity. Accumulation of cyclic GMP causes activation of cyclic GMP-dependent protein kinase that is involved in the adenosine triphosphate (ATP)-sensitive K⁺-channel opening to produce spinal or peripheral antinociception¹⁸⁻²⁰ and in Na⁺,K⁺-ATPase activation.²¹ Cyclic GMP-dependent protein kinase phosphorylates the serotonin transporter at Thr-276 and increases its activity by modifying the substrate permeation.^{22,23} Cyclic GMP is degraded by phosphodiesterase type 5 to 5'-GMP. Cyclic GMP is degraded by phosphodiesterase type-5 to 5'-GMP.

Nonadrenergic noncholinergic inhibitory responses to autonomic nerve stimulation are mainly mediated through nitric oxide synthesized by nNOS; Nitric oxide plays a crucial role as a neurotransmitter from the peripheral efferent nerves in cerebral blood vessels.²⁴ Afferent nitrenergic nerves control some aspects of sensory information processing. There is evidence that nitric oxide is the neurotransmitter released from primary sensory nerves that mediates mesenteric vasodilatation.²⁵ Nitric oxide offers important roles in afferent signaling of pain through the dorsal horn of the spinal cord and in autonomic control through nitrenergic innervation.²⁶ Many of the homeostatic actions of spinal afferents are brought about by release of transmitters (nitric oxide and calcitonin gene-related peptide) from their peripheral endings.²⁷

Nitric oxide functions mainly as a neuromodulator in the brain. Nitric oxide signaling appears to be essential for neural plasticity; that is, long-term potentiation in the hippocampus and long-term depression in the cerebellum. Glutamate participates mainly in synaptic interactions; with the help of nitric oxide, the strength of excitatory input might be nonsynaptically signaled to the surrounding monoaminergic neurons in the brain. Nitric oxide formed by *N*-methyl-D-aspartate (NMDA)-receptor activation diffuses to adjacent nerve terminals to modulate neurotransmitter release.²⁸ Nitric oxide may act at several levels of the nervous system to develop hyperexcitability, resulting in hyperalgesia or allodynia.^{29,30} On the other hand, nitric oxide-releasing molecules, such as nitroparacetamol (NOX-701) resulting from the combination of paracetamol and a nitrooxybutyrol moiety, has been shown to be effective in acute nociception and in neuropathic pain.^{31,32}

Under pathologic conditions (*e.g.*, during inflammation), high levels of nitric oxide are produced after iNOS expression is induced, mainly in macrophages. Nitric oxide possesses protective/destructive duality inherent in every other major component of the immune response. This labile molecule exerts beneficial effects by acting as an antibacterial, antiparasitic, antiviral agent or as a tumoricidal agent; on the other hand, high levels of nitric oxide, if

uncontrolled, elicits detrimental effects that are produced because nitric oxide reacts with concomitantly produced superoxide anions, thereby generating highly toxic compounds such as peroxynitrite and hydroxyl radicals.

Effects of Morphine on NO Synthesis and Release

Morphine on NOS

Acute and chronic morphine treatment produced an increase in Ca²⁺-dependent NOS in mouse brain; the acute effect of morphine was blocked by coadministration of naloxone.³³ In rat spinal cord, repeated administration of morphine increased the NOS mRNA level, the effect being accompanied by an increase in both the number of NOS-positive cells and the optical density of NOS-immunoreactivity, indicating that repeated morphine administration increases NOS biosynthesis.³⁴ At 24 h after treatment with morphine (25 mg in pellet), there was decreased NOS activity in all brain regions and the spinal cord of the mouse, and NOS activity increased at 48 and 72 h after the treatment in both the cerebellum and cortex; implantation of a naltrexone pellet in conjunction with a morphine pellet blocked the changes in NOS activity.³⁵ The authors suggested that the initial decrease in NOS activity is related to enhanced motor activity, whereas the increase in NOS activity is associated with tolerance-physical dependence development. Morphine increased the invertebrate immunocyte intracellular Ca²⁺ level that was mediated by μ 3-opioid receptors and was associated with stimulating nitric oxide production.³⁶ The opioid stimulation of intracellular Ca²⁺ levels appears to regulate constitutive NOS activity. Higher numbers of nNOS-positive cells were observed in the hippocampal dentate gyrus of wild-type mice repeatedly treated with either morphine or cocaine than in saline-treated wild-type mice; moreover, μ -opioid receptor knockout mice showed higher morphine- or cocaine-induced nNOS expression levels in the dentate gyrus than saline-treated wild-type mice.³⁷ The knockout mice showed a higher morphine-induced nNOS expression level or a lower cocaine-induced nNOS expression level than morphine- or cocaine-treated wild-type mice. The authors suggested that morphine and cocaine sensitization is differentially regulated by the μ -opioid receptors in the knockout mice *via* the nNOS systems in the dentate gyrus.

In contrast, Barjavel and Bhargava³⁸ demonstrated that NOS activity, as determined by the rate of conversion of [³H]arginine into [³H]citrulline, was inhibited by the κ -opioid receptor agonist U-50,488H but only at a high concentration (0.1 mM) and was not affected by selective μ - and δ -opioid receptor agonists in rat cerebral cortex, indicating that drugs acting at μ -, δ -, and κ -receptors have no direct action on central NOS activity *in vitro*. NOS activity was found to be unchanged in the brainstem and cerebellum of mice treated with morphine.³⁹

Systemic administration of diacetylmorphine reduced the number of reduced nicotinamide adenine dinucleotide phosphate (NADPH) diaphorase-positive (nitric oxide-synthesizing) neurons in the rat brain raphe nuclei, and this effect was blocked by naloxone.⁴⁰ Acute and chronic administration of morphine suppressed NADPH diaphorase-positive neurons in rat brain cervical nuclei, and naloxone reversed the morphine actions.⁴¹

Morphine on NO Release

Acute exposure of human saphenous or internal thoracic artery endothelium or rat microvascular endothelial cells to morphine resulted in nitric oxide release *via* the μ 3-opiate receptor subtype.^{42,43} Intravenous morphine increased norepinephrine, acetylcholine, and nitrite in spinal dorsal horn microdialysate in anesthetized sheep, and these effects were antagonized by intrathecal injection of the α_2 adrenoceptor blocker idazoxan, atropine, or L-NMMA.⁴⁴ Spinally released nitric oxide appears to play a role in the analgesic effects of systemic opioids. Fimiani *et al.*⁴⁵ noted that nitric oxide release was mediated through the μ 3-opioid receptor, and morphine-stimulated nitric oxide release was higher in human surgical specimens of nonsmall-cell lung carcinoma than in those of the normal lung. Zhu *et al.*⁴⁶ provided evidence that morphine biosynthesis occurs in rat brain amygdala, and morphine releases nitric oxide in limbic tissues. Morphine and dopamine induced a transient surge of nitric oxide production in endometrial glandular epithelial cells, free of endothelial cells, isolated from human endometrial specimens.⁴⁷ How increased nitric oxide release contributes to regulation of endometrial cell functions is indeed an intriguing mechanism to investigate. The μ -opioid receptor-specific antagonist β -funaltrexamine inhibited nitric oxide release from endomorphine 1-treated rodent and human immune cells.⁴⁸ There was evidence suggesting that by inhibiting nNOS and reducing nitric oxide levels, asymmetric dimethylarginine decreases μ -opioid receptor constitutive activity in mice.⁴⁹

On the other hand, Pu *et al.*⁵⁰ obtained findings that suggest the existence of a dual-control mechanism composed of the excitatory NMDA and the inhibitory μ -opioid receptors in modulating cyclic GMP/nitric oxide release in the medial preoptic area of the rat brain.

Modulation of Morphine Actions by NO Analgesia

Supraspinal Site of Action

The central analgesic effect of morphine as tested by the rat paw pressure and tail flick tests was inhibited by intracerebroventricular injection of methylene blue and potentiated by an inhibitor of cyclic GMP phosphodiesterase, but it was not blocked by the NOS inhibitors L-NMMA and *N*-iminoethyl-L-ornithine.^{51,52} Therefore, activation of the cyclic GMP system, not *via* nitric oxide

release, may be involved in the mechanism of the central analgesic effect of morphine. Antinociception induced by the muscarinic receptor agonist (+)-cis-dioxolane, but not β -endorphin, given supraspinally is likely mediated by the direct activation of a nitric oxide/cyclic GMP system, and the activation by the muscarinic agonist potentiates the antinociception induced by intracerebroventricular β -endorphin, but not that by μ -, δ -, or κ -opioid receptor agonists.⁵³ Intracerebroventricular L-NA diminished the morphine-induced analgesia, and L-arginine administered by the same route increased the analgesic effect of morphine, indicating that increased nitric oxide synthesis may potentiate morphine analgesia.⁵⁴ L-Arginine and the nitric oxide donor 3-morpholinosydnoimine intracerebroventricularly administered to mice produced antinociceptive effects that were blocked by naloxone and also by intracerebroventricular administration of a rabbit antiserum against rat dynorphin 1-13; the antinociceptive effect of L-arginine was antagonized by an inhibitor of nNOS (table 1).⁵⁵ The mechanisms for the antinociceptive action of L-arginine and 3-morpholinosydnoimine appear to be mediated by dynorphin and dependent on nitric oxide. Javanmardi *et al.*⁵⁶ noted that mesencephalic morphine antinociception was reduced when MK-801 and L-NAME were microinjected sequentially into the rostral ventromedial medulla in rats, and this reduction was not significantly different from the effects of MK-801 or L-NAME alone, implying that NMDA receptors and nitric oxide production in the rostral ventromedial medulla modulate the transmission of opioid pain-inhibitory signals from the periaqueductal gray.

On the other hand, L-NAME produced an opioid-independent antinociception in the mouse, probably by a direct effect within the brain.⁵⁷ In awake dogs, L-NA or morphine perfused through the fourth ventricle increased the nociceptive threshold, and combined morphine/L-NA perfusions produced a greater antinociceptive effect than seen when morphine was given alone, suggesting that endogenous nitric oxide, produced at supraspinal sites, acts as a nociceptive mediator.⁵⁸ Additional information in large mammals is required to extrapolate data to humans. In mice, chronic intraperitoneal administration of L-arginine decreased the antinociceptive response to subcutaneously administered morphine-6- β -D-glucuronide, a potent metabolite of morphine, whereas the response to intracerebroventricularly administered morphine-6- β -D-glucuronide was unaffected by L-arginine treatment; the decreased antinociceptive response to subcutaneous morphine-6- β -D-glucuronide was reversed by L-NA, suggesting that the decreased antinociceptive response of peripherally administered morphine-6- β -D-glucuronide by L-arginine may be related to a decrease in morphine-6- β -D-glucuronide entry into brain structures responsible for antinociceptive action.⁵⁹ Similar results were also obtained with morphine.^{60,61} Bhargava *et al.*⁶² obtained evidence suggesting that chronic intraperitoneal administration of

Table 1. Evidence for NO as an Analgesic Mediator at Supraspinal, Spinal, and Peripheral Sites in Experimental Animals

Author, Year	Species	Method	Analgesic (Route)	Treatment (Route)	Effect	Analgesic Mechanism
Pataki and Telegdy, 1998 ⁵⁴	Mouse	Tail-flick test	Morphine (icv)	L-arginine (icv)	Potential of analgesia	NO
Javanmardi <i>et al.</i> , 2005 ⁵⁶	Rat	Tail-flick test	Morphine (PAG) MK-801 (RVM)	L-NAME (RVM)	Inhibition of analgesia Inhibition of analgesia	NO NMDA receptor
Kolesnikov <i>et al.</i> , 1997 ⁶⁶	Mouse		Morphine (sc, ith)	AS to nNOS-2 (ith)	Block of analgesia	NO <i>via</i> nNOS-2
Song <i>et al.</i> , 1998 ⁶⁷	Rat	Paw withdrawal from heat source	Morphine (iv)	AS to nNOS-1 (ith) NOSI, NO scavenger or α_2 -AB (ith)	No effect Inhibition of analgesia	NO, α_2 -Adrenoceptor
Hayashida <i>et al.</i> , 2003 ⁶⁸	Rat	Formalin-test	BLF (ith) Morphine (ith) BLF + morphine*	L-NAME (ith) μ -Antagonist (ith)	Inhibition of analgesia Inhibition of analgesia	NO μ -Opioid receptor
Chen <i>et al.</i> , 2003 ⁶⁹	DM rat	Paw pressure test	DPDPE (ith)	L-NMMA (ith) or carboxy-PTIO (ith)	Inhibition of analgesia	NO
Durate <i>et al.</i> , 1990 ⁷⁶	Rat	Paw pressure test	ACh (paw) ACh and SNP (paw)	L-NMMA (ip) MY 5445 (paw)	Inhibition of analgesia Potentiation of analgesia	NO/cyclic GMP
Aquirre-Banuelos and Granados-Soto, 1999 ⁸⁰	Rat	Formalin-test	Morphine (paw) Morphine + dipyrone* (paw)	L-NAME (paw) L-NMMA (paw)	Inhibition of analgesia Inhibition of analgesia	NO NO
Nozaki and Yamamoto, 1998 ⁷⁹	Rat	Formalin-test	Morphine (paw) morphine + FK409* (paw)	carboxy-PTIO (paw)	Inhibition of analgesia	NO α_2 -Adrenoceptor
Mixcoatl-Zexuatl <i>et al.</i> , 2000 ⁸¹	Rat	Formalin-test	Morphine (paw)	L-NAME (paw) or ODQ (paw) Sildenafil (paw)	Inhibition of analgesia Potentiation of analgesia	NO/cyclic GMP
Jain <i>et al.</i> , 2003 ⁸²	Rat and mouse	Carrageenin (rat) Acetic acid-writhing (mouse)	Sildenafil (paw, ip) Morphine + sildenafil* (paw)	L-NAME (paw, ip) MB (paw, ip)	Inhibition of analgesia Inhibition of analgesia	NO Cyclic GMP
Hayashida <i>et al.</i> , 2004 ⁸³	Rat	Formalin-test	Morphine (paw) Morphine + BLF (paw)	BLF (paw) L-NAME (paw)	Potentiation of analgesia Inhibition of analgesia	NO
Ortiz <i>et al.</i> , 2005 ⁸⁴	Rat	Formalin-test	Codeine (paw)	L-NAME, MB, GLC, 4-AP, or TEA (paw)	Inhibition of analgesia	NO/cyclic GMP/ K ⁺ -channel

* Potentiation by combined use.

4-AP = 4-aminopyridine; α_2 -AB = α_2 -adrenoceptor blocker; ACh = acetylcholine; AS = antisense; BLF = bovine lactoferrin; carboxy PTIO = nitric oxide scavenger; DM = diabetes mellitus; DPDPE = [D-Pen²,D-Pen⁵]-enkephalin; FK409 = nitric oxide releaser; GLC = glibenclamide; GMP = guanosine monophosphate; icv = intracerebroventricular; ip = intraperitoneal; ith = intrathecal; iv = intravenous; L-NAME = N^G-nitro-L-arginine methylester; L-NMMA = N^G-monomethyl-L-arginine; MB = methylene blue; MY 5445 = cyclic GMP phosphodiesterase inhibitor; nNOS = neuronal nitric oxide synthase; NO = nitric oxide; NOSI = NOS inhibitor; ODQ = soluble guanylyl cyclase inhibitor; PAG = periaqueductal grey; RVM = rostral ventromedial medulla; sc = subcutaneous; SNP = sodium nitroprusside; TEA = tetraethylammonium.

L-arginine reduced the antinociceptive effect of morphine by increasing brain NOS activity and by decreasing the concentration of morphine in certain brain regions and the spinal cord in mice. Acute activation of the nitric oxide system by L-arginine administration attenuated morphine antinociception, possibly by inhibiting its uptake in central sites (midbrain and spinal cord) involved in antinociceptive actions.⁶³ Streptozotocin-induced diabetes in mice markedly decreased the antinociceptive effect of intracerebroventricularly administered morphine and in-

creased the urinary nitrite concentration; administration of aminoguanidine improved the effect of morphine and attenuated the increase in urinary nitrite concentration, indicating that an increase in nitric oxide formation by iNOS may be responsible for the observed decrease in antinociceptive effect of morphine in diabetic mice.⁶⁴

Spinal Site of Action

Nitric oxide acts as a modulator of dorsal horn spinal cord nociceptive pathways. NOS immunoreactivity was

Table 2. Evidence for NO as an Algesic Mediator at Supraspinal, Spinal, or Systemic Sites in Experimental Animals

Author, Year	Animal	Method	Analgesic (Route)	Treatment (Route)	Effect	Mechanism
Pelligrino <i>et al.</i> , 1996 ⁵⁸	Dog	Hindpaw withdrawal threshold	Morphine (icv)	L-NA (icv)	Increased nociceptive threshold	NO as a nociceptive
Grover <i>et al.</i> , 2000 ⁶⁴	DM mouse	Tail-flick test	Morphine (icv)	Aminoguanidine (ip)	Potential of analgesia	NO formed by iNOS as a nociceptive
Przewlocki <i>et al.</i> , 1993 ⁷¹	Rat	Tail-flick and paw pressure tests	Morphine (ith)	L-NAME (ith) or Hemoglobin (ith)	Potential of analgesia	NO as a nociceptive
Yamaguchi and Naito, 1996 ⁷⁰	Rat	Tail-flick on hot plate	Morphine (ith, ep, iv) L-NAME (ith, ep, iv)	SIN-1 L-NAME (ith, ep, iv)	Inhibition of analgesia Potential of analgesia	NO as a nociceptive
Machelska <i>et al.</i> , 1997 ⁷²	Rat	Tail-flick and paw pressure tests	Morphine, μ -agonist, or δ -agonist (ith)	L-NAME or 7-NI (ith)	Potential of analgesia	NO as a nociceptive μ - and δ -opioid receptor
Li and Clark, 2001 ⁷³	Mouse	Hot plate test	Morphine (ith)		Increase in cGMP production	
			Morphine (ith)	nNOS/HO-2 null	No effect on cGMP	
			Morphine (ith)	L-NAME (ith)	Inhibition of morphine-stimulated cGMP	NO as a nociceptive
				Sn-P (ith)		CO as a nociceptive
Brignola <i>et al.</i> , 1994 ⁶⁹	Mouse	Hot plate, tail-flick, and acetic acid-writhing	Morphine (icv)	L-arginine (ip) L-NMMA (ip) L-NAME (ip)	Inhibition of analgesia Block of L-arginine action	L-arginine/NO as a nociceptive

7-NI = 7-nitroindazole; CO = carbon monoxide; DM = diabetes mellitus; ep = epidural; HO-2 = heme oxygenase-2; icv = intracerebroventricular; iNOS = inducible nitric oxide synthase; ip = intraperitoneal; ith = Intrathecal; iv = intravenous; L-NA = N^G-nitro-L-arginine; L-NAME = L-NA methylester; L-NMMA = N^G-monomethyl-L-arginine; NO = nitric oxide; SIN-1 = nitric oxide donor; Sn-P = HO inhibitor.

present in both humans and rats with similar distribution, being present in primary sensory neurons of dorsal root ganglia and their afferent terminals in the dorsal horn of spinal cord.⁶⁵ nNOS-immunoreactive interneurons were found in the superficial layer of the dorsal horn and the intermediolateral cell column.

Kolesnikov *et al.*⁶⁶ noted that an antisense probe selectively targeting nNOS-2 blocked morphine analgesia and suggested that the facilitating nNOS-2 system predominates at the spinal level over the supraspinal level. Intravenous administration of morphine produced antinociception in rats, and an α_2 -adrenoceptor antagonist, NOS inhibitors, and a nitric oxide scavenger that were intrathecally injected produced attenuation of morphine-induced antinociception.⁶⁷ It appears that a spinal α_2 -adrenergic mechanism is involved in antinociception from intravenously administered morphine and that spinal nitric oxide mediates antinociception produced by morphine. Spinally applied bovine milk-derived lactoferrin (BLF) produced μ -opioid receptor-mediated analgesia that was reversed by coadministration of L-NAME in the rat formalin test, and it potentiated the analgesia induced

by morphine, suggesting that BLF acts as an enhancer of the spinal μ -opioidergic system *via* a nitric oxide-mediated mechanism.⁶⁸ Rats rendered diabetic with streptozotocin developed a mechanical hyperalgesia, and intrathecal [D-Pen2,D-Pen5]-enkephalin, a δ -opioid receptor agonist, increased the withdrawal threshold in response to noxious pressure in diabetic rats to a greater extent than in normal rats; intrathecal L-NMMA or the nitric oxide scavenger carboxy PTIO diminished the analgesic action of the δ -opioid receptor agonist in both normal and diabetic rats.⁶⁹ Spinal endogenous nitric oxide seems to contribute to the analgesic action of intrathecal [D-Pen2,D-Pen5]-enkephalin in both normal and diabetic neuropathic pain conditions. Table 1 summarizes the data indicating that morphine and nitric oxide show antinociceptive effects and that inhibitors of the nitric oxide/cyclic GMP pathway attenuate the analgesic effect of morphine or other opioids.

In contrast, there are findings indicating that NOS inhibition results in analgesic action and potentiates morphine-induced antinociception (table 2). Morphine or L-NAME given intrathecally, epidurally, or

intravenously produced antinociceptive effects as assessed by tail flick latency in response to thermal stimulation, and coadministration of small doses of L-NAME and morphine produced reductions of the median effective doses for morphine.⁷⁰ L-NAME given *via* three different routes appears to have a synergistic antinociceptive interaction with morphine in response to thermal stimulation. L-NAME and morphine coadministered intrathecally elicited a profound and long-lasting antinociception, which was abolished by intrathecal administration of the NO donor 3-morpholinolinosydnoimine.⁷¹ Intrathecally injected 7-NI and L-NAME enhanced antinociception induced by morphine or by agonists of μ - and δ -opioid receptors in rat tail-flick, paw pressure, and formalin tests; however, coadministration of L-NAME and a κ -opioid receptor agonist produced antinociception in the paw pressure test only, showing that the inhibition of spinal NOS appears to potentiate the μ - and δ -mediated spinal antinociception and, to a lesser extent, κ -mediated spinal antinociception.⁷² The use of the nNOS-selective inhibitor 7-NI excludes the possible involvement of cerebral vasoconstriction and systemic blood pressure increase in the enhancing effect of NOS inhibition on morphine actions. L-NAME lost the ability to potentiate the analgesic actions of intrathecally administered morphine in nNOS null-mutant mice, and the heme oxygenase inhibitor no longer potentiated morphine-induced analgesia in mice lacking a functional heme oxygenase gene.⁷³ In addition, the intrathecal injection of the cyclic GMP analog caused hyperalgesia in the hot plate assay; in spinal cord slices from either nNOS or heme oxygenase null-mutant mice, morphine did not stimulate cyclic GMP production. The authors suggested that spinal monoxide generation modifies the acute analgesic actions of morphine.

Behavioral responses, such as vocalization and agitation, to intrathecal injection of high-dose morphine in rats were not reversed by naloxone but were inhibited by pretreatment with NMDA-receptor antagonists and L-NAME; the intrathecal injection of morphine evoked increases in nitric oxide metabolites and glutamine in the extracellular fluid of dorsal spinal cord that were reduced by antagonists against L-NAME and NMDA receptors, suggesting that the excitatory action of high-dose intrathecal morphine may be mediated by an NMDA-nitric oxide cascade in the spinal cord.⁷⁴ These authors⁷⁵ also noted that injections of formalin into the plantar surface of the paw evoked a biphasic spinal release of nitrite/nitrate and a transient release of glutamate; these effects, together with flinching and licking/biting, were reduced by intrathecal, combined administration of L-NAME and morphine, suggesting that L-NAME may enhance morphine-induced antinociception through an increased inhibition of

nitrite/nitrate and glutamate releases evoked by formalin injection at the spinal cord level.

Peripheral Site of Action

Acetylcholine and sodium nitroprusside, which releases nitric oxide nonenzymatically, caused antinociception in the rat paw made hyperalgesic with prostaglandin E₂, and these analgesic effects were enhanced by intraplantar injection of the inhibitor of cyclic GMP phosphodiesterase MY5445 and blocked by the guanylyl cyclase inhibitor methylene blue; the analgesia induced by acetylcholine, but not sodium nitroprusside, was blocked by L-NMMA, and L-arginine had no effect on prostaglandin-induced hyperalgesia but caused analgesia in paws inflamed with carrageenin.⁷⁶ Peripheral analgesia induced by acetylcholine and morphine was potentiated by MY5445 and blocked by L-NMMA, whereas central morphine analgesia was potentiated by MY5445 but not affected by L-NMMA, suggesting that nitric oxide causes peripheral analgesia *via* stimulation of the nitric oxide/guanylyl cyclase system and that the central analgesic effect of morphine is associated with activation of the cyclic GMP system that is not mediated by nitric oxide.⁵¹ Involvement of the L-arginine/nitric oxide/cyclic GMP pathway in peripheral morphine analgesia was also reported.^{77,78} FK409, a nitric oxide releaser, alone had no effect on the number of flinches induced by formalin injection in rats; however, when administered after intraplantar morphine, FK409 depressed the agitation behavior, and this inhibitory effect was reversed by naloxone and carboxy-PTIO.⁷⁹ In the rat formalin test, morphine (10 μ g, ipsilateral intraplantar injection) produced antinociception. However, contralateral injection of morphine did not produce any antinociceptive effect, an indication that the local administration of morphine did not result in a systemic drug distribution.⁸⁰ Moreover, in the dipyrrone-morphine combination study, the dose of morphine used (1.25 μ g) was lower than the dose used for the study with morphine alone. Coadministration of sildenafil, an inhibitor of phosphodiesterase-5, enhanced the antinociceptive effect of morphine; and pretreatment of the paw with L-NAME, 1H[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one, or naloxone blocked the effect of the combination of sildenafil and morphine, suggesting that sildenafil itself produces antinociception and increases that induced by morphine, probably through the inhibition of cyclic GMP degradation.⁸¹ Sildenafil also exhibited an antinociceptive effect against the paw pressure test in rats and the writhing test in mice, and pretreatment with L-NAME, methylene blue, or naloxone (intraplantar injection) blocked the effect of the sildenafil-morphine combination.⁸² Peripheral coadministration into the paw of morphine with a subeffective dose of BLF produced a potentiated antinociceptive effect compared to that of morphine alone in the rat formalin test, and this poten-

tiating effect was reversed by L-NAME and a μ -opioid receptor antagonist.⁸³ Local peripheral injection of codeine produced an antinociception in the rat formalin test, and local pretreatment of paws with L-NAME, methylene blue, ATP-sensitive K⁺ channel inhibitors (glybenclamide and tolbutamide), nonselective voltage-dependent K⁺ channel inhibitors (4-aminopyridine and tetraethylammonium), or naloxone prevented codeine-induced antinociception.⁸⁴ Codeine appears to activate the opioid receptor-nitric oxide-cyclic GMP-K⁺ channels pathway to produce its effect.

Stanojevic *et al.*⁸⁵ obtained data suggesting that the rat strain (Albino Oxford and Dark Agouti)-dependent opposing effects of β -endorphin on paw inflammation are mediated through δ - and κ -opioid receptors and probably involve changes in the production of reactive oxygen species.

Actions of Systemically Administered Morphine

On the basis of the phenyl benzoquinone-induced abdominal constriction test⁸⁶ in mice, morphine, mepyramine (H₁-receptor antagonist), and L-arginine produced antinociception; L-arginine increased the antinociceptive effect of morphine and mepyramine, and L-NAME decreased the antinociception induced by these agents in combination with L-arginine.⁸⁷ It appears that morphine and mepyramine produce peripheral antinociception through involvement of the L-arginine/nitric oxide cascade or other related pathways of nociceptive processes induced by nitric oxide. Brief and continuous footshock stress (3 min) induced a naloxone-insensitive antinociception that was not altered by either L-NAME, aminoguanidine, or L-arginine in mice; in contrast, prolonged and intermittent footshock (30 min) induced a naloxone-reversible antinociceptive effect that was blocked by L-NAME but not by aminoguanidine or L-arginine. In addition, morphine increased the antinociceptive effect of prolonged footshock, and this increase was inhibited by L-NAME but not by aminoguanidine.⁸⁸ Based on these observations, the authors concluded that nitric oxide of constitutive origin may be selectively involved in an opioid-mediated type of footshock stress antinociception in mice.

However, there is evidence suggesting that nitric oxide counteracts the analgesic actions of morphine. L-Arginine administered orally or intraperitoneally, but not intracerebroventricularly, reduced the antinociceptive effect of morphine assessed in mice by using the hot plate, tail flick, and acetic acid-induced writhing tests; L-NMMA and L-NAME reversed the effects of L-arginine.⁸⁹ Although L-NAME alone did not show any antinociceptive activity, it potentiated morphine-induced analgesia in mice; L-NAME dramatically augmented the analgesic effect of morphine in the late dark period at 19 h after the lights were turned on.⁹⁰ Morphine, L-NAME, or both increased the nociceptive threshold for a criterion response to thermal stimuli for rats, and exposure to a


magnetic field abolished the analgesic effects of morphine or L-NAME when applied separately but not when injected together relative to rats that had received these drugs and had been exposed to the sham field.⁹¹ Homayoun *et al.*⁹² demonstrated that acute administration of cyclosporin A, known to decrease nitric oxide production in nervous tissues, or L-NAME enhanced the antinociception induced by administration of morphine, whereas chronic pretreatment with cyclosporin A or L-NAME did not affect morphine-induced antinociception. L-NAME, but not the iNOS inhibitor L-canavanine, enhanced the pain threshold and potentiated morphine-induced analgesia in mice.⁹³ L-NAME changed the non-analgesic effect of codeine to highly significant analgesia, and naloxone abolished the L-NAME-codeine-induced analgesia, indicating that the nitric oxide modulatory effect on the opioid analgesic codeine may be exerted, at least in part, through opioid receptors.⁹⁴ Experimental diabetes in mice decreased the antinociceptive effect of morphine, and green tea extract reversed the morphine analgesia; coadministration of green tea extract and L-NAME further increased the antinociceptive effect of morphine, the stimulating effect being attenuated by administration of L-arginine. In addition, diabetes increased plasma nitrite levels that were decreased by green tea extract.⁹⁵ Increased nitric oxide formation may be responsible for the decreased antinociceptive effect of morphine in diabetic mice, and green tea extract appears to restore the antinociceptive effect of morphine by inhibition of nitric oxide production.

The analgesic effect of morphine injected intraperitoneally exhibited biologic time-dependent differences in the thermally induced algesia in mice, whereas exogenously administered peroxynitrite exhibited either an algesic or analgesic effect, depending on the circadian time of its injection; concomitant administration of peroxynitrite and morphine reduced morphine-induced analgesia.⁹⁶

Human Studies

Randomized studies on patients with cancer pain who reported pain despite taking 80–90 mg of oral morphine daily indicated that the daily consumption of oral morphine on day 30 was greater in the group that received an additional 20 mg of oral morphine and the group that received 500 mg of oral dipyron at 6-h intervals than in the group that received a 5-mg nitroglycerin (nitric oxide donor) patch daily and the group that received 0.5 mg/kg oral ketamine at 12-h intervals, suggesting that low-dose ketamine and transdermal nitroglycerin are effective adjuvant analgesics.⁹⁷ In conjunction with their opioid tolerance-sparing function, joint delivery of ketamine or nitric oxide donors with opiates may be of benefit in cancer pain management. In another randomized, double-blind study on cancer pain patients⁹⁸ who complained of pain despite taking 80 to 90 mg of oral

Fig. 1. Possible mechanism underlying interactions between μ -opioid receptor agonists and nitric oxide (NO) generated via NMDA-receptor stimulation in antinociceptive processes. Effects of NO on tolerance to and dependence on opioids and their withdrawal syndrome are also presented. NMDA-R, *N*-methyl-D-aspartate-receptor; opioid-R, opioid receptor; $[Ca^{2+}]_i$, intracellular concentrations of Ca^{2+} ; CaM, calmodulin; BH₄, tetrahydrobiopterin; NADPH, reduced nicotinamide adenine dinucleotide phosphate; +, stimulation; -, inhibition; 7-NI, 7-nitroindazol; ADMA, asymmetrical dimethyl arginine; L-NA, N^G-nitro-L-arginine; L-NAME, L-NA methylester; nNOS, neuronal nitric oxide synthase.



morphine daily and were allowed to freely manipulate their daily morphine consumption at the time the test drug (placebo patch or 5 mg/24-h nitroglycerin patch) was administered, these authors found that daily consumption of oral morphine was smaller in the nitroglycerin group compared with the placebo group, when evaluated on the fourteenth day after initiation of the study; patients from the placebo group in general complained of somnolence, compared with the nitroglycerin group. Delivery of nitric oxide donors together with opioids may be of significant benefit in cancer pain management. This idea was further evaluated by Iohom *et al.*⁹⁹ who performed studies on patients undergoing breast surgery with axillary clearance who were randomly allocated into one of two groups: group S received a standard intraoperative and postoperative analgesic regimen (morphine, diclofenac, dextropropoxyphene + acetaminophen), and group N received a continuous paravertebral block (for 48 h) and acetaminophen and parecoxib. They found that 80% of the patients in group S and no patient in group N developed chronic postsurgical pain, and the plasma concentrations of nitrite/nitrate were greater in group N compared with group S 48 h postoperatively. Whether increased nitric oxide production is involved in the analgesic efficacy could not be determined.

Figure 1 summarizes the scheme of NMDA-receptor stimulation that mediates nNOS activation *via* an increase in intracellular Ca^{2+} concentrations and thereby the release of nitric oxide that modifies the opioidergic analgesic cascade. The nNOS activity and nitric oxide release are enhanced mainly by μ -opioid receptor stimulation. Up-to-date reports in the literature on studies employing experimental animals have led to the conclusion that nitric oxide appears to participate in potentiation of analgesia induced by morphine administered to peripheral sites (paw) *via* μ -opioid receptors; the nitric oxide/cyclic GMP pathway may be involved (table 1). However, some studies with intrathecally administered mor-

phine and NOS inhibitors have led to conclusions that endogenous nitric oxide, possibly formed through nNOS, plays significant roles as an antinociceptive agent and/or an enhancer of morphine antinociception (table 1), but other researchers report that NOS inhibitors enhance the antinociceptive effect of morphine or other opioids (table 2). Differences in animal species, NOS inhibitors, and nitric oxide scavengers employed in the various experiments do not appear to account for the opposite findings. A difference is only seen in diabetic mice, in which nitric oxide derived from iNOS acts as a nociceptive when morphine is administered intraventricularly.⁵⁴ Nitric oxide formed in large amounts by iNOS may contribute to lowering the pain threshold. Sousa and Prado¹⁰⁰ noted that low doses of the nitric oxide donor 3-morpholinonyldimethylamine reduced and higher doses enhanced or had no effect on the mechanical allodynia evoked by chronic ligature of rat sciatic nerve, suggesting that nitric oxide produces a dual effect in a model of neuropathic pain. However, it might be too speculative to postulate that excessive concentrations of nitric oxide produced through iNOS or nNOS *via* activation of NMDA receptors during nociceptive stimuli (tail-flick, hot plate tests mainly used in studies appearing in table 2) participate in promoting nociception. At present, there is only limited information available to support the beneficial use of nitric oxide as an analgesic in patients. However, data on patients are quite important because one frequently faces the problem of species variations (primate *vs.* subprimate mammals) in the actions and mechanisms of action of endogenous molecules or therapeutic compounds. More data on healthy individuals and patients are required to introduce a qualified strategy of opioidergic analgesic treatment.

Tolerance

The analgesic response to subcutaneous morphine given daily was abolished within 5 days in mice, and coadministration of L-NA with morphine prevented the

development of tolerance for at least 11 days.¹⁰¹ Together with the finding that the NMDA-receptor antagonist MK-801 prevented the development of tolerance to morphine, the authors suggested that morphine tolerance involves the activation of NMDA receptors followed by the subsequent release of nitric oxide. Preventive effects of competitive and noncompetitive NMDA antagonists and L-NA against the development of morphine tolerance were also reported.¹⁰² Administration of L-NA or L-NAME along with morphine prevented the development of tolerance to morphine and also attenuated some signs of morphine dependence in mice.¹⁰³ Downregulation by antisense treatment of nNOS-1 prevented the development of morphine tolerance in mice.⁶⁶ Neurons in rat locus coeruleus expressed opioid receptors, and these neurons exhibited tolerance to chronic administration of opioids; the average median effective dose (obtained from curves for morphine dose *vs.* single cell extracellular activity response) for morphine of the locus coeruleus cells from rats who received L-NA injections and morphine pellets was similar to that in cells from control rats, and the median effective dose of cells from morphine pelletted rats who received saline injections was substantially higher.¹⁰⁴ Nitric oxide inhibition appears to attenuate the development of tolerance to morphine in locus coeruleus neurons. Cyclosporin A that decreases nitric oxide production in nervous tissues, L-NAME, and a combination of the two at *per se* noneffective doses inhibited the induction and expression of tolerance to morphine-induced antinociception, and aminoguanidine did not alter morphine tolerance, suggesting the involvement of decreasing nitric oxide production through constitutive NOS, but not iNOS, in the modulation of morphine tolerance.¹⁰⁵

After implantation of morphine pellets in mice, the analgesic response was abolished on the third day, coadministration of L-NA with the pellets markedly retarded the development of tolerance, and L-NA slowly reversed preexisting tolerance; L-NA did not prevent tolerance to analgesia mediated by a $\kappa 1$ or $\kappa 3$ agonist.¹⁰⁶ Multiple injections of [D-Pen2,D-Pen5]enkephalin ($\delta 1$ -opioid receptor agonist), deltorphin II ($\delta 2$ -opioid receptor agonist), or morphine resulted in the development of tolerance to their analgesic action in mice; concurrent administration of L-NA or L-NMMA had no effect on the development of [D-Pen2,D-Pen5]enkephalin or deltorphin tolerance but inhibited the development of morphine tolerance.^{107,108} Development of tolerance to the antinociceptive activity of morphine and the κ -opioid receptor agonist U-50,488H was inhibited by 7-NI, which did not modify the development of tolerance to the activity of [D-Pen2,D-Pen5]enkephalin in mice.¹⁰⁹ It appears that inhibition of nNOS activity inhibits tolerance to the antinociceptive activity of μ - and κ -opioid receptor agonists but not δ -opioid receptor agonists. Develop-

ment of acute antinociceptive tolerance to intracerebroventricular morphine in mice was blocked by L-NAME, L-NMMA, 7-NI, and 3-bromo-7-NI and also by a guanylyl cyclase inhibitor.¹¹⁰ Nitric oxide formed by nNOS acting through the cyclic GMP pathway appears to mediate the development of acute antinociceptive tolerance. Subchronic administration of 7-NI attenuated the development of morphine tolerance to the cellular and analgesic effects of μ -opioid receptor agonists in rats.¹¹¹ Herrera-Baranda *et al.*¹¹² provided evidence indicating that κ -opioid receptors and nNOS in the same intracellular network of the rat periaqueductal gray appears to control the development of morphine tolerance and dependence. Blockade of nitric oxide overproduction, the consequence of NMDA-receptor activation by aminoguanidine *via* inhibition of iNOS, attenuated the development of morphine tolerance and dependence.¹¹³

In mice implanted subcutaneously with morphine pellets, NOS activity measured from the rate of conversion of [³H]arginine to [³H]citrulline in the cerebral cortex and cerebellum increased; morphine-implanted mice had higher Vmax values, but the Km values did not differ from those of control mice.¹¹⁴ Chronic treatment with morphine seems to increase NOS activity in the brain without modifying its substrate affinity. L-Arginine accelerated tolerance when it was coadministered with morphine and decreased morphine's potency in mice when given alone.³⁹ In superfused rat hippocampal slices, the amplitude of population spikes recorded by a glass microelectrode was increased by morphine; after continuous morphine superfusion, this effect was deteriorated (*i.e.*, tolerance developed), which also increased the nitrite level in the superfusate; cosuperfusion of L-arginine with morphine further increased the nitrite level and facilitated the development of morphine tolerance.¹¹⁵ Morphine administration increased the nitric oxide production in hippocampal microdialysate in rats; the time course of altered nitric oxide production coincided with the development of antinociceptive tolerance.¹¹⁶ After sustained morphine administration, nNOS-deficient mice exhibited less tolerance development compared to the control group, although measurable tolerance still occurred; mice deficient in endothelial NOS showed a degree of tolerance similar to that of the control animals.¹¹⁷ In addition, tolerance development appears to be predominantly a nitric oxide-mediated process, but it is likely augmented by a secondary (non-nitric oxide) pathway. Heinzen *et al.*¹¹⁸ provided evidence for important nitric oxide-induced alterations in μ -opioid receptor functionality that directly lead to the development of opioid antinociceptive tolerance. Asymmetric dimethylarginine, a major circulating form of methylarginines in humans and animals, competes with L-arginine for the active site of NOS isoforms.¹¹⁹ Kielstein *et al.*¹²⁰ noted that increased nitric oxide production in mice resulted in an enhanced development of tolerance

to morphine; nitric oxide increased constitutive μ -receptor activity; 7-NI attenuated morphine withdrawal in opioid-dependent rats. It was hypothesized that by inhibiting nNOS and reducing nitric oxide levels, asymmetric dimethylarginine may decrease μ -opioid receptor constitutive activity, resulting in alteration of the analgesic dose-response curve of morphine. According to Muscoli *et al.*,¹²¹ morphine-induced antinociceptive tolerance in mice was associated with increased formation of proinflammatory cytokines and oxidative DNA, and inhibition of nitric oxide synthesis or removal of superoxide blocked these biochemical changes and inhibited the development of tolerance; coadministration of morphine with a peroxynitrite decomposition catalyst attenuated protein nitration and the observed biochemical changes and prevented the development of tolerance. Peroxynitrite decomposition catalysts may have therapeutic potential as adjuncts to opioids in relieving suffering from chronic pain.

Chronic administration of morphine resulted in the development of tolerance to the analgesic action of morphine, and L-NMMA attenuated the tolerance in rats at a higher dose than in mice.¹²² Induction phase L-arginine slowed the development of opioid tolerance and physical dependence, while L-NAME and L-NMMA led to a higher degree of tolerance but had no effect on the development of physical dependence; expression-phase NOS inhibition attenuated morphine tolerance and reduced the incidence of withdrawal signs.¹²³ A 5-day treatment with increasing doses of morphine in rats induced antinociceptive tolerance, which was attenuated by L-NAME, whereas tolerance to the effect of morphine on thyrotropin (decrease) and prolactin (increase) levels was not modified by L-NAME.¹²⁴ Both nitric oxide-dependent and nitric oxide-independent mechanisms may be involved in the development of tolerance to the various effects of morphine. Additional experimentation is needed to determine the mechanism underlying the nitric oxide action on morphine tolerance.

On the other hand, combined administration of BLF with morphine retarded the development of tolerance to morphine in mice; this effect of BLF was partially blocked by L-NAME or methylene blue and completely blocked by 7-NI, suggesting that BLF blocks the development of tolerance to morphine, possibly *via* the selective activation of nNOS.¹²⁵

Spinal Tolerance

Intrathecal coadministration of L-NAME with spinal morphine produced only a small attenuation of tolerance in rats, showing little effect of nitric oxide on morphine tolerance at spinal sites.¹²⁶ There was evidence indicating that spinal cyclooxygenase activity, and to a lesser extent NOS activity, may contribute to the development and expression of opioid tolerance.¹²⁷ In a rat spinal model, coadministration of MK-801 inhibited the development of morphine tolerance; the binding affinity of

[³H]MK-801 was higher in lumbar spinal cords of morphine-tolerant rats than control rats, and constitutive expression of nNOS protein was also higher in the morphine-tolerant group, this upregulation being prevented by MK-801.¹²⁸ The authors suggested that morphine tolerance affects NMDA-receptor binding activity and increases nNOS expression in rat spinal cord. Pretreatment with midazolam inhibited the development of acute and chronic morphine tolerance in mice, and this was reversed by intrathecal injection of L-arginine; in chronic morphine-tolerant rats, midazolam decreased formalin-induced expression of Fos and Fos/NADPH diaphorase double-labeled neurons in the contralateral spinal cord and NADPH diaphorase-positive neurons in the bilateral spinal cord.¹²⁹ It appears that decreases in both the activity and expression of NOS (nNOS and iNOS) contribute to the inhibitory effect of midazolam on the development of morphine tolerance. Intrathecal 7-NI attenuated not only the development of morphine antinociceptive tolerance, but also the activation of p38 mitogen-activated protein kinase in the spinal microglia induced by chronic intrathecal administration of morphine.¹³⁰ Attenuation of morphine tolerance by nNOS inhibition appears to be associated with reducing p38 mitogen-activated protein kinase activation in the spinal microglia. On the basis of studies measuring the spinal gene expression of heme oxygenase, NOS, soluble guanylyl cyclase, and cyclic GMP-dependent protein kinase, Liang and Clark¹³¹ noted that the carbon monoxide/nitric oxide-cyclic GMP signaling pathway was upregulated after chronic morphine exposure in mice. Xu *et al.*¹³² provided evidence suggesting that activations of metabotropic glutamate receptor subtype-5 and NMDA receptors occur after the appearance of antinociceptive tolerance to morphine in rats and that the activations of these receptors appear to play a role in the development of tolerance and expression of spinal NOS through increased concentration of $[Ca^{2+}]_i$ and activation of protein kinase C.

In summary, morphine tolerance involves the activation of NMDA receptors followed by subsequent release of nitric oxide formed by nNOS but not by iNOS (fig. 1); the nitric oxide/carbon monoxide-cyclic GMP pathway may participate in the induced tolerance. Morphine tolerance also appears to be partly mediated by nitric oxide-independent mechanisms.

Dependence

Opioid tolerance and dependence are distinct phenomena, developing independently of each other.^{133,134} Administration of L-NA reduced dependence to morphine in mice with implanted morphine pellets and reversed previously established dependence.¹⁰⁶ Chronic administration of morphine resulted in the development of physical dependence as evidenced by the appearance of a variety of symptoms including a stereotyped jump-

ing response following naltrexone injection; concurrent treatment with L-NMMA inhibited the naltrexone-induced jumping response, but not other responses.¹²² L-NAME attenuated the expression phase of morphine dependence, but it did not modify the induction phase of morphine dependence and tolerance in mice.⁹³ Acute and chronic administration of agmatine that inhibits NOS activity¹³⁵ prevented morphine dependence/withdrawal in wild-type mice; in contrast, agmatine reduced only peripheral signs, not the central signs, of morphine physical dependence in nNOS knockout mice, indicating that the action of agmatine in reducing the central signs requires functional nNOS.¹³⁶ In addition to inhibition of NOS activity, the effects of agmatine are mediated through imidazoline receptors, α_2 -adrenoceptors, and blockade of NMDA receptors that may involve peripheral signs of morphine dependence.

On the other hand, Pataki and Telegdy⁵⁴ reported that neither L-NA nor L-arginine affected the signs of morphine dependence, as assessed by naloxone-precipitated withdrawal in mice. In morphine-dependent rats, 7-NI did not affect naloxone's discriminative stimulus effects, but it decreased naloxone-induced weight loss and abolished expression of withdrawal signs, such as diarrhea, scream on touch, tremor, and "wet dog" like shaking, suggesting different mechanisms for subjective and somatic components of opioid withdrawal.¹³⁷ In morphine-dependent rats, pretreatment with L-NA potentiated the cataleptic response to morphine and blocked morphine-induced hyperthermia.¹³⁸ In the induction phase of morphine dependence, the α_2 -adrenoceptor agonist clonidine intensified and yohimbine attenuated the degree of dependence; L-NAME did not affect the development of dependence, but it potentiated the effect of clonidine.¹³⁹ The α_2 -adrenergic pathway seems to functionally link with the nitric oxide pathway in the modulation of opioid dependence.

In heroin abusers, the levels of lipoperoxides in plasma and erythrocytes and the plasma level of nitric oxide increased with prolonged abuse and with increased daily quantity, whereas the plasma levels of vitamins C and E and β -carotene and the erythrocyte levels of superoxide dismutase, catalase, and glutathione peroxidase decreased.¹⁴⁰ The balance between oxidation and antioxidation in the heroin abusers appears to be seriously impaired.

In summary, NOS inhibitors prevent morphine dependence; central signs of morphine dependence may be associated with nitric oxide derived from nNOS (fig. 1). Some investigators report that NOS inhibitors do not affect or even potentiate the signs of morphine dependence.

Withdrawal Syndrome

L-NA and L-NAME intraperitoneally administered 1 h before naloxone reduced "wet dog" shakes and weight loss that were evoked in morphine-dependent rats given

naloxone¹⁴¹ or decreased naloxone-precipitated withdrawal jumping and diarrhea in morphine-dependent mice,¹⁴² suggesting the involvement of nitric oxide in morphine withdrawal syndrome. NOS inhibition may contribute to treatment of the opioid withdrawal syndrome. Similar results were also obtained in studies using L-NAME and isosorbide dinitrate, a nitric oxide donor, injected shortly before naloxone in morphine-dependent rats.¹⁴³ Thorat *et al.*¹⁴⁴ obtained data suggesting that NOS inhibitors may be more beneficial than NMDA-receptor antagonists in managing the symptoms of morphine abstinence syndrome. L-NA and 7-NI attenuated naloxone-precipitated withdrawal signs, such as rearing, jumping, ptosis, rhinorrhoea, and irritability on touch, in morphine-dependent rats.¹⁴⁵ Mainly central, but not endothelial, nitric oxide may be involved in the expression of some opioid withdrawal symptoms. L-NA, L-NAME, 7-NI, and N^5 -(1-iminoethyl)-L-ornithine, a potent inhibitor of endothelial NOS, administered shortly before morphine withdrawal produced decreases in weight loss, diarrhea, wet dog shakes, and grooming; 7-NI also reduced mastication, salivation, and genital effects, and clonidine produced similar effects to 7-NI, indicating that the nNOS inhibitor attenuates more signs of opioid withdrawal than inhibitors of other types of NOS without causing hypertension.¹⁴⁶ Insofar as hypertension is a component of opioid withdrawal in humans, the ability of 7-NI to attenuate morphine withdrawal in rats without eliciting a vasopressor response suggests that 7-NI may have human therapeutic potential.¹⁴⁷ These authors¹⁴⁸ also obtained data suggesting that constitutive NOS isoforms, but not iNOS, have a primary role in nitric oxide-mediated processes that modulate the opioid withdrawal syndrome in the rat. There was evidence suggesting that nNOS and phospholipase A₂, but not iNOS, play an important role in the expression of morphine-induced withdrawal syndrome in mice, possibly by increasing free radicals.¹⁴⁹ When administered intracerebroventricularly, both L-NA and L-NMMA inhibited naltrexone-induced stereotyped jumping behavior in morphine-dependent mice.¹⁵⁰ Brain nitric oxide likely plays an important role in the expression of behavioral signs of morphine withdrawal syndrome. In morphine-dependent mice, subcutaneous L-NAME reduced the number of escape jumps and other motor symptoms of abstinence, together with a decrease in NOS activity in the cerebellum, indicating a hyperactivity of the L-arginine/nitric oxide pathway in opiate withdrawal.¹⁵¹ Pretreatment with subcutaneous lamotrigine, a new antiepileptic compound, reversed the withdrawal-induced increase in cerebellar Ca²⁺-dependent NOS activity and reduced the number of escape jumps and other motor symptoms of abstinence; MK-801 also showed similar effects on cerebellar NOS activity and motor symptoms.¹⁵² Treatment of morphine-dependent rats with L-NAME enhanced osmotically-stimulated oxytocin secre-

tion during naloxone-precipitated withdrawal, and sodium nitroprusside inhibited oxytocin neurons during naloxone-precipitated morphine withdrawal.¹⁵³

Systemic and intracerebroventricular pretreatment of rats with L-NAME in a dose sufficient to inhibit brain NOS activity blocked the naloxone-precipitated locus coeruleus withdrawal response as measured with an *in vivo* voltammetric approach, and it reduced the naloxone-induced increase in the catechol oxidation current signal.¹⁵⁴ Nitric oxide appears to play an intermediary role in the locus coeruleus neuronal hyperactivity associated with both acute and chronic morphine withdrawal. Jhamandas *et al.*¹⁵⁵ provided evidence by way of NADPH-diaphorase histochemistry for the activation of select populations of nitric oxide-synthesizing neurons in the paraventricular and supraoptic nuclei, and to a lesser extent in the brainstem nucleus tractus solitarius, during the opioid withdrawal syndrome. The soluble guanylyl cyclase may play an intermediary role in the genesis of locus coeruleus neuronal hyperactivity and behavioral signs of morphine withdrawal.¹⁵⁶ In anesthetized rats chronically treated with morphine, intraperitoneal L-NAME attenuated some signs of opioid withdrawal and also reduced the withdrawal-induced hyperactivity of locus coeruleus neurons; intraperitoneal 7-NI caused a complete blockade of the withdrawal-induced hyperactivity, and application of 7-NI to the vicinity of the locus coeruleus also caused a partial blockade, leading to the conclusion that opioid withdrawal may be mediated by nitric oxide acting as an intermediate messenger in the locus coeruleus.¹⁵⁷ In morphine-treated freely moving rats, acute pretreatment with 7-NI or L-NA-p-nitroaniline (nNOS inhibitors) before naltrexone challenge attenuated the behavioral expression of morphine withdrawal and reduced the withdrawal-induced increase in 3,4-dihydroxyphenylacetic acid in the rat locus coeruleus, suggesting a role for nitric oxide in the expression of morphine withdrawal syndrome that may be mediated, at least in part, by locus coeruleus noradrenergic neurons.¹⁵⁸ Other noradrenergic nuclei may also be involved in the action of NOS inhibition.

Mice first made dependent to morphine that were then withdrawn by removal of pellets followed by a sublethal dose of lipopolysaccharide exhibited 100% lethality, and these animals had elevated serum tumor necrosis factor- α and nitric oxide levels and depressed interleukin-12 levels compared to controls; anti-tumor necrosis factor- α antibody given at the same time as the lipopolysaccharide challenge afforded protection to morphine-withdrawn mice.¹⁵⁹ Morphine withdrawal may sensitize the animals to lipopolysaccharide lethality *via* increased production of tumor necrosis factor- α .

Spinal Sites of Action

Intrathecal administration of NMDA receptor antagonists, MK-801 and AP-7, reduced the expression of naloxone-precipitated cardiovascular and behavioral symp-

oms; L-NAME produced L-arginine-reversible inhibition of the cardiovascular component of withdrawal, but it had no effect on the expression of behavioral signs; in contrast, L-NA and L-NMMA inhibited only the expression of the behavioral signs.¹⁶⁰ Both spinal NMDA receptors and a nitric oxide-generating system are suggested to play a role in the expression of both the cardiovascular and behavioral components of naloxone-precipitated withdrawal. However, the fact that different structural analogs of L-arginine have different profiles of activity is puzzling. Naloxone-precipitated morphine withdrawal increased the expression of Fos protein, NADPH-diaphorase-positive neurons, and Fos/NADPH diaphorase double-labeled neurons in all the laminae of the rat spinal cord; intrathecal injection of nNOS antisense oligonucleotides inhibited the increase in Fos expression and NMDA_{1A}-receptor mRNA expression during morphine withdrawal and decreased the scores of morphine withdrawal symptoms.¹⁶¹ The authors concluded that nitric oxide seems to mediate the increase of Fos protein and NMDA_{1A}-receptor mRNA expression in the spinal cord during morphine withdrawal. On the basis of immunohistochemical studies, morphine induced c-Fos expression in the striatum, cerebral cortex, and midline/intralaminar nuclei of the thalamus in rats; expression in the striatum, but not the thalamus or cortex, was blocked by 7-NI, and there was no colocalization of c-Fos and nNOS in any brain region, suggesting a role for nNOS in the neural circuits activated by morphine.¹⁶² Cao *et al.*¹⁶³ provided evidence supporting the idea that cross talk between nitric oxide and the extracellular signal regulated kinase 1 and 2 signaling pathway mediates morphine withdrawal and withdrawal-induced spinal neuronal sensitization in morphine-dependent rats. Intrathecal injection of muscarinic M₂-receptor antisense oligonucleotides decreased the scores of morphine withdrawal symptoms; the expression of nNOS-positive neurons in the locus coeruleus increased in morphine-dependent rats and increased to a greater extent during morphine withdrawal, and intrathecal injection of M₂ antisense oligonucleotide inhibited the increase in nNOS expression.¹⁶⁴ M₂ muscarinic receptors of the spinal cord appear to mediate the increase of nNOS expression in the locus coeruleus during morphine withdrawal.

As presented so far, naloxone-precipitated withdrawal responses are attenuated by NOS inhibitors, possibly *via* the activation of guanylyl cyclase; nNOS-selective inhibitors are more effective than inhibitors of other types of NOS isoforms (fig. 1) and do not raise systemic blood pressure; these effects may be therapeutically beneficial for the prevention of withdrawal syndrome.

Summary and Conclusion

The current article includes a summary of the interactions between morphine/other opioids and nitric oxide

in antinociception, dependence, tolerance, and withdrawal syndrome. Some investigators have reported that NOS inhibitors attenuate the antinociceptive effects of morphine; in contrast, others have shown that NOS inhibition augments morphine-induced analgesia in experimental animals. Limited studies on humans suggest the involvement of nitric oxide in morphine-induced antinociception. Dependence on morphine is either enhanced or inhibited by endogenous nitric oxide; however, tolerance to morphine and morphine withdrawal syndrome are potentiated by nitric oxide, as so far reported. Although the reason for the controversial results on morphine-nitric oxide interactions in experimental animals remains to be determined, information included in the present review should contribute to the construction of advanced strategies for therapy with morphine, with the goals of effectively eliminating nociception and minimizing side effects. However, further, consolidated studies on human materials, healthy individuals, and patients are eagerly awaited.

References

- Furchgott RF, Zawadzki JV: The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature (Lond)* 1980; 288:373-6
- Furchgott RF: Studies on relaxation of rabbit aorta by sodium nitrite: The basis for the proposal that the acid-activatable inhibitory factor from bovine retractor penis is inorganic nitrite and the endothelium-derived relaxing factor is nitric oxide. *Vasodilatation: Vascular Smooth Muscle, Autonomic Nerves, and Endothelium*. Edited by Vanhoutte PM, New York, Raven Press, 1988, pp. 401-14
- Ignarro LJ, Buga GM, Wood KS, Byrns RE, Chaudhuri G: Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proc Natl Acad Sci U S A* 1987; 84:9265-9
- Palmer RM, Ferrige AB, Moncada S: Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature (Lond)* 1987; 327:524-6
- Bredt DS, Snyder GH: Isolation of nitric oxide synthetase, a calmodulin-requiring enzyme. *Proc Natl Acad Sci U S A* 1990; 87:682-5
- Hebel JM, White KA, Marletta MA: Purification of the inducible murine macrophage nitric oxide synthase: identification as a flavo-protein. *J Biol Chem* 1991; 266:22789-91
- Boolell M, Allen MJ, Ballard SA, Gepi-Attee S, Muirhead G, Naylor AM, Osterloch IH, Gennell C: Sildenafil: an orally active type 5 cyclic GMP-specific phosphodiesterase inhibitor for the treatment of penile erectile dysfunction. *Int J Impot Res* 1996; 8:47-52
- Toda N, Ayajiki K, Okamura T: Nitric oxide and penile erectile function. *Pharmacol Ther* 2005; 106:233-66
- Toda N, Toda H, Hatano Y: Nitric oxide: Involvement in the effects of anesthetic agents. *ANESTHESIOLOGY* 2007; 107:822-42
- Förstermann U, Pollock JS, Schmidt HH, Heller M, Murad F: Calmodulin-dependent endothelium-derived relaxing factor/nitric oxide synthase activity is present in the particulate and cytosolic fractions of bovine aortic endothelial cells. *Proc Nat Acad Sci U S A* 1991; 88:1788-92
- Palmer RMJ, Rees DD, Ashton DS, Moncada S: L-Arginine is the physiological precursor for the formation of nitric oxide in endothelium-dependent relaxation. *Biochem Biophys Res Commun* 1988; 153:1251-6
- Rees DD, Palmer RMJ, Schulz R, Hodson HF, Moncada S: Characterization of three inhibitors of endothelial nitric oxide synthase *in vivo* and *in vitro*. *Br J Pharmacol* 1990; 101:746-52
- Toda N, Minami Y, Okamura T: Inhibitory effects of L-N^G-nitro-arginine on the synthesis of EDRF and the cerebroarterial response to vasodilator nerve stimulation. *Life Sci* 1990; 47:345-51
- Vallance P, Leone A, Calver A, Collier J, Moncada S: Endogenous dimethylarginine as an inhibitor of nitric oxide synthesis. *J Cardiovasc Pharmacol* 1992; 20(Suppl12):S60-2
- Moore PK, Babbidge RC, Wallace P, Gaffen ZA, Hart SL: 7-Nitroindazole, an inhibitor of nitric oxide synthase, exhibits antinociceptive activity in the mouse without increasing blood pressure. *Br J Pharmacol* 1993; 108:296-7
- Griffiths MJ, Messent M, MacAllister RJ, Evans TW: Aminoguanidine selectively inhibits inducible nitric oxide synthase. *Br J Pharmacol* 1993; 110:963-8
- Garthwaite J, Southam E, Boulton CL, Nielsen EB, Schmidt K, Mayer B: Potential and selective inhibition of nitric oxide sensitive guanylate cyclase by IH[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-on. *Mol Pharmacol* 1995; 48:184-8
- Mixcoatl-Zecuatl T, Flores-Murrieta FJ, Granados-Soto V: The nitric oxide-cyclic GMP-protein kinase G-K⁺ channel pathway participates in the antiallo-dynamic effect of spinal gabapentin. *Eur J Pharmacol* 2006; 531:87-95
- Brito GA, Sachs D, Cunha FQ, Vale ML, Lotufo CM, Ferreira SH, Ribeiro RA: Peripheral antinociceptive effect of pertussis toxin: activation of the arginine/NO/cGMP/PKG/ATP-sensitive K channel pathway. *Eur J Neurosci* 2006; 24:1175-81
- Ortiz MI, Medina-Tat DA, Sarmiento-Heredia D, Palma-Martinez J, Granados-Soto V: Possible activation of the NO-cyclic GMP-protein kinase G-K⁺ channels pathway by gabapentin on the formalin test. *Pharmacol Biochem Behav* 2006; 83:420-7
- Scavone C, Munhoz CD, Kawamoto EM, Glezer I, de Sa Lima L, Marcourakis T, Markus RP: Age-related changes in cyclic GMP and PKG-stimulated cerebellar Na,K-ATPase activity. *Neurobiol Aging* 2005; 26:907-16
- Ramamoorthy S, Samuvel DJ, Buck ER, Rudnick G, Jayanthi LD: Phosphorylation of threonine residue 276 is required for acute regulation of serotonin transporter by cyclic GMP. *J Biol Chem* 2007; 282:11639-47
- Zhang YW, Gesmonde J, Ramamoorthy S, Rudnick G: Serotonin transporter phosphorylation by cGMP-dependent protein kinase is altered by a mutation associated with obsessive compulsive disorder. *J Neurosci* 2007; 27:10878-86
- Toda N, Okamura T: Possible role of nitric oxide in transmitting information from vasodilator nerve to cerebral arterial muscle. *Biochem Biophys Res Commun* 1990; 170:308-13
- Zheng Z, Shimamura K, Anthony TL, Travagli RA, Kreulen DL: Nitric oxide is a sensory nerve neurotransmitter in the mesenteric artery of guinea pig. *J Auton Nerv Syst* 1997; 29:137-44
- Zochodne DW, Levy D: Nitric oxide in damage, disease and repair of the peripheral nervous system. *Cell Mol Biol* 2005; 51:255-67
- Holzer P: Efferent-like roles of afferent neurons in the gut: blood flow regulation and tissue protection. *Auton Neurosci* 2006; 125:70-5
- Kiss JP, Vizi ES: Nitric oxide: a novel link between synaptic and nonsynaptic transmission. *Trends Neurosci* 2001; 24:211-5
- Riedel W, Neeck G: Nociception, pain, and antinociception: current concepts. *Z Rheumatol* 2001; 60:404-15
- Levy D, Zochodne DW: NO pain: potential roles of nitric oxide in neuropathic pain. *Pain Pract* 2004; 4:11-8
- Bargaud JL, Riffaud JP, Del Soldato P: Nitric-oxide releasing molecules: a new class of drugs with several major indications. *Curr Pharm Des* 2002; 8:201-13
- Romero-Sandoval EA, Curros-Criado MM, Gaitan G, Molina Herrero JF: Nitroparacetamol (NCX-701) and pain: first in a series of novel analgesics. *CNS Drug Rev* 2007; 13:279-95
- Leza JC, Lizasoain I, San-Martin-Clark O, Lorenzo P: Morphine-induced changes in cerebral and cerebellar nitric oxide synthase activity. *Eur J Pharmacol* 1995; 285:95-8
- Machelska H, Ziolkowska B, Mika J, Prezewlocka B, Prezewlocki R: Chronic morphine increases biosynthesis of nitric oxide synthase in the rat spinal cord. *Neuroreport* 1997; 8:2743-7
- Kumar S, Bhargava HN: Time course of the changes in central nitric oxide synthase activity following chronic treatment with morphine in the mouse: reversal by naltrexone. *Gen Pharmacol* 1997; 223-7
- Nieto-Fernandez FE, Mattocks D, Cavani F, Salzet M, Stefano GB: Morphine coupling to invertebrate immunocyte nitric oxide release is dependent on intracellular calcium transients. *Comp Biochem Physiol B Biochem Mol Biol* 1999; 123:295-9
- Yoo JH, Cho JH, Lee SY, Lee S, Loh HH, Ho IK, Jang CG: Differential effects of morphine- and cocaine-induced nNOS immunoreactivity in the dentate gyrus of hippocampus of mice lacking μ -opioid receptors. *Neurosci Lett* 2006; 395:98-102
- Barjavel MJ, Bhargava HN: Effect of opioid receptor agonists on nitric oxide synthase activity in rat cerebral cortex homogenate. *Neurosci Lett* 1994; 181:27-30
- Babey AM, Kolesnikov Y, Cheng J, Inturrisi CE, Trifilletti RR, Pasternak GW: Nitric oxide and opioid tolerance. *Neuropharmacology* 1994; 33:1463-70
- Dyuzen IV, Motavkin PA, Shorin VV: Dynamics of NADPH diaphorase activity in raphe neurons during chronic treatment with opiates. *Bull Exp Biol Med* 2001; 132:918-20
- Dyuzen IV, Deridovich II, Kurbatskii RA, Shorin VV: NO-ergic neurons of the cervical nucleus of the rat brain in normal conditions and after administration of opiates. *Neurosci Behav Physiol* 2004; 34:621-6
- Stefano GB, Hartman A, Bilfinger TV, Magazine HI, Liu Y, Casares F, Goligorsky MS: Presence of the μ 3 opiate receptor in endothelial cells. Coupling to nitric oxide production and vasodilation. *J Biol Chem* 1995; 270:30290-3
- Stefano GB, Salzet M, Bilfinger TV: Long-term exposure of human blood vessels to HIV gp120, morphine, and anandamide increases endothelial adhesion of monocytes: uncoupling of nitric oxide release. *J Cardiovasc Pharmacol* 1998; 31:862-9
- Xu Z, Tong C, Pan HL, Cerda SE, Eisenach JC: Intravenous morphine increases release of nitric oxide from spinal cord by an α -adrenergic and cholinergic mechanism. *J Neurophysiol* 1997; 78:2072-8
- Fimiani C, Arcuri E, Santoni A, Rialas CM, Bilfinger TV, Peter D, Salzet B:

- Stefano GB: μ_3 Opiate receptor expression in lung and lung carcinoma: ligand binding and coupling to nitric oxide release. *Cancer Lett* 1999; 146:45-51
46. Zhu W, Ma Y, Bell A, Esch T, Guarna M, Bilfinger TV, Bianchi E, Stefano GB: Presence of morphine in rat amygdala: evidence for the μ_3 opiate receptor subtypes *via* nitric oxide release in limbic structures. *Med Sci Monit* 2004; 10:BR433-9
47. Tseng L, Mazella J, Goligorsky MS, Rialas CM, Stefano GB: Dopamine and morphine stimulate nitric oxide release in human endometrial glandular epithelial cells. *J Soc Gynecol Investig* 2000; 7:343-7
48. Sarić A, Balog T, Sobocanec S, Marotti T: Endomorphine 1 activates nitric oxide synthase 2 activity and downregulates nitric oxide synthase 2 mRNA expression. *Neuroscience* 2007; 144:1454-61
49. Kielstein A, Tsikas D, Galloway GP, Mendelson JE: Asymmetric dimethylarginine (ADMA)—a modulator of nociception in opiate tolerance and addiction? *Nitric Oxide* 2007; 17:55-9
50. Pu S, Horvath TL, Diano S, Naftolin F, Kalra PS, Kalra SP: Evidence showing that β -endorphin regulates cyclic guanosine 3',5'-monophosphate (cGMP) efflux: anatomical and functional support for an interaction between opiates and nitric oxide. *Endocrinology* 1997; 138:1537-43
51. Ferreira SH, Duarte ID, Lorenzetti BB: Molecular base of acetylcholine and morphine analgesia. *Agents Actions Suppl* 1991; 32:101-6
52. Duarte ID, Ferreira SH: The molecular mechanism of central analgesia induced by morphine or carbachol and the L-arginine-nitric oxide-cGMP pathway. *Eur J Pharmacol* 1992; 221:171-4
53. Xu JY, Tseng LF: Role of nitric oxide/cyclic GMP in i.c.v. administered β -endorphin- and (+)-cis-dioxolane-induced antinociception in the mouse. *Eur J Pharmacol* 1994; 262:223-31
54. Pataki I, Telegdy G: Further evidence that nitric oxide modifies acute and chronic morphine actions in mice. *Eur J Pharmacol* 1998; 357:157-62
55. Chung E, Burke B, Bieber AJ, Doss JC, Ohgami Y, Quock RM: Dynorphin-mediated antinociceptive effects of L-arginine and SIN-1 (an NO donor) in mice. *Brain Res Bull* 2006; 70:245-50
56. Javanmardi K, Parviz M, Sadr SS, Keshavarz M, Minaii B, Dehpour AR: Involvement of N-methyl-D-aspartate receptors and nitric oxide in the rostral ventromedial medulla in modulating morphine pain-inhibitory signals from the periaqueductal grey matter in rats. *Clin Exp Pharmacol Physiol* 2005; 32:585-9
57. Moore PK, Oluoyomi AO, Babbidge RC, Wallace P, Hart SL: L-N^G-nitroarginine methyl ester exhibits antinociceptive activity in the mouse. *Br J Pharmacol* 1991; 102:198-202
58. Pelligrino DA, Laurito CE, VadeBoncouer TR: Nitric oxide synthase inhibition modulates the ventilatory depression and antinociceptive actions of fourth ventricular infusions of morphine in the awake dog. *ANESTHESIOLOGY* 1996; 85:1367-77
59. Bhargava HN, Bian JT: Effects of acute and chronic administration of L-arginine on the antinociceptive action of morphine-6- β -D-glucuronide. *Pharmacology* 1997; 55:165-72
60. Bhargava HN, Bian JT: N^G-nitro-L-arginine reversed L-arginine induced changes in morphine antinociception and distribution of morphine in brain regions and spinal cord of the mouse. *Brain Res* 1997; 749:351-3
61. Bian JT, Bhargava HN: Effect of chronic administration of L-arginine, N^G-nitro-L-arginine or their combination on morphine concentration in peripheral and urine of the mouse. *Gen Pharmacol* 1998; 30:753-7
62. Bhargava HN, Bian JT, Kumar S: Mechanism of attenuation of morphine antinociception by chronic treatment with L-arginine. *J Pharmacol Exp Ther* 1997; 281:707-12
63. Bhargava HN, Bian JT: Effects of acute administration of L-arginine on morphine antinociception and morphine distribution in central and peripheral tissues of mice. *Pharmacol Biochem Behav* 1998; 61:29-33
64. Grover VS, Sharma A, Singh M: Role of nitric oxide in diabetes-induced attenuation of antinociceptive effect of morphine in mice. *Eur J Pharmacol* 2000; 399:161-4
65. Terenghi G, Riveros-Moreno V, Hudson LD, Ibrahim NB, Polak JM: Immunohistochemistry of nitric oxide synthase demonstrates immunoreactive neurons in spinal cord and dorsal ganglia of man and rat. *J Neurol Sci* 1993; 118:34-7
66. Kolesnikov YA, Pan YX, Babey AM, Jain S, Wilson R, Pasternak GW: Functionally differentiating two neuronal nitric oxide synthase isoforms through antisense mapping: evidence for opposing NO actions on morphine analgesia and tolerance. *Proc Natl Acad Sci U S A* 1997; 94:8220-5
67. Song HK, Pan HL, Eisenach JC: Spinal nitric oxide mediates antinociception from intravenous morphine. *ANESTHESIOLOGY* 1998; 89:215-21
68. Hayashida K, Takeuchi T, Shimizu H, Ando K, Harada E: Lactoferrin enhances opioid-mediated analgesia *via* nitric oxide in the rat spinal cord. *Am J Physiol* 2003; 285:R302-5
69. Chen SR, Pan HL: Spinal nitric oxide contributes to the analgesic effect of intrathecal [d-pen2,d-pen5]-enkephalin in normal and diabetic rats. *ANESTHESIOLOGY* 2003; 98:217-22
70. Yamaguchi H, Naito H: Antinociceptive synergistic interaction between morphine and N^ω-nitro-L-arginine methyl ester on thermal nociceptive tests in the rats. *Can J Anaesth* 1996; 43:975-81
71. Przewlocki R, Machelska H, Przewlocka B: Inhibition of nitric oxide synthase enhances morphine antinociception in the rat spinal cord. *Life Sci* 1993; 53:PL1-5
72. Machelska H, Labuz D, Przewlocki R, Przewlocka B: Inhibition of nitric oxide synthase enhances antinociception mediated by μ , δ , and κ opioid receptors in acute and prolonged pain in the rat spinal cord. *J Pharmacol Exp Ther* 1997; 282:977-84
73. Li X, Clark JD: Spinal cord nitric oxide synthase and heme oxygenase limit morphine induced analgesia. *Mol Brain Res* 2001; 95:96-102
74. Watanabe C, Sakurada T, Okuda K, Sakurada C, Ando R, Sakurada S: The role of spinal nitric oxide and glutamate in nociceptive behaviour evoked by high-dose intrathecal morphine in rats. *Pain* 2003; 106:269-83
75. Watanabe C, Okuda K, Sakurada C, Ando R, Sakurada T, Sakurada S: Evidence that nitric oxide-glutamate cascade modulates spinal antinociceptive effect of morphine: a behavioural and microdialysis study in rats. *Brain Res* 2003; 990:77-86
76. Durate ID, Lorenzetti BB, Ferreira SH: Peripheral analgesia and activation of the nitric oxide-cyclic GMP pathway. *Eur J Pharmacol* 1990; 186:289-93
77. Ferreira SH, Duarte ID, Lorenzetti BB: The molecular mechanism of action of peripheral morphine analgesia: stimulation of the cGMP system *via* nitric oxide release. *Eur J Pharmacol* 1991b; 201:121-2
78. Granados-Soto V, Rufino MO, Gomes Lopes LD, Ferreira SH: Evidence for the involvement of the nitric oxide-cGMP pathway in the antinociception of morphine in the formalin test. *Eur J Pharmacol* 1997; 340:177-80
79. Nozaki-Taguchi N, Yamamoto T: The interaction of FK409, a novel nitric oxide releaser, and peripherally administered morphine during experimental inflammation. *Anesth Analg* 1998; 86:367-73
80. Aquirre-Banuelos P, Granados-Soto V: Evidence for a peripheral mechanism of action for the potentiation of the antinociceptive effect of morphine by dipyrone. *J Pharmacol Toxicol Methods* 1999; 42:79-85
81. Mixcoatl-Zecuatl T, Aguirre-Banuelos P, Granados-Soto V: Sildenafil produces antinociception and increases morphine antinociception in the formalin test. *Eur J Pharmacol* 2000; 400:81-7
82. Jain NK, Patil CS, Singh A, Kulkarni SK: Sildenafil, a phosphodiesterase5 inhibitor, enhances the antinociceptive effect of morphine. *Pharmacology* 2003; 67:150-6
83. Hayashida K, Takeuchi T, Harada E: Lactoferrin enhances peripheral opioid-mediated antinociception *via* nitric oxide in rats. *Eur J Pharmacol* 2004; 484:175-81
84. Ortiz MI, Castro-Olguin J, Pena-Samaniego N, Castaneda-Hernandez G: Probable activation of the opioid receptor-nitric oxide-cyclic GMP-K⁺ channels pathway by codeine. *Pharmacol Biochem Behav* 2005; 82:695-703
85. Stanojevic S, Mitic K, Vujic J, Kovacevic-Jovanovic V, Dimitrijevic M: Beta-endorphine differentially affects inflammation in two inbred rat strains. *Eur J Pharmacol* 2006; 549:157-65
86. Okun R, Liddon SC, Lasagnal L: The effects of aggregation, electric shock, and adrenergic blocking drug on inhibition of the "writhing syndrome." *J Pharmacol Exp Ther* 1963; 139:107-9
87. Abacioglu N, Ozmen R, Cakici I, Tunctan B, Kanzik I: Role of L-arginine/nitric oxide pathway in the antinociceptive activities of morphine and mepyrmine in mice. *Arzneimittelforschung* 2001; 51:977-83
88. Homayoun H, Khavandgar S, Dehpour AR: The selective role of nitric oxide in opioid-mediated foodshock stress antinociception in mice. *Physiol Behav* 2003; 79:567-73
89. Brignola G, Calignano A, Di Rosa M: Modulation of morphine antinociception in the mouse by endogenous nitric oxide. *Br J Pharmacol* 1994; 113:1372-6
90. Guney HZ, Gorgun CZ, Tunctan B, Uludag O, Hodoglugil U, Abacioglu N, Zengil H: Circadian-rhythm-dependent effects of L-N^G-nitroarginine methyl ester (L-NAME) on morphine-induced analgesia. *Chronobiol Int* 1998; 15:283-9
91. Dixon SJ, Persinger MA: Suppression of analgesia in rats induced by morphine or L-NAME but not both drugs by micro Tesla, frequency-modulated magnetic fields. *Int J Neurosci* 2001; 108:87-97
92. Homayoun H, Khavandgar S, Namiranian K, Dehpour AR: The effect of cyclosporine A on morphine tolerance and dependence: involvement of L-arginine/nitric oxide pathway. *Eur J Pharmacol* 2002; 452:67-75
93. Ozek M, Uresin Y, Gungor M: Comparison of the effects of specific and nonspecific inhibition of nitric oxide synthase on morphine analgesia, tolerance and dependence in mice. *Life Sci* 2003; 72:1943-51
94. Khattab MM, El-Hadiyah TM, Al-Shabanah OA, Raza M: Modification by L-NAME of codeine induced analgesia: possible role of nitric oxide. *Receptors Channels* 2004; 10:139-45
95. Singal A, Anjaneyulu M, Chopra K: Modulatory role of green tea extract on antinociceptive effect of morphine in diabetic mice. *J Med Food* 2005; 8:386-91
96. Altug S, Uludag O, Tunctan B, Cakici I, Zengil H, Abacioglu N: Biological time-dependent difference in effect of peroxynitrite demonstrated by the mouse hot plate pain model. *Chronobiol Int* 2006; 23:583-91
97. Lauretti GR, Lima IC, Reis MP, Prado WA, Pereira NL: Oral ketamine and transdermal nitroglycerin as analgesic adjuvants to oral morphine therapy for cancer pain management. *ANESTHESIOLOGY* 1999; 90:1528-33
98. Lauretti GR, Perez MV, Reis MP, Pereira NL: Double-blind evaluation of transdermal nitroglycerin as adjuvant to oral morphine for cancer pain management. *J Clin Anesth* 2002; 14:83-6
99. Iohom G, Abdalla H, O'Brien J, Szarvas S, Larney V, Buckley E, Butler M, Shorten GD: The associations between severity of early postoperative pain, chronic postoperative pain and plasma concentration of stable nitric oxide products after breast surgery. *Anesth Analg* 2006; 103:995-1000
100. Sousa AM, Prado WA: The dual effect of a nitric oxide donor in nociception. *Brain Res* 2001; 897:9-19

101. Kolesnikov YA, Pick CG, Pasternak GW: N^G-nitro-L-arginine prevents morphine tolerance. *Eur J Pharmacol* 1992; 221:399-400
102. Elliott K, Minami N, Kolesnikov YA, Pasternak GW, Inturrisi CE: The NMDA receptor antagonists, LY274614 and MK-804, and the nitric oxide synthase inhibitor, N^G-nitro-L-arginine, attenuate analgesic tolerance to the μ -opioid morphine but not to κ opioids. *Pain* 1994; 56:69-75
103. Majeed NH, Przewlocka B, Machelska H, Przewlocki R: Inhibition of nitric oxide synthase attenuates the development of morphine tolerance and dependence in mice. *Neuropharmacology* 1994; 33:189-92
104. Highfield DA, Grant S: N^G-nitro-L-arginine, an NOS inhibitor, reduces tolerance to morphine in the rat locus coeruleus. *Synapse* 1998; 29:233-9
105. Homayoun H, Khavandgar S, Namiranian K, Dehpour AR: The effect of cyclosporine A on morphine tolerance and dependence: involvement of L-arginine/nitric oxide pathway. *Eur J Pharmacol* 2002; 452:67-75
106. Kolesnikov YA, Pick CG, Ciszewska G, Pasternak GW: Blockade of tolerance but not to κ opioids by a nitric oxide synthase inhibitor. *Proc Natl Acad Sci USA* 1993; 90:5162-6
107. Bhargava HN, Zhao GM: Effect of nitric oxide synthase inhibition on tolerance to the analgesic action of D-Pen2,D-Pen5 enkephalin and morphine in the mouse. *Neuropeptides* 1996; 30:219-23
108. Zhao GM, Bhargava HN: Nitric oxide synthase inhibition attenuates tolerance to morphine but not to [D-Ala2, Glu4]deltorphin II, a δ 2-opioid receptor agonist in mice. *Peptides* 1996; 17:619-23
109. Bhargava HN, Cao YJ, Zhao GM: Effect of 7-nitroindazole on tolerance to morphine, U-50,488H and [D-Pen2,D-Pen5]enkephalin in mice. *Peptides* 1997; 18:797-800
110. Xu JY, Hill KP, Bidlack JM: The nitric oxide/cyclic GMP system at the supraspinal site is involved in the development of acute morphine antinociceptive tolerance. *J Pharmacol Exp Ther* 1998; 284:196-201
111. Santamarta MT, Ulibarri I, Pineda J: Inhibition of neuronal nitric oxide synthase attenuates the development of morphine tolerance in rats. *Synapse* 2005; 57:38-46
112. Herraez-Baranda LA, Carretero J, Gonzalez-Sarmiento R, Laorden ML, Milanes MV, Rodriguez RE: Evidence of involvement of the nNOS and the κ -opioid receptor in the same intracellular network of the rat periaqueductal gray that controls morphine tolerance and dependence. *Brain Res Mol Brain Res* 2005; 137:166-73
113. Abdel-Zaher AO, Hamdy MM, Aly SA, Abdel-Hady RH, Abdel-Rahman SRH: Attenuation of morphine tolerance and dependence by aminoguanidine in mice. *Eur J Pharmacol* 2006; 540:60-6
114. Bhargava HN, Kumar S, Barjavel MJ: Kinetic properties of nitric oxide synthase in cerebral cortex and cerebellum of morphine tolerant mice. *Pharmacology* 1998; 56:252-6
115. Lue WM, Su MT, Lin WB, Tao PL: The role of nitric oxide in the development of morphine tolerance in rat hippocampal slices. *Eur J Pharmacol* 1999; 383:129-35
116. Heinzen EL, Pollack GM: Pharmacodynamics of morphine-induced neuronal nitric oxide production and antinociceptive tolerance development. *Brain Res* 2004; 1023:175-84
117. Heinzen EL, Pollack GM: The development of morphine antinociceptive tolerance in nitric oxide synthase-deficient mice. *Biochem Pharmacol* 2004; 67:735-41
118. Heinzen EL, Booth RG, Pollack GM: Neuronal nitric oxide modulates morphine antinociceptive tolerance by enhancing constitutive activity of the μ -opioid receptor. *Biochem Pharmacol* 2005; 69:679-88
119. Leiper J, Vallance P: Biological significance of endogenous methylarginines that inhibit nitric oxide synthases. *Cardiovasc Res* 1999; 43:542-8
120. Kielstein A, Tsikas D, Galloway GP, Mendelson JE: Asymmetric dimethylarginine (ADMA)—a modulator of nociception in opiate tolerance and addiction? *Nitric Oxide* 2007; 17:55-9
121. Muscoli C, Cuzzocrea S, Ndengele MM, Mollace V, Porreca F, Fabrizi F, Esposito E, Masini E, Matsuschak GM, Salvemini D: Therapeutic manipulation of peroxynitrite attenuates the development of opiate-induced antinociceptive tolerance in mice. *J Clin Invest* 2007; 117:3530-9
122. Bhargava HN: Attenuation of tolerance to, and physical dependence on, morphine in the rat by inhibition of nitric oxide synthase. *Gen Pharmacol* 1995; 26:1049-53
123. Dambisiya YM, Lee TL: Role of nitric oxide in the induction and expression of morphine tolerance and dependence in mice. *Br J Pharmacol* 1996; 117:914-8
124. Rauhala P, Idanpaan-Heikkila JJ, Tuominen RK, Mannisto PT: N-nitro-L-arginine attenuates development of tolerance to antinociceptive but not to hormonal effects of morphine. *Eur J Pharmacol* 1994; 259:57-64
125. Tsuchiya T, Takeuchi T, Hayashida K, Shimizu H, Ando K, Harada E: Milk-derived lactoferrin may block tolerance to morphine analgesia. *Brain Res* 2006; 1068:102-8
126. Dunbar S, Yaksh TL: Effect of spinal infusion of L-NAME, a nitric oxide synthase inhibitor, on spinal tolerance and dependence induced by chronic intrathecal morphine in the rat. *Neurosci Lett* 1996; 207:33-6
127. Powell KJ, Hosokawa A, Bell A, Sutak M, Milne B, Quirion R, Jhamandas K: Comparative effects of cyclo-oxygenase and nitric oxide synthase inhibition on the development and reversal of spinal opioid tolerance. *Br J Pharmacol* 1999; 127:631-44
128. Wong CS, Hsu MM, Chou YY, Tao PL, Tung CS: Morphine tolerance increases [³H]MK-904 binding affinity and constitutive neuronal nitric oxide synthase expression in rat spinal cord. *Br J Anaesth* 2000; 85:587-91
129. Cao JL, Ding HL, He JH, Zheng LC, Duan SM, Zeng YM: The spinal nitric oxide involved in the inhibitory effect of midazolam on morphine-induced analgesia tolerance. *Pharmacol Biochem Behav* 2005; 80:493-503
130. Liu W, Wang CH, Cui Y, Mo LQ, Zhi JL, Sun SN, Wang YL, Yu HM, Zhao CM, Feng JQ, Chen PX: Inhibition of neuronal nitric oxide synthase antagonizes morphine antinociceptive tolerance by decreasing activation of p38 MAPK in the spinal microglia. *Neurosci Lett* 2006; 410:174-7
131. Liang DY, Clark JD: Modulation of the NO/CO-cGMP signaling cascade during chronic morphine exposure in mice. *Neurosci Lett* 2004; 365:73-7
132. Xu T, Jiang W, Du D, Xu Y, Zhou Q, Pan X, Lou Y, Xu L, Ma K: Inhibition of MPEP on the development of morphine antinociceptive tolerance and the biosynthesis of neuronal nitric oxide synthase in rat spinal cord. *Neurosci Lett* 2008; 436:214-8
133. Wüster M, Schulz R, Herz A: Opioid tolerance and dependence. *Trend Pharmacol Sci* 1985; 6:64-7
134. Johnson SM, Fleming WW: Mechanisms of cellular adaptive sensitivity changes: applications to opioid tolerance and dependence. *Pharmacol. Rev* 1989; 41:435-88
135. Galea E, Regunathan S, Eliopoulos V, Feinstein DL, Reis DJ: Inhibition of mammalian nitric oxide synthases by agmatine, an endogenous polyamine formed by decarboxylation of arginine. *Biochem J* 1996; 316:247-9
136. Aricioglu F, Paul IA, Regunathan S: Agmatine reduces only peripheral-related behavior signs, not the central signs, of morphine withdrawal in nNOS deficient transgenic mice. *Neurosci Lett* 2004; 354:153-7
137. Medvedev IO, Dravolina OA, Beshpalov AY: Differential effects of nitric oxide synthase inhibitor, 7-nitroindazole, on discriminative stimulus and somatic effects of naloxone in morphine-dependent rats. *Eur J Pharmacol* 1999; 377:183-6
138. Afify EA, Daabees TT, Gabra BH, Abou Zein-Har MS: Role of nitric oxide in catalepsy and hyperthermia in morphine-dependent rats. *Pharmacol Res* 2001; 44:533-9
139. Zarrindast MR, Homayoun H, Khavandgar S, Fayaz-Dastgerdi M, Fayaz-Dastgerdi M: The effects of simultaneous administration of α 2-adrenergic agents with L-NAME or L-arginine on the development and expression of morphine dependence in mice. *Behav Pharmacol* 2002; 13:117-25
140. Zhou JF, Yan ZF, Ruan ZR, Peng FY, Cai D, Yuan H, Sun L, Ding DY, Xu SS: Heroin abuse and nitric oxide, oxidation, peroxidation, lipoperoxidation. *Biomed Environ Sci* 2000; 13:131-9
141. Kimes AS, Vaupel DB, London ED: Attenuation of some signs of opioid withdrawal by inhibitors of nitric oxide synthase. *Psychopharmacology* 1993; 112:521-4
142. Cappendijk SL, de Vries R, Dzolic MR: Inhibitory effect of nitric oxide (NO) synthase inhibitors on naloxone-precipitated withdrawal syndrome in morphine-dependent mice. *Neurosci Lett* 1993; 162:97-100
143. Adams ML, Kalicki JM, Meyer ER, Cicero TJ: Inhibition of the morphine withdrawal syndrome by a nitric oxide synthase inhibitor, N^G-nitro-L-arginine methyl ester. *Life Sci* 1993; 52:PL245-9
144. Thorat SN, Barjavel MJ, Matwysyn GA, Bhargava HN: Comparative effects of N^G-monomethyl-L-arginine and MK-801 on the abstinence syndrome in morphine-dependent mice. *Brain Res* 1994; 642:153-9
145. Cappendijk SL, Duval SY, de Vries R, Dzolic MR: Comparative study of normotensive and hypertensive nitric oxide synthase inhibitors on morphine withdrawal syndrome in rats. *Neurosci Lett* 1995; 183:67-70
146. Vaupel DB, Kimes AS, London ED: Comparison of 7-nitroindazole with other nitric oxide synthase inhibitors as attenuators of opioid withdrawal. *Psychopharmacology* 1995; 118:361-8
147. Vaupel DB, Kimes AS, London ED: Nitric oxide synthase inhibitors. Preclinical studies of potential use for treatment of opioid withdrawal. *Neuro-psychopharmacology* 1995; 13:315-22
148. Vaupel DB, Kimes AS, London ED: Further *in vivo* studies on attenuating morphine withdrawal: isoform-selective nitric oxide synthase inhibitors differ in efficacy. *Eur J Pharmacol* 1997; 324:11-20
149. Mori T, Ito S, Matsubayashi K, Sawaguchi T: Comparison of nitric oxide synthase inhibitors, phospholipase A₂ inhibitor and free radical scavengers as attenuators of opioid withdrawal syndrome. *Behav Pharmacol* 2007; 18:725-9
150. Bhargava HN, Thorat SN: Evidence for a role of nitric oxide of the central nervous system in morphine abstinence syndrome. *Pharmacology* 1996; 52:86-91
151. Leza JC, Lizasoain I, Cuellar B, Moro MA, Lorenzo P: Correlation between brain nitric oxide synthase activity and opiate withdrawal. *Naunyn Schmiedeberg Arch Pharmacol* 1996; 353:349-54
152. Lizasoain I, Leza JC, Cuellar B, Moro MA, Lorenzo P: Inhibition of morphine withdrawal by lamotrigine: involvement of nitric oxide. *Eur J Pharmacol* 1996; 299:41-5
153. Bull PM, Ludwig M, Blackburn-Munro GJ, Delgado-Cohen H, Brown CH, Russell JA: The role of nitric oxide in morphine dependence and withdrawal excitation of rat oxytocin neurons. *Eur J Neurosci* 2003; 18:2545-51
154. Hall S, Milne B, Jhamandas K: Nitric oxide synthase inhibitors attenuate acute and chronic morphine withdrawal response in the rat locus coeruleus: an *in vivo* voltammetric study. *Brain Res* 1996; 739:182-91
155. Jhamandas JH, Harris KH, Petrov T, Jhamandas KH: Activation of nitric oxide-synthesizing neurons during precipitated morphine withdrawal. *Neuroreport* 1996; 7:2843-6
156. Sullivan ME, Hall SR, Milne B, Jhamandas K: Suppression of acute and

chronic opioid withdrawal by a selective soluble guanylyl cyclase inhibitor. *Brain Res* 2000; 859:45-56

157. Pineda J, Torrecilla M, Martín-Ruiz R, Ugedo L: Attenuation of withdrawal-induced hyperactivity of locus coeruleus neurons by inhibitors of nitric oxide synthase in morphine-dependent rats. *Neuropharmacology* 1998; 37:759-67

158. Javelle N, Berod A, Renaud B, Lambas-Senas L: NO synthase inhibitors attenuate locus coeruleus catecholamine metabolism and behavior induced by morphine withdrawal. *Neuroreport* 2002; 13:725-8

159. Feng P, Meissler JJ, Adler MW, Eisenstein TK: Morphine withdrawal sensitizes mice to lipopolysaccharide: elevated TNF- α and nitric oxide with decreased IL12. *J Neuroimmunol* 2005; 164:57-65

160. Buccafusco JJ, Terry AV, Shuster L: Spinal NMDA receptor-nitric oxide mediation of the expression of morphine withdrawal symptoms in the rat. *Brain Res* 1995; 679:189-99

161. Gao JL, Zeng YM, Zhang LC, Gu J, Liu HF, Zhou WH, Yang GD: NO mediated increase of Fos protein and NMDA1A R mRNA expression in rat spinal cord during morphine withdrawal. *Acta Pharmacol Sin* 2001; 22:505-11

162. Harlan RE, Webber DS, Garcia MM: Involvement of nitric oxide in morphine-induced c-Fos expression in the rat striatum. *Brain Res Bull* 2001; 54:207-12

163. Cao JL, Liu HL, Wang JK, Zeng YM: Cross talk between nitric oxide and ERK1/2 signaling pathway in the spinal cord mediates naloxone-precipitated withdrawal in morphine-dependent rats. *Neuropharmacology* 2006; 51:315-26

164. Chou WB, Zeng YM, Duan SM, Zhou WH, Gu J, Yang GD: M₂ muscarinic receptor of spinal cord mediated increase of nNOS expression in locus coeruleus during morphine withdrawal. *Acta Pharmacol Sin* 2002; 23: 691-7

■ ANESTHESIOLOGY REFLECTIONS

Bronze of M.T. "Pepper" Jenkins, M.D.



Anesthesiologist-in-Chief at Dallas' University of Texas Southwest Medical Center (UTSWMC) from 1948–1981, Marion Thomas "Pepper" Jenkins (1917–1994) served as UTSWMC's first McDermott Chair after his professorship was funded by a grateful founding family of Texas Instruments, Inc. A codesigner of teaching laryngoscopes, Dr. Jenkins received worldwide attention after he pronounced the death of President John F. Kennedy at Parkland Memorial Hospital on November 22, 1963. After serving as American Society of Anesthesiologists (ASA) president in 1972, Dr. Jenkins received Distinguished Service Awards from the ASA in 1978 and from the American Medical Association a decade later. A trustee and vice-president of the Wood Library-Museum of Anesthesiology (WLM), Pepper delivered the WLM's 1993 Lewis H. Wright Memorial Lecture, titled "Epochs in Intravenous Fluid Therapy: From the Goose Quill and Pig Bladder to Balanced Salt Solutions." Afterwards he quipped, "Yes, Pepper loves salt." Pictured here is a bronze bust, "M.T. Pepper Jenkins, M.D.," completed by his friend and colleague, surgeon-sculptor Ben Wilson, M.D., just 1 yr after Dr. Jenkins' passing. (Copyright © the American Society of Anesthesiologists, Inc. This image appears in the *Anesthesiology Reflections* online collection available at www.anesthesiology.org.)

George S. Bause, M.D., M.P.H., Honorary Curator, ASA's Wood Library-Museum of Anesthesiology, Park Ridge, Illinois, and Clinical Associate Professor, Case Western Reserve University, Cleveland, Ohio. UJYC@aol.com.