

placement of a sciatic catheter in a patient presenting an atypical response to neurostimulation. *Reg Anesth Pain Med* 2006;31: 482-3

29. Sites BD, Gallagher J, Sparks M: Ultrasound-guided popliteal block demonstrates an atypical motor response to nerve stimulation in 2 patients with diabetes mellitus. *Reg Anesth Pain Med* 2003; 28:479-82

30. Szerb J, Persaud D: Long current impulses may be required for nerve stimulation in patients with ischemic pain. *Can J Anaesth* 2005; 52:963-6

31. Adler AI, Boyko EJ, Ahroni JH, Stensel V, Forsberg RC, Smith DG: Risk factors for diabetic peripheral sensory neuropathy. Results of the Seattle Prospective Diabetic Foot Study. *Diabetes Care* 1997; 20:1162-7

32. Partanen J, Niskanen L, Lehtinen J, Mervaala E, Siitonen O, Uusitupa M: Natural history of peripheral neuropathy in patients with non-insulin-dependent diabetes mellitus. *N Engl J Med* 1995; 333:89-94

33. Koscielniak-Nielsen ZJ: Ultrasound-guided peripheral nerve blocks: What are the benefits? *Acta Anaesthesiol Scand* 2008; 52:727-37

34. Beach ML, Sites BD, Gallagher JD: Use of a nerve stimulator does not improve the efficacy of ultrasound-guided supraclavicular nerve blocks. *J Clin Anesth* 2006; 18:580-4

35. Schwemmer U, Papenfuss T, Greim C, Brederlau J, Roewer N: Ultrasound-guided interscalene brachial plexus anaesthesia: Differences in success between patients of normal and excessive weight. *Ultraschall Med* 2006; 27:245-50

36. Moayeri N, Bigeleisen PE, Groen GJ: Quantitative architecture of the brachial plexus and surrounding compartments, and their possible significance for plexus blocks. *ANESTHESIOLOGY* 2008; 108:299-304

37. Hogan Q: Finding nerves is not simple. *Reg Anesth Pain Med* 2003; 28:367-71

38. Eichenberger U, Greher M, Kirchmair L, Curatolo M, Moriggl B: Ultrasound-guided blocks of the ilioinguinal and iliohypogastric nerve: Accuracy of a selective new technique confirmed by anatomical dissection. *Br J Anaesth* 2006; 97:238-43

39. Chan V, Perlas A, Rawson R, Odukoya O: Ultrasound-guided supraclavicular brachial plexus block. *Anesth Analg* 2003; 97:1514-7

40. Maecken T, Zenz M, Grau T: Ultrasound characteristics of needles for regional anesthesia. *Reg Anesth Pain Med* 2007; 32:440-7

41. Kerr AT: The brachial plexus of nerves in man, the variations in its formation and branches. *Am J Anat* 1918; 23:285-395

42. McFarlane RM, McGrouther DA: Complications and their management, Dupuytren's Disease. Edited by McFarlane RMM, Flint DA. Edinburgh, Churchill Livingstone, 1990, pp 348-64

43. Sennwald GR: Fasciectomy for treatment of Dupuytren's disease and early complications. *J Hand Surg [Am]* 1990; 15:755-61

## ■ ANESTHESIOLOGY REFLECTIONS

### Draeger Pulmotor



Around 1908 the Draeger Oxygen Apparatus Company of Pittsburgh produced Pulmotors for resuscitating miners. The Pulmotor was designed as an automatic ventilator for pumping air or oxygen into the airways of victims of fire or mining accidents. By the 1910s, these devices were recommended for resuscitating electrocuted or drowned individuals. Rescue workers using Pulmotors were soon widely known as “Draegermen.” (Copyright © the American Society of Anesthesiologists, Inc. This image appears in the *Anesthesiology Reflections* online collection available at [www.anesthesiology.org](http://www.anesthesiology.org).)

*George S. Bause, M.D., M.P.H., Honorary Curator, ASA's Wood Library-Museum of Anesthesiology, Park Ridge, Illinois, and Clinical Associate Professor, Case Western Reserve University, Cleveland, Ohio. UJYC@aol.com.*