


10. Boretti FS, Buehler PW, D'Agnillo F, Kluge K, Glaus T, Butt OI, Jia Y, Goede J, Pereira CP, Maggiorini M, Schoedon G, Alayash AI, Schaer DJ: Sequestration of extracellular hemoglobin within a haptoglobin complex decreases its hypertensive and oxidative effects in dogs and guinea pigs. *J Clin Invest* 2009; 119:2271–80
11. Hod EA, Spitalnik SL: Stored red blood cell transfusions: Iron, inflammation, immunity, and infection. *Transfus Clin Biol* 2012; 19:84–9
12. Roback JD: Vascular effects of the red blood cell storage lesion. *Hematology Am Soc Hematol Educ Program* 2011; 2011:475–9
13. Donadee C, Raat NJ, Kanias T, Tejero J, Lee JS, Kelley EE, Zhao X, Liu C, Reynolds H, Azarov I, Frizzell S, Meyer EM, Donnenberg AD, Qu L, Triulzi D, Kim-Shapiro DB, Gladwin MT: Nitric oxide scavenging by red blood cell microparticles and cell-free hemoglobin as a mechanism for the red cell storage lesion. *Circulation* 2011; 124:465–76
14. Baek JH, D'Agnillo F, Vallelian F, Pereira CP, Williams MC, Jia Y, Schaer DJ, Buehler PW: Hemoglobin-driven pathophysiology is an *in vivo* consequence of the red blood cell storage lesion that can be attenuated in guinea pigs by haptoglobin therapy. *J Clin Invest* 2012; 122:1444–58
15. Baron DM, Yu B, Lei C, Bagchi A, Beloiartsev A, Stowell CP, Steinbicker AU, Malhotra R, Bloch KD, Zapol WM: Pulmonary hypertension in lambs transfused with stored blood is prevented by breathing nitric oxide. *ANESTHESIOLOGY* 2012; 116:637–47
16. Yu B, Lei C, Baron DM, Steinbicker AU, Bloch KD, Zapol WM: Diabetes augments and inhaled nitric oxide prevents the adverse hemodynamic effects of transfusing syngeneic stored blood in mice. *Transfusion* 2012; 52:1410–22
17. Berra L, Coppadoro A, Yu B, Lei C, Spagnolli E, Steinbicker AU, Bloch KD, Lin T, Sammy FY, Warren HS, Fernandez BO, Feelisch M, Dzik WH, Stowell CP, Zapol WM: Transfusion of stored autologous blood does not alter reactive hyperemia index in healthy volunteers. *ANESTHESIOLOGY* 2012; 117:56–63

ANESTHESIOLOGY REFLECTIONS FROM THE PIERRE VIARS MUSEUM

Epidural Anesthesia at the Pitié-Salpêtrière Hospital: From Fernand Cathelin (1901) to Jeanne Seebacher (1974)



Avec passion le Docteur Jeanne Seebacher a combattu pour la liberté de la femme et pour l'analgesie obstetricale. Surmontant bien des obstacles, elle a realisé la premiere peridurale obstetricale dans cette maternite en 1974. Son action est le sursaut de developpement et de la generalisation des techniques d'anesthesies rachidiennes en obstetricale en France.

The discovery of the epidural technique for locoregional anesthesia has been attributed to two independent French physicians, Jean Athanase Sicard (1872–1929) and Fernand Cathelin (1873–1945). However, for many French historians, it seems that the credit of the discovery should be attributed to Cathelin and that Sicard was something of an opportunist, being aware of the research conducted by Cathelin at the Pitié-Salpêtrière hospital. An image from the book of Cathelin published in 1903 is shown (*left*) and represents the main historical steps in the development of epidural injections from 1885 to 1901. In 1974, in the Pitié-Salpêtrière, the anesthesiologist Jeanne Seebacher (*top right*) introduced epidural anesthesia in obstetrics for the first time in France. In the obstetric department, a memorial tablet celebrates that event (*bottom right*): “With passion, Jeanne Seebacher struggled for the freedom of women and for obstetric analgesia. Overcoming many obstacles, she performed the first obstetric epidural anesthesia in 1974 in France in this hospital. Her action was the first step toward a widespread use of epidural and intrathecal anesthesia for obstetrical analgesia in France.”

Jean-Bernard Cazalaà, M.D., *President of Club d'Histoire de l'Anesthésie et de la Réanimation (French Association for the History of Anesthesiology and Critical Care), France (www.char-fr.net), and Musée Viars, CHU Pitié-Salpêtrière, Paris, France.*