

On tooth movement

Robert J. Isaacson, DDS, MSD, PhD; Steven J. Lindauer, DMD, MDS; and Moshe Davidovitch, DMD, MMSc

Biologically based tooth movement studies recognize that inter/intra/extra-cellular interactions are ultimately responsible for physical tooth movement. Such interactions include the 'coupling' phenomena of bone resorption and formation¹ with its associated local factors,²⁻¹⁰ systemic hormonal control of bone-cell activity,¹¹⁻¹⁵ as well as responses to endothelial,¹⁶ neural,¹⁷⁻¹⁹ inflammatory,^{20,21} and immune signals.²² Investigations have also been focused on alveolar bone as an organ.^{3,24} This characterization of bone has been analyzed employing ionic (electric) currents,^{25,26} whether stress-generated²⁷ or actively administered,^{28,29} to affect tooth movement. In addition, bone metabolism has been investigated with regard to the direct influence of force-dependent tooth movement on alveolar response.³⁰ The common factor throughout many of these types of

studies is the use of mechanical forces, or the signals they evoke, to elicit the experimental response(s).

Orthodontic tooth movements are currently described using biomechanical Newtonian mathematical models which explain tooth movement from a pure physics point of view where only two possible ways exist to apply a force system to a tooth.³¹ The first is by a single force, which in practice almost never acts through the center of resistance. Therefore, a single force results in displacement of the center of resistance in the direction of the line of the force and tipping as shown in Figure 1. This tendency for rotation is called the *moment of the force* whose magnitude is equal to the magnitude of the force multiplied by its perpendicular distance from the center of resistance of the tooth. This endows the moment of the

Abstract

Current concepts of clinical orthodontic tooth movement are misleading. Traditionally, these have been described as dependent on force levels and being tipping/bodily in nature. The following is a reevaluation of the relationships between tooth movement and orthodontic mechanotherapy not usually considered when analyzing tooth movement clinically.

It is our contention that confusion as to the biological response(s) to mechanical force is derived from *in vivo* orthodontic trials where conditions are undefined. This correlates with variable biological responses within the periodontium.

Constantly changing moment to force ratios inherent to the dynamic load deflection rates of any system used to move a tooth cause changing tendencies for crown/root tipping. This implies that the accepted notion of static translatory tooth movement, although a useful didactic concept, is currently unattainable. In actuality, it has never been documented empirically using any conventional orthodontic appliance.

Key Words

Tooth movement • Orthodontic • Mechanotherapy

Submitted: January 1993

Accepted for publication: April 1993

Angle Orthod 63:305-309

Figure 1

A two-dimensional schematic representation of a typical orthodontic point force applied away from the center of resistance. This creates a tendency for rotation referred to as the *moment of the force* (M_F), causing the resultant proportional distribution of strain in the periodontal fibers represented by the lines of tension radiating from the root. This application of a force will cause the center of resistance (o) of the tooth to move parallel to, and in the same direction as, the force. The center of rotation (X) will be located apical to the center of resistance (o).




Figure 1

force with a clockwise or counter-clockwise 'sense' (direction) around a center of rotation which will always be apical to the center of resistance of the tooth.

The second method by which tooth movement can be effected is through the application of a pair of equal forces which are parallel, noncolinear, and of opposite direction, termed a couple (Figure 2). This system, applied anywhere on a tooth, creates only a tendency for rotation referred to as the *moment of the couple* whose magnitude is equal to one of the forces of the couple multiplied by the inter-force distance. The center of rotation resulting from the moment of the couple is always coincident with the center of resistance of the tooth irrespective of its point of application.

Connoting teeth as rigid bodies, and accepting the limitation that orthodontic forces do not act directly through the center of resistance, all appliance systems must cause tooth movement through one of these two processes, either alone or in combination. The traditional paradigm describes tooth movement as rotational (tipping) and/or translational (bodily movement), as shown in Figures 1 and 3.³² Figure 1 depicts the effect of a single force whose point of application is away from the center of resistance. This configuration causes rotation or tipping, with the center of resistance moving parallel to the line of force.

The latter descriptor of tooth movement, that of translation, is somewhat more complex (Figure 3). Here, as with rotation, the force is applied away from the center of resistance, again causing rota-




Figure 2

tion or tipping. However, for translation to occur there must concurrently exist a couple with an opposite sense tending to tip the root in the opposite direction as the crown. Under these conditions, the relative amount of crown tipping (moment of the force) and root tipping (moment of the couple) expressed at any given moment in time determines the location of an instantaneous center of rotation. When these two oppositely directed moments are equal in magnitude, the center of rotation is at infinity and tooth translation occurs (Figure 3). This determinant is also expressed as the moment to force ratio.³¹⁻³³

It is important to realize that no empirical or experimental evidence has ever been reported to verify this concept of tooth translation. Any anecdotal time-lapse radiographic 'evidence' of changes in tooth position without alterations in inclination does not prove the mathematical concept of translation. With existing appliance systems, the force and its accompanying moment to tip the crown are derived from activation (deformation) of one part of the appliance system that has its own particular load deflection rate. However, the moment of the couple to tip the root is derived from one or more different components of the appliance system endowed with its own unique load deflection rate. Since the decay rates of the moment of the force and the moment of the couple are not equal, the ratio between them constantly changes.³⁴ This results in continuous repositioning of the (instantaneous) center of rotation. Therefore, *in vivo* the biomechanical concept of

Figure 3
A two-dimensional schematic representation illustrating the traditional theoretical concept of translation. The rotational tendencies of the force and couple, being opposite in direction and equal in magnitude, will nullify each other. The remaining force will tend to cause tooth movement in a bodily manner, and result in a uniform distribution of strain in the periodontal fibers as shown.




Figure 3

translation can only exist instantaneously, and cannot be sustained over time by any existing appliance system.

As a consequence, a continuously changing pattern of strain will exist throughout the periodontal ligament (Figures 1-3). This in turn influences the responses within the periodontium.^{35,36} Continuous variability in the quality/quantity of periodontal strain is not consistent or compatible with the concept of parallel tooth displacement³³ (Figure 3). In reality, the illusion of translatory tooth movement requires ongoing clinical titration of the moments of the force and the couple. In turn, the constantly changing pattern of periodontal strain should be expected to evoke variable biological responses in the periodontal ligament and its associated structures.

The orthodontic literature is replete with examples of the concept that compressive and tensile forces, with some relation to duration and intensity, elicit varying biological responses.³⁵⁻⁴⁵ This dogma is firmly rooted in the early work of Reitan (1947) where histologic 'evidence' is produced to promote the notion of "bodily tooth movement".⁴⁶ Examples are cited of fractions of the root surface area behaving as expected according to the above theory (Figure 3). However, given the previous contentions it seems just as plausible (if not more so) that when mechanical forces are used as stimuli in tooth movement studies, the specific quantity, and in some instances even the quality, of the strain transferred to any given portion of the periodontium varies

continuously and is indeterminable. Fluctuations in the quality/quantity of these forces during their application could impact further on these responses in an inconsistent manner. Moreover, the resulting temporal overlap of these responses increases the difficulty in interpreting the corresponding biological findings. This use of a variable stimulus will impact reproducibility as well as applicability of any data gathered. That the need for a constant quantity/quality of mechanical force is recognized as desirable in orthodontics^{34,47,48} is evidenced indirectly through the evolution of orthodontic materials. For example, the super-elastic property of NiTi alloy wires is touted as being able to "generate a physiologic tooth movement because of the relatively constant force delivered."⁴⁹

It is unrealistic to expect tooth displacement to occur in any manner other than crown movement exceeding root movement (i.e. the moment of the force exceeding the moment of the couple), or *vice versa*. In this way tooth movement is analogous to alternating current (AC), whereas the idealized version is likened to direct current (DC). The net result is that the periodontium is not homogeneously affected as described by an idealized model (Figure 3). In fact, this mathematical ideal probably only rarely occurs, and if/when it does it is only momentary in duration. Clinical tooth movement is accompanied by a constantly changing spectrum of strain in the periodontium that ranges from compression to tension. In order to gain an understanding of the biological mechanism(s) of tooth movement, carefully controlled appliance systems or measures of strain in the periodontium are important to meaningfully relate the stimulus of periodontal strain to the biologic events recorded.

Author Address

Dr. Robert J. Isaacson
 Department of Orthodontics
 VCU/MCV School of Dentistry
 Richmond, VA 23298-0566
 Office: (804) 786-9326

R.J. Isaacson is Professor & Chairman, Department of Orthodontics, School of Dentistry, Medical College of Virginia, Virginia Commonwealth University.

S.J. Lindauer is Assistant Professor, Department of Orthodontics, School of Dentistry, Medical College of Virginia, Virginia Commonwealth University.

M. Davidovitch is Assistant Professor, Department of Orthodontics, School of Dentistry, Medical College of Virginia, Virginia Commonwealth University.

References

1. Mundy GR, Roodman GD. Osteoclast ontogeny and function. *In*: Peck WA (ed), Bone and Mineral Research V (Elsevier: Amsterdam) 1987; 209-280.
2. Han VK, D'Ercole JD, Lund PK. Cellular localization of somatomedin (insulin-like growth factor) messenger RNA in the human fetus. *Sci* 1987; 136:193-197.
3. Canalis E, McCarthy T, Centrella M. Isolation and characterization of insulin-like growth factor I (somatomedin C) from cultures of fetal rat calvariae. *Endocrin* 1988; 122:22-27.
4. Hauschka PV, Mavrakos AE, Iafrairi MD, Doleman SE, Klagsbrun M. Growth factors in bone matrix. *J Biol Chem* 1986; 261:12665-12674.
5. Canalis E, Lorenzo J, Burgess WH, Maciag T. Effects of endothelial cell growth factor on bone remodeling *in vitro*. *J Clin Invest* 1987; 79:52-58.
6. Bertolini DR, Nedwin GE, Bringman TS, Smith DD, Mundy GR. Stimulation of bone resorption and inhibition of bone formation *in vitro* by human tumor necrosis factor. *Nature* 1986; 319:516-518.
7. Krane SM, Goldring MB, Goldring SR. Cytokines. *Ciba Foundation Symposium* 1988; 136:239-256.
8. Chambers TJ, Ali NN. Inhibition of osteoclast motility by prostaglandins I₂, E₁, E₂, and 6-oxoE₁. *J Path* 1983; 139:383-397.
9. Triffitt JT. The special proteins of bone tissue. *Clin Sci* 1987; 72:399-408.
10. Heingård D, Hultenby K, Oldberg A, Reinholt F, Wendel M. Macromolecules in bone matrix. *Conn Tis Res* 1989; 21:3-14.
11. Segre GV. Secretion, metabolism and circulating heterogeneity of parathyroid hormone. *In*: Favus MJ (ed), Primer on the metabolic bone diseases and disorders of mineral metabolism. 1990; 43-44.
12. Kronenberg HM. Parathyroid hormone: Mechanism of action. *In*: Favus MJ (ed), Primer on the metabolic bone diseases and disorders of mineral metabolism. 1990; 44-46.
13. Henry HL, Norman AW. Vitamin D: Metabolism and biological actions. *Ann Rev Nutr* 1984; 4:493-520.
14. Fisher JA, Born W. Calcitonin gene products: evolution, expression and biological targets. *Bone and Min* 1987; 2:347-353.
15. Reeve J, Zanelli JM. Parathyroid hormone and bone. *Clin Sci* 1986; 71:231-238.
16. Doty SB. *In*: Davidovitch Z (ed), Intercellular communication between blood vessels in bone and bone cells. *The Biological Mechanisms of Tooth Movement and Craniofacial Adaptation*. 1992; 73-83.
17. Davidovitch Z, Nicolay OF, Ngan PW, Shanfeld JL. Neurotransmitters, cytokines, and the control of alveolar bone remodeling in orthodontics. *Dent Clin N Am* 1988; 31:411.
18. Hohmann EL, Levine L, Tashjian AH Jr. Vasoactive intestinal peptide stimulates bone resorption via a cyclic adenosine 3',5'-monophosphate-dependent mechanism. *Endocrin* 1983; 1121:1233.
19. Hohmann EL, Elde RP, Rysavy JA, Einzig S, Gebhard RL. Innervation of periosteum and bone by sympathetic vasoactive intestinal peptide-containing nerve fibers. *Sci* 1986; 232:868.
20. Saito S, Saito M, Ngan PW, Shanfeld JL, Davidovitch Z. Interleukin 1- β and prostaglandin E are involved in the response of periodontal cells to mechanical stress *in vivo* and *in vitro*. *Am J Orthod Dentofac Orthop* 1993; In Press.
21. Dewhirst FE, Stashenko PP, Mole JE, Tsurumachi T. Purification and partial sequence of human osteoclast-activating factor: Identity with interleukin 1- β . *J Immunol* 1985; 135:2562-2568.
22. Schneider GB. Natural immunity and bone remodeling. *In*: Davidovitch Z (ed), *The Biological Mechanisms of Tooth Movement and Craniofacial Adaptation* 1992; 121-126.
23. DeAngelis V. Observations on the response of alveolar bone to orthodontic force. *Am J Orthodont* 1970; 58:284.
24. Grimm FM. Bone bending, a feature of orthodontic tooth movement. *Am J Orthod* 1972; 62:384.
25. Marino AA. Electrical stimulation in orthopedics: past, present and future. *J Bioelect* 1984; 3:235.
26. Friedenber ZB, Brighton CT. Bioelectricity and fracture healing. *Plastic Reconstr Surg* 1981; 68:435.
27. Otter M, Shoening J, Williams WS. Evidence for different sources of stress-generated potentials in wet and dry bone. *J Orthop Res* 1985; 3:321.
28. Beeson DC, Johnston LE, Wisotzky J. Effect of constant currents on orthodontic tooth movement in the cat. *J Dent Res* 1975; 54:251.
29. Davidovitch Z, Finkelson MD, Steigman S, Shanfeld JL, Montgomery PC, Korostoff E. Electric currents, bone remodeling, and orthodontic tooth movement. II. Increase in rate of tooth movement and periodontal cyclic nucleotide levels by combined force and electric current. *Am J Orthod* 1980; 77:33.
30. Storey E. Bone changes associated with tooth movement: a histological study of the effect of force in the rabbit, guinea pig, and rat. *Aust J Dent* 1955; 59:147.
31. Smith RJ, Burstone CJ. Mechanics of tooth movement. *Am J Orthod* 1986; 85:294-307.
32. Proffit WR. *Contemporary orthodontics*, 1st ed., St. Louis: CV Mosby 1986; 228-238.
33. Marcotte MR. *Biomechanics in orthodontics*, 1st ed., Philadelphia: B.C. Decker Inc. 1990; 13-15.
34. Burstone CJ. The segmented arch approach to space closure. *Am J Orthod* 1982; 82(5):361-378.
35. Reitan K. The initial tissue reaction incident to orthodontic tooth movement as related to the influence of function. *Acta Odontol Scand* 1951; suppl 6.
36. Reitan K. Tissue rearrangement during retention of orthodontically rotated teeth. *Angle Orthod* 1959; 29:105.
37. Edwards JG. A study of the periodontium during orthodontic rotation of the teeth. *Am J Orthod* 1968; 54:441.
38. Gianelly AA. Force-induced changes in the vascularity of the periodontal ligament. *Am J Orthod* 1969; 55:5-11.
39. Reitan K. Evaluation of orthodontic forces as related to histologic and mechanical factors. *Schweiz Monatsschr Zahnheilkd* 1970; 80:579.
40. Atherton JD. The gingival response to orthodontic tooth movement. *Am J Orthod* 1970; 58:179.

41. Justus R, Luft HJ. A mechanochemical hypothesis for bone remodeling induced by mechanical stress. *Calcif Tis Res* 1970; 5:222.
42. Baumrind S, Buck DL. Rate changes in cell replication and protein synthesis in the periodontal ligament incident to tooth movement. *Am J Orthod* 1970; 57:109.
43. Rygh P. Ultrastructural changes in pressure zones of human periodontium incident to orthodontic tooth movement. *Acta Odontol Scand* 1973; 31:109.
44. Storey E. Bone changes associated with tooth movement. A histological study of the effect of force for varying durations in the rabbit, guinea pig and rat. *Aust J Dent* 1955; 59:209.
45. Davidovitch Z (ed). *Biological mechanisms of tooth movement and craniofacial adaptation*. Birmingham, Alabama, EBSCO Media, 1992.
46. Reitan K. Continuous bodily tooth movement and its histological significance. *Acta Odontol Scand* 1947; 7:115.
47. Burstone CJ. The mechanics of the segmented arch techniques. *Angle Orthod* 1966; 36(2):99-120.
48. Burstone CJ, Koenig HA. Force systems from an ideal arch. *Am J Orthod* 1974; 65(3):270-289.
49. Miura F, Mogi M, Ohura Y, Hamanaka H. The super-elastic property of the Japanese NiTi alloy wire for use in orthodontics. *Am J Orthod Dentofac Orthop* 1986; 90(1):1-10.

Commentary: On tooth movement

Sunil Kapila, DDS, MS

This well-written article provides some insights into new concepts related to biomechanics and biology of tooth movement. The authors contend that currently held opinions on constant and "static" moment to force ratios and their associated types of tooth movements may not be valid since moments and forces in orthodontic force systems vary as the tooth moves and the appliance deactivates.

The concept of dynamically changing moments and forces is logical and, indeed, has been discussed previously (Burstone C. *Am J Orthod* 1982; 85:361-378). However, what the response of the periodontal tissues is to these variable moments and forces has not been adequately developed or discussed in this article. It is possible, using information currently available in orthodontic and orthopedic literature, to further develop this concept.

With regard to translational and tipping movements, one generally considers these to be the net result of an activation as observed by the clinician at the next activation visit. It is highly likely that in the time between activations a variety of types of tooth movements occur, but the end result is

dependent upon what the average moment to force ratio was over the period between activations. Therefore, although the moments and forces may vary over time, their net result between visits is largely determined by the appliance design and the activation. The above discussion highlights the differences in our understanding of the definition of the words translation and tipping tooth movements, and is not meant to detract from the concept proposed by the authors on variable moments and forces and their related tooth movements occurring between visits.

In summary, although this paper examines the concepts of biomechanics and tooth movement from a new perspective, it lacks further development of these concepts. The proposed concepts need to be further validated with information available from other current literature in orthodontics and orthopedics. Furthermore, a greater synthesis of this information would also provide a stronger rationale for the proposed concepts.

S. Kapila is Assistant Professor in the Department of Growth and Development, Division of Orthodontics, School of Dentistry, University of California, San Francisco.