

## Letters From Our Readers

**To: Editor, *The Angle Orthodontist***

**Re: Response to: Gummy smile and facial profile correction using miniscrew anchorage by Masato Kaku, Shunichi Kojima, Hiromi Sumi, Hiroyuki Koseki, Sara Abedini, Masahide Motokawa, Tadashi Fujita, Junji Ohtani, Toshitsugu Kawata, and Kazuo Tanne. *Angle Orthod.* 2012;82:170–177.**

The nature of an excessive gingival display can be divided into several categories according to etiologic factors: abnormal dental eruption; excessive vertical growth of maxilla; and muscular.

Problem.1 In this case N-A was 61.5 mm (mean value of adult female was 63.7), showing no excessive vertical growth of the maxilla. The patient's FMA was 32.3° which is a standard for Japanese, indicating that she was an average angle case. Based on this information, we diagnosed that the patient had a dentoalveolar gummy smile without abnormal vertical growth of the maxilla. However, we agree that the relaxation of the maxillary labial muscle after the retraction might contribute to the correction of a gummy smile.

Actually, we can observe root resorption of incisors by panoramic radiograph. According to the previous study, the degree of root resorption after anterior teeth intrusion was score 2.2. We think score 2 was not so severe degree if we take account the 4mm of intrusion.

*Masato Kaku*

*Department of Orthodontics and Craniofacial  
Developmental Biology, Hiroshima University  
Graduate School of Biomedical Sciences  
1-2-3 Kasumi, Minami-ku  
Hiroshima 734-8553, Japan  
mkaku@hiroshima-u.ac.jp*

### REFERENCES

1. Monaco A, Streni O, Marci MC, Marzo G, Gatto R, Giannoni M. Gummy smile: clinical parameters useful for diagnosis and therapeutical approach. *J Clin Pediatr Dent.* 2004;29:19–25.
2. Malmgren O, Goldson L, Hill C, Orwin A, Petrini L, Lundberg M. Root resorption after orthodontic treatment of traumatized teeth. *Am J Orthod.* 1982;82:487–491.