

First Occurrence of *Paramphistomum microbothrium* (Fischoeder 1901) in Roe Deer (*Capreolus capreolus*) in Serbia

Ivan Pavlović,¹ Božidar Savić,^{1,3} Snežana Ivanović,¹ and Duško Ćirović² ¹ Scientific Veterinary Institute of Serbia, Vojvode Toze 14, 11000 Belgrade, Serbia; ² Faculty of Biology, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia; ³ Corresponding author (email: sb.boban@sezampro.rs)

ABSTRACT: *Paramphistomum* parasites identified by histology as *Paramphistomum microbothrium* were found in 18 of 34 roe deer (*Capreolus capreolus*) intestines from central Serbia, examined between 1998 and 2004. This represents the first record of *P. microbothrium* in roe deer in Serbia.

Parasitic infections caused by helminths can cause serious health problems in roe deer (*Capreolus capreolus*), and they often involve nematodes (Bernard et al., 1988; Kutzer et al., 1988; Drozd and Dudzinski, 1993; Shimalov and Shimalov, 2003). Although infections with trematodes are less frequent, they can also cause serious health problems, including fasciolosis and distomatosis (Pavlović et al., 2007). Paramphistomiasis is a seldom-reported plathyhelminth infection in ruminants, including roe deer. The development of *Paramphistomum* sp. includes an intermediate host—a snail of the genus *Bulinus*. After the ingestion of the metacercaria by the final host, the development is completed after the passage through the rumen, abomasum, and small intestine (Vujić, 1965). The disease is characterized by sporadic epizootics with acute parasitic gastroenteritis, followed by high morbidity and mortality of predominantly young animals (Seck et al., 2007). In southern and Eastern Europe, the species *Paramphistomum microbothrium*, *Paramphistomum cervi*, and *Paramphistomum ichikawai* (Horak, 1971; Vishnyakov, 1980) have been recorded in domestic and wild ruminants but not in roe deer. In Serbia, *P. microbothrium* has been found both in sheep and cattle (Vujić and Petrović, 1971), as well as in red deer (*Cervus elaphus*; Pavlović et al., 2007). We report

the first findings of this parasite in roe deer from Serbia or the western Balkans.

We examined offal from 34 roe deer shot by hunters in central Serbia during the official hunting seasons in May–June and September–October, 1988–2004. Examinations were performed in the laboratory of the Scientific Veterinary Institute of Serbia. Information on animal age or sex is not available. Trachea, lungs, heart, gastrointestinal tract, liver, kidneys, and bladder were incised or opened, and all visible parasites were removed. The content of the gastrointestinal tract was sieved. Paramphistomidae parasites were fixed in 10% buffered formalin, and other helminths were fixed in Raillet and Henry's acetic formalin (920 ml of 0.8% NaCl, 30 ml of 40% formalin, 50 ml of glacial acetic acid). Selected parasites were embedded in paraffin, sectioned medio-sagittal to 5–6 µm, and stained with hematoxylin-eosin. Species determination followed Samnaliev (1981). The acetabulum was examined for determination of genera, and the genital atrium and acetabulum were examined for determination of Paramphistomidae species.

Of 34 roe deer examined, Paramphistomidae parasites were found in 18 (53%). On average, 139 (range, 38–234; standard deviation [SD], 46.72) adult parasites per host were found. The largest number of adult parasites was found in the rumen and, to a lesser extent, in the omasum and reticulum. Young parasites were found attached in a brownish-pink cluster in the mucosa of the duodenum, just distal to the pylorus, with the wall and folds so thickened that the intestinal lumen was almost completely occluded. Erosions and minor hemorrhages were visible in the




FIGURE 1. Photomicrograph of medio-sagittal section of dorsal part of the acetabulum of a *Paramphistomum microbothrium* showing the muscle series of the acetabulum (mes) and radial muscle fibers (rf). The dorsal exterior (de) and dorsal interior (di) circular muscle layers are used to identify the genus *Paramphistomum*. The number of circular muscle fibers is specific for each species of *Paramphistomum*. Dorsal external muscles are grouped into two parts: de1 contains 16–19 muscles, and de2 contains 8–14 muscles. The dorsal internal muscles contain 42–49 fibers. The number of fibers in the ventral external (ve) muscle layer is 12–16, and the number of fibers in the ventral internal (vi) layer is 45–50. The acetabulum is in a subterminal position, measuring 0.921×1.1 mm, in proportion to body length 1:3–(3.3).

mucosa, and the intestinal content was discolored red. At the primary site of infestation, the rumen, destruction of the papillae was detected, as well as hyperplasia of the epithelium and inflammatory reaction with the lymphocytes, similar to that described by Pavlović et al. (2007) and Seck et al. (2007).

Histologic examination of nine *Paramphistomidae* parasites originating from separate hosts identified them as *P. microbothrium* (Fischoeder, 1901; Eduardo, 1982a, b). The dorsal part of the acetabulum was characteristic. The dorsal circular muscle was divided into two parts, the dorsal exterior circular muscle series 1 and the dorsal exterior circular muscle series 2. These circular muscle layers are used for the determination of the genus *Param-*

phistomum (Fig. 1). The ventral exterior circular muscle series, the ventral interior circular muscle series, the radial muscle fibers, the external longitudinal muscle series, and median circular muscle series of the acetabulum specifically identified the parasites as *P. microbothrium* (Vishnyakov, 1980; Samnaliev, 1981). The prepatent period is unknown for roe deer, but it is 8 wk in cattle and 10 wk in sheep (Rangel-Ruiz et al., 2003), and under normal conditions, the complete infection cycle takes 3–4 mo.

This work was partially funded by the Serbian Ministry of Science and Technological Development (project TR 31009).

LITERATURE CITED

- BERNARD, J., W. BIESEMANS, AND P. MATHY. 1988. Nématodes parasites gastro-intestinaux des Ongulés gibier dans les Ardennes belges. *Schweizer Archiv für Tierheilkunde* 130: 77–103.
- DROZDZ, J., AND W. DUDZINSKI. 1993. Changes in the intensity of infection of roe deer, *Capreolus capreolus* (L.), with abomasum nematodes in relation to host density in a hunting ground. *Acta Parasitologica* 38: 29–32.
- EDUARDO, S. L. 1982a. The taxonomy of the family Paramphistomidae Fischoeder, 1901, with special reference to the morphology of species occurring in ruminants. I. General considerations. *Systematic Parasitology* 4: 7–57.
- . 1982b. The taxonomy of the family Paramphistomidae Fischoeder, 1901, with special reference to the morphology of species occurring in ruminants. II. Revision of the genus *Paramphistomum* Fischoeder, 1901. *Systematic Parasitology* 4: 189–238.
- FISCHODER, F. 1901. Die Paramphistomiden der Säugethiere. *Zoologischer Anzeiger* 24: 367–375.
- HORAK, I. G. 1971. Paramphistomosis of domesticated ruminants. In *Advances in parasitology*, Vol. 9, B. Davies (ed.). Academic Press, New York, New York, pp. 33–72.
- KUTZER, E., L. SUGAR, AND S. BUCHACHER-TONITZ. 1988. Contribution to the parasite fauna of wild ruminants in Hungary. II. Dynamics of parasite infections of roe deer (*Capreolus c. capreolus*). *Parasitologica Hungarica* 21: 85–97.
- PAVLOVIĆ, I., V. IVETIĆ, AND B. SAVIĆ. 2007. Occurrence of *Paramphistomum microbothrium* (Fischoeder 1901) in deer (*Cervus elaphus*). *Lucrări Științifice* 50: 601–602.
- RANGEL-RUIZ, L. J., S. T. ALBORES-BRAHMS, AND J. GAMBOA-AGUILAR. 2003. Seasonal trends of

- Paramphistomum cervi* in Tabasco, Mexico. *Veterinary Parasitology* 116: 217–222.
- SAMNALIEV, P. 1982. Morphologic and biologic difference between larvae and parthenites of *Paramphistomum microbothrium* and *Paramphistomum microbothrioides*. *Khelmintologiya*, 13: 61–66.
- SECK, M. T., B. MARCHAND, AND C. T. BA. 2007. Etude histopathologique du rumen de bovins infestés par *Carnymerius marchandi* (Gastrothylacidae) et par *Paramphistomum microbothrium* (Paramphistomidae), dans la région sud du Sénégal. *Annales de Médecine Vétérinaire* 151: 200–206.
- SHIMALOV, V. V., AND V. T. SHIMALOV. 2003. Helminth fauna of cervids in Belorussian Polesie. *Parasitology Research* 85: 75–76.
- VISHNYAKOV, Yu. I. 1980. Differential diagnosis of paramphistome infections in ruminants. *Doklady Vsesoyuzno i Akademii Selskohozyalastvenih Nauk imeni V. I. Lenin* 6: 25–26. [In Russian.]
- VUJIĆ, B. 1965. Paramphistomosis in ruminants and methods to its identification. *Veterinaria* 14: 471–478. [In Serbian.]
- VUJIĆ, B., AND Z. PETROVIĆ. 1971. A contribution to the knowledge of paramphistomides in Yugoslavia and their determination. In *Proceedings of the Première Multicolloque Européen de Parasitologie*, Rennes, France, 1–4 September 1971, Federation Européenne de Parasitologie, pp. 388–391.

Submitted for publication 3 December 2009.

Accepted 27 October 2011.