

Influence of Carbon Source, Bile Salts and Incubation Temperature on Recovery of *Enterobacteriaceae* from Foods Using MacConkey-type Agars

D. A. A. MOSSEL*, INEKE EELDERINK, MIRESE KOOPMANS and FRANCINE VAN ROSSEM

Laboratory of Microbiology, Department of Science of Food of Animal Origin, Faculty of Veterinary Medicine, The University of Utrecht, Biltstraat 172, 3572 BP Utrecht, The Netherlands

(Received for publication September 21, 1978)

ABSTRACT

The plea made many years ago (27) to replace the ill-defined coli-aerogenes ('coliform') bacteria as indicator organisms in foods processed for safety with the *Enterobacteriaceae* which are taxonomically accurately defined and as a rule more abundant has of late been more generally accepted. This called for development of a rigorously standardized formula for violet red bile glucose agar and for assessment of the optimal incubation temperature. Four reference strains of *Enterobacteriaceae*, 120 samples of minced meat and 100 samples of frozen broiler chickens were used in these studies. Considerable differences in the performance of commercially available dried formulae, when used as poured plates were observed. These applied both to productivity and to the type of colony produced by a given pure culture. As expected, replacement of lactose plus glucose by an equimolar amount of glucose did not influence the performance of the medium. Intrinsic toxicity of some batches of medium to non-stressed *Enterobacteriaceae* appeared to be mainly responsible for substandard performance. It could be overcome by careful selection of the preparations of crystal violet and particularly the bile salts (39) used in the formulae. Incubation at 30 C led to higher confirmed colony counts in minced meat than at 37 C. However, incubation temperature did not greatly influence similar counts in broiler drip. This observation could be substantiated by identification of the types of *Enterobacteriaceae* isolated from the two commodities. Psychrotrophic species predominated in minced meats, which are often made from raw materials stored for some time under refrigeration, whereas mesophilic species were in the majority on frozen broilers, which are generally frozen shortly after slaughter.

In monitoring foods processed for safety (11,33), it is advantageous to replace the coli-aerogenes bacteria, indicator organisms used in the past, by the taxonomically better defined and more abundant *Enterobacteriaceae* (38). Subsequently it has been suggested to apply *Enterobacteriaceae* counts also to unprocessed foods (4,8,19,25,44). Extreme care should be exercised in doing so. Particularly, any conclusions about lack of safety or mishandling drawn from high counts should be based on studies on the mesophilic or psychrotrophic character of the *Enterobacteriaceae* isolated (28,38,40).

For enumeration of the entire *Enterobacteriaceae* group MacConkey's agar with glucose added was originally recommended (29). For various reasons it seemed better to replace lactose in MacConkey's medium with glucose. The effect of this substitution had to be assessed experimentally. In addition, we had repeatedly

¹Brand names are presented for the purpose of identification of materials used in this investigation. Their mentioning does not imply recommendation.

observed, that the properties of the bile salts, and to a lesser extent of the triphenyl methane dyes, used in this type of medium could significantly influence its productivity (32,34,36,37,39). This investigation was designed to study this effect in more detail.

Furthermore, the temperature of incubation of MacConkey-type agar plates is worthy of consideration. Occurrence of psychrotrophic *Enterobacteriaceae* in foods has repeatedly been demonstrated (6,10,12,15,21,24,28,41,42,45,46,47,56). Incubation at 35-37 C leads to suppression of many of these strains (14) and hence to an underestimation of the degree of recontamination of products processed for safety. On the other hand, we observed that incubation at lower temperatures occasionally led to the growth of non-fermentative bacteria in MacConkey-type agars, even when these were overlaid with sterile agar (Harrewijn and Mossel, 1971, unpublished). In this investigation the influence of the incubation temperature in the range 30 to 37 C on gross and confirmed counts of typical colonies in MacConkey type agars was therefore also studied.

MATERIALS AND METHODS

Pure cultures

The pure cultures studied in this investigation originated from the collection of the Laboratory of Microbiology, Department of the Science of Food of Animal Origin. They consisted of *Escherichia coli*, *Klebsiella* sp., *Salmonella typhimurium* and *Shigella sonnei* cultures. Before use they were streaked on lactose lauryl sulfate agar (11) and incubated overnight at 37 C. After their mode of attack on lactose had been verified, their key properties were determined by deep stabbing into gram negative diagnostic tubes (36).

Minced meat samples

One hundred and twenty samples of minced meat were obtained from 10 different butcher's shops within 11 months. The minced meat was composed of approximately equal parts of pork and beef.

Samples of drip from frozen broilers

Ten samples from each of 10 different consignments of frozen broilers were defrosted for about 24 h at 4 ± 2 C. Thereupon the packs were hand-pressed (33) and about 5 ml of drip were withdrawn using a sterile syringe.

Culture media

In the pure culture studies two non-selective media, tryptone glucose yeast extract agar (Difco¹) and tryptone soya peptone agar (Oxoid¹) were used. The media were prepared following the manufacturers' instructions, and after sterilization, tempered at 47-51 C to prepare poured plates.

The violet red bile agars used in this investigation originated from four different manufacturers. They will be indicated as A, B, C and D. Media A and B contained 10 g of lactose/l (VRBL), and a further 10 g of glucose/l was added, as done previously (VRBLG). Medium C contained 10 g of glucose/l only (VRBG). From brand D two different formulae were tested, D₁ the lactose-containing formula with glucose added (VRBLG) and D₂ the medium containing glucose only (VRBG). For the remainder, the media were identical in composition, containing per l of distilled water: peptone, 7 g; yeast extract, 3 g; NaCl, 5 g; bile salts, 1.5 g; neutral red, 30 mg; crystal violet, 2 mg; and agar 15 g.

All media were prepared by bringing to a boil to dissolve the ingredients, and were not further sterilized. One brand, chosen at random (A), was also tested after sterilization for 15 min at 121 C. This medium is indicated as A(S).

Plating procedure

Pure cultures were incubated overnight at 37 C on nutrient agar (Oxoid). From these a series of decimal dilutions in peptone-saline solution was prepared in such a way, that at least one could be expected to contain 150 ± 50 cfu/ml. As a rule, three successive decimal dilutions had to be examined to attain this objective.

Minced meat samples were homogenized in 25-g portions with 250 ml of peptone-saline solution in a "stomacher" and, after allowing resuscitation of stressed cells for 2 h at room temperature (20,30,49,50,51,52,59), further decimal dilutions up to 10⁻⁴ were prepared. A series of three dilutions was examined in every experiment, again with the purpose of obtaining at least one series of duplicate plates containing approximately 150 colonies.

Broiler drip was kept for 2 h at room temperature before dilutions were made, once more to allow resuscitation of cells injured by freezing.

One-ml portions of dilutions of pure cultures, minced meat and drip were transferred to 9-cm petri dishes, thoroughly mixed with 15 ml of tempered agar, and allowed to solidify. Subsequently the plates were overlaid with 10 ml of sterile medium, tempered at 47-51 C. Incubation was done at 30 or 37 C for 18-20 h.

RESULTS

Pure culture studies

The colony counts obtained with the various media have been summarized in Table 1. An assessment of the types of colonies is presented in Table 2. This criterion is of some relevance since at one time it was thought that the diameter of the colony and of the precipitation zone around it could help in recognition of particular *Enterobacteriaceae* types (19).

It is quite obvious that considerable differences in performance of the various media occurred. The differences in the numbers of colonies remained within reasonable limits, although the recovery was usually maximal in D₁ and D₂. Considerable differences in colony size, variation in colony size and in particular the character of the precipitation zones were observed. The media D₁ and D₂ were markedly better than the other brands. The type of carbon source, lactose + glucose or only glucose did not influence the counts or colony types.

TABLE 1. Recovery of pure cultures of *Enterobacteriaceae* in a variety of MacConkey-type selective agar media.

| Strain | Number of replicates | Recovery, in vertical order min., max., mean, as fraction of colonies obtained from same inoculum in the non selective reference agar medium | | | | | |
|-------------------------------|----------------------|--|----------------|------------|--------|-------------------------|---------------------|
| | | Medium and type of sugar | | | | | |
| | | A ^a L + G ^b | A (S) L + G | B L + G | C G | D ₁ L + G | D ₂ G |
| <i>Escherichia coli</i> | 6 | | | 0.54 | 0.59 | 0.83 | 0.65 |
| | | NT | NT | 0.83 | 0.90 | 1.02 | 0.92 |
| | | | | 0.68 | 0.78 | 0.93 | 0.76 |
| <i>Klebsiella</i> | 4 | 0.42 | — | 0.30 | 0.24 | 0.49 | 0.43 |
| | | 0.42 | — | 0.64 | 0.57 | 0.50 | 0.50 |
| | | 0.42 | 0.31 | 0.47 | 0.41 | 0.50 | 0.46 |
| <i>Salmonella typhimurium</i> | 6 | NT | NT | 0.48 | 0.41 | 0.88 | 0.81 |
| | | | | 0.84 | 0.90 | 1.15 | 0.91 |
| | | | | 0.75 | 0.65 | 0.99 | 0.86 |
| <i>Shigella sonnei</i> | 4 | 0.85 | — | 1.09 | 0.83 | 1.02 | 0.87 |
| | | 0.95 | — | 1.16 | 0.96 | 1.12 | 1.11 |
| | | 0.89 | 0.98 | 1.13 | 0.90 | 1.07 | 0.99 |

^aA-D = different manufacturers (cf. text), (S) = medium sterilized instead of pasteurized, as is customary.

^bL = lactose, G = glucose, NT = not tested.

TABLE 2. Character of colonies obtained from pure cultures in a variety of MacConkey-type selective agar media.^a

| Strain | Medium and type of sugar | | | | | |
|-------------------------------|--------------------------|----------------|------------|--------|-------------------------|---------------------|
| | A L + G | A (S) L + G | B L + G | C G | D ₁ L + G | D ₂ G |
| <i>Escherichia coli</i> | | | 0.60 | 0.70 | 0.85 | 1.00 |
| | NT | NT | 0.50 | 0.50 | 1.0 | 0.87 |
| | | | 2 | 1 | 3 | 3 |
| <i>Klebsiella</i> | 0.50 | 0.50 | 0.75 | 0.50 | 1.0 | 0.75 |
| | 0 | 0 | 0.50 | 0 | 0.50 | 0.62 |
| | 0 | 0 | 2 | 0 | 3 | 2 |
| <i>Salmonella typhimurium</i> | | | 0.95 | 0.65 | 1.10 | 1.12 |
| | NT | NT | 1.2 | 0.50 | 1.0 | 1.50 |
| | | | 2 | 0 | 3 | 3 |
| <i>Shigella sonnei</i> | 0.75 | 0.75 | 1.0 | 1.0 | 1.3 | 1.12 |
| | 0 | 0.50 | 0.60 | 0.50 | 0.80 | 0.50 |
| | 0 | 0.50 | 2 | 1 | 3 | 2 |

^aParameter combination: colony size, average, mm;/ precipitation zone, do./; quality of precipitation zone, expressed in 3 = excellent, 2 = good, 1 = fair, 0 = absent.

Consequently another medium component, most probably the bile preparation (5,7,37,39) influenced performance of the MacConkey-type media.

Lactose versus glucose at 30 and 37 C

To study the effect of lactose in comparison to that of glucose, agars purchased from manufacturer D were used in all experiments, because the pure culture studies reported in the previous section indicated that much more easily countable colonies were obtained from these agars. In Tables 3 and 4 the data obtained with VRBG and VRBL are recorded. Minced meat was used as the inoculum. Colony counts were confirmed by testing the square root of the total number of colonies obtained, chosen at random, for the criteria of *Enterobacteriaceae*: negative oxidase reaction and fermentative attack on glucose. Three-layer gram negative diagnostic tubes, as described elsewhere (36) were used for this purpose.

TABLE 3. Effect of glucose on the recovery of *Enterobacteriaceae* from minced meat using VRB agars^a incubated at 37 C.

| Percentage of the samples | $\log_{10}\text{cfu}_{\text{VRBG}} - \log_{10}\text{cfu}_{\text{VRBL}}$ |
|---------------------------|---|
| 14 | 0.00 |
| 22 | 0.00-0.10 |
| 53 | 0.11-0.50 |
| 11 | > 0.50 |

Av. $\log_{10}\text{cfu}_{\text{VRBG net}} = 3.61$; av. $\log_{10}\text{cfu}_{\text{VRBL net}} = 3.53$.

^aVRBG = violet red bile glucose agar; VRBL = violet red bile lactose agar; net = confirmed as *Enterobacteriaceae*.

TABLE 4. Effect of glucose on the recovery of *Enterobacteriaceae* from minced meat using VRB-agars incubated at 30 C.

| Percentage of the samples | $\log_{10}\text{cfu}_{\text{VRBG}} - \log_{10}\text{cfu}_{\text{VRBL}}$ |
|---------------------------|---|
| 6 | 0.00 |
| 19 | 0.00-0.10 |
| 41 | 0.11-0.50 |
| 34 | > 0.50 |

Av. $\log_{10}\text{cfu}_{\text{VRBG net}} = 4.26$; av. $\log_{10}\text{cfu}_{\text{VRBL net}} = 3.92$.
Difference 0.34.

In analyzing the data, the results of earlier studies on reproducibility of counts in MacConkey type agars (27,38) were used as the base line. These had indicated that the confidence interval of such counts was of the order of a factor 2; therefore differences below 0.3 \log_{10} were not considered as significant. Applying this approach to the data in Tables 3 and 4 it appears that net counts at 37 C did not depend on the carbon source, but those at 30 C did.

The figures in Tables 3 and 4 also indicate that net recoveries at the lower temperature are considerably higher than at 37 C. This is further analyzed in Tables 5 and 6. The differences between confirmed counts at 30 and 37 C were greater in VRBG than in VRBL.

As indicated earlier, we had observed in preliminary studies that the rate of confirmation as *Enterobacteriaceae* of typical colonies in MacConkey-type agars increased with increasing temperature of incubation. The data obtained in this investigation were analyzed for this effect in Tables 7-10. Such a difference was not

TABLE 5. Influence of the incubation temperature on the recovery of *Enterobacteriaceae* from minced meat using VRBG-agar.

| Percentage of the samples | $\log_{10}\text{cfu}_{30\text{C}} - \log_{10}\text{cfu}_{37\text{C}}$ |
|---------------------------|---|
| 5 | 0.00 |
| 9 | 0.00-0.10 |
| 41 | 0.10-0.50 |
| 45 | > 0.50 |

Av. $\log_{10}\text{cfu}_{30\text{C net}} = 4.26$; av. $\log_{10}\text{cfu}_{37\text{C net}} = 3.61$.
Difference 0.65.

TABLE 6. Influence of the incubation temperature on the recovery of *Enterobacteriaceae* from minced meat using VRBL-agar.

| Percentage of the samples | $\log_{10}\text{cfu}_{30\text{C}} - \log_{10}\text{cfu}_{37\text{C}}$ |
|---------------------------|---|
| 15 | 0.00 |
| 23 | 0.00-0.10 |
| 36 | 0.10-0.50 |
| 26 | > 0.50 |

Av. $\log_{10}\text{cfu}_{30\text{C net}} = 3.92$; av. $\log_{10}\text{cfu}_{37\text{C net}} = 3.53$.
Difference 0.39.

TABLE 7. Confirmation rate of typical colonies obtained from minced meat using VRBG-agar incubated at 37 C.

| Percentage of the samples | $\log_{10}\text{cfu}_{\text{gross}} - \log_{10}\text{cfu}_{\text{net}}$ |
|---------------------------|---|
| 71 | 0.00 |
| 5 | 0.01-0.05 |
| 5 | 0.06-0.10 |
| 19 | 0.10 max. |

Av. $\log_{10}\text{cfu}_{\text{net}} = 3.61$; av. $\log_{10}\text{cfu}_{\text{gross}} = 3.76$.
Difference 0.15.

TABLE 8. Confirmation rate of typical colonies obtained from minced meat using VRBG-agar incubated at 30 C.

| Percentage of the samples | $\log_{10}\text{cfu}_{\text{gross}} - \log_{10}\text{cfu}_{\text{net}}$ |
|---------------------------|---|
| 75 | 0.00 |
| 6 | 0.01-0.05 |
| 6 | 0.06-0.10 |
| 13 | 0.10 max. |

Av. $\log_{10}\text{cfu}_{\text{net}} = 4.26$; av. $\log_{10}\text{cfu}_{\text{gross}} = 4.31$.

TABLE 9. Confirmation rate of typical colonies obtained from minced meat using VRBL-agar incubated at 37 C.

| Percentage of the samples | $\log_{10}\text{cfu}_{\text{gross}} - \log_{10}\text{cfu}_{\text{net}}$ |
|---------------------------|---|
| 86 | 0.00 |
| 2 | 0.01-0.05 |
| 7 | 0.06-0.10 |
| 5 | > 0.10 |

Av. $\log_{10}\text{cfu}_{\text{net}} = 3.53$; av. $\log_{10}\text{cfu}_{\text{gross}} = 3.55$.

TABLE 10. Confirmation rate of typical colonies obtained from minced meat using VRBL-agar incubated at 30 C.

| Percentage of the samples | $\log_{10}\text{cfu}_{\text{gross}} - \log_{10}\text{cfu}_{\text{net}}$ |
|---------------------------|---|
| 79 | 0.00 |
| 2 | 0.01-0.05 |
| 12 | 0.06-0.10 |
| 7 | > 0.10 |

Av. $\log_{10}\text{cfu}_{\text{net}} = 3.91$; av. $\log_{10}\text{cfu}_{\text{gross}} = 3.94$.

observed in any of the data, the confirmation rates of typical colonies being very high in all instances. This disagreement with the results of previous studies may be caused by two factors. First, the earlier investigations had been carried out with quite different MacConkey-type agar formulae; as we have observed that some of such media lead to poorly discernible colonies, our previous results may have suffered from this shortcoming of the media used. In addition, minced meat might contain less interfering types of organisms than the commodities studied earlier, which were mainly dried foods.

Lactose plus glucose versus glucose at 30 and 37 C

The results of the experiments with pure cultures recorded in Tables 1 and 2, point to the absence of any influence of the use of glucose alone or in combination with lactose on the productivity of MacConkey-type agars as expected. This required yet further substantiation by the use of the two types of media for examination of foods containing considerable numbers of *Enterobacteriaceae*. Minced meat and chicken drip were chosen as such substrates.

The data obtained with minced meat are summarized in Table 11. Once more, considering differences below $\log_{10} = 0.30$ as non-significant because within the confidence interval of the mode of examination, the results for glucose and glucose plus lactose were identical. Again, the differences between net and gross counts were negligible. And, once more the counts at 30 C were higher than those at 37 C.

TABLE 11. *The influence of carbon source and incubation temperature on the recovery of Enterobacteriaceae from minced meats.*

| Incubation temperature (C) | Gross (\log_{10} cfu/g) | | Confirmed (\log_{10} cfu/g) | |
|----------------------------|----------------------------|------|--------------------------------|------|
| | VRBLG | VRBG | VRBLG | VRBG |
| 30 | 4.46 | 4.42 | 4.37 | 4.40 |
| 37 | 4.14 | 4.11 | 4.14 | 3.90 |

The results of the examination of the samples of broiler drip are summarized in Table 12. In agreement with the observations made on minced meat, gross and confirmed counts did not differ, and the carbon source did not exert any influence. However, with broiler drip the temperature of incubation did not influence the colony counts. We had observed earlier that the types of *Enterobacteriaceae* isolated from minced meat were quite different from those present in chicken drip. Among the latter *Escherichia coli* seemed to predominate, whereas from minced meat mostly psychrotrophic types were isolated. To investigate this matter in more detail, 72 colonies taken at random from plates of minced meat, and 68 picked similarly from plates of chicken drip were identified. The following procedure was used for this purpose: (a) the three layer tubes referred to earlier, allowing assessment of oxidase reaction, mode of attack on glucose, motility and formation of H_2S and indole; (b) slants of Simmons citrate agar, DNA agar, urea agar and gelatin agar. The

TABLE 12. *The effects of carbon source and incubation temperature on the recovery of Enterobacteriaceae from broiler drip.*

| Incubation temperature (C) | Gross (\log_{10} cfu/g) | | Confirmed (\log_{10} cfu/g) | |
|----------------------------|----------------------------|------|--------------------------------|------|
| | VRBLG | VRBG | VRBLG | VRBG |
| 30 | 1.97 | 2.03 | 1.97 | 2.03 |
| 37 | 1.96 | 1.94 | 1.96 | 1.94 |

results obtained are presented in Table 13. Over 75% of the isolates obtained from minced meat were typical psychrotrophic types. These are mostly considered to be primarily of aqueous or vegetable origin. Psychrotrophic types occurred only to a total of 30% in chicken drip, with *E. coli* predominating there. It is therefore not surprising that *Enterobacteriaceae* counts in chicken were almost equal at 37 C and 30 C, but higher at the lower temperature when minced meat was examined.

TABLE 13. *Types of Enterobacteriaceae isolated from minced meat and drip from frozen broilers in violet red bile media at 30 C.*

| Species or genus ^a | Percentage of isolates from | |
|---|-----------------------------|--------------|
| | Minced meat | Broiler drip |
| <i>Serratia liquefaciens</i> ^b | 47 | 12 |
| <i>Escherichia coli</i> ^b | 9 | 35 |
| <i>Enterobacter</i> | 21 | 6 |
| <i>Erwinia</i> | 10 | 12 |
| <i>Hafnia</i> | 7 | 13 |
| <i>Citrobacter</i> | 4 | 13 |
| <i>Klebsiella</i> | < 1 | 6 |
| <i>Proteus</i> | < 1 | 3 |

^aThe order is that of decreasing summated frequency.

^bIncluding a minority of irregular types.

DISCUSSION

The investigations reported in this paper substantiate observations made repeatedly in the past.

First, we have demonstrated once more that indiscriminate use of dried formulae of MacConkey-type agars may lead to drastically reduced recoveries of *Enterobacteriaceae* and hence to serious underestimation of health or spoilage risks in foods. This confirms observations on similar and related media by numerous authors (3,13,22,23,26,34,43,53,54,55,57,58). These results call for constant monitoring of purchased dried culture media. Such testing should include (a) pure culture studies to assess productivity, and (b) practical challenge tests to estimate selectivity (31).

Furthermore, the possible influence of the incubation temperature on the results of testing for *Enterobacteriaceae* has been demonstrated once more. Incubation at ca. 30 C may lead to consistently higher results than at 37 C, as could be anticipated from physiological studies and ecological surveys carried out many years ago. The consequence of these observations for ecological studies are obvious. However, for practical monitoring the inference is less strict. Results of the bacteriological examination of foods depend strongly on the method by which they were obtained. Consequently Reference Values (35) carry little weight, unless accompanied by prescribed methods. Hence, any of the temperatures customarily used so far for enumeration of *Entero-*

bacteriaceae may be applied, provided Reference Values are adapted accordingly.

One new, although barely surprising, element emerged from our investigation. Using glucose or glucose plus lactose in MacConkey-type media does not in the least influence confirmed counts of *Enterobacteriaceae* at a given temperature. However, the composition of the media in other respects is of great significance. Particularly characteristics of the bile preparations may result in dramatic differences in recoveries (5,7,37,39). For practical purposes the conclusion is obvious: glucose-base media can be used in confidence, provided the bile preparation to be incorporated has been carefully selected by the manufacturer. Nevertheless it remains an obligation of the bacteriologist to check every new batch purchased from suppliers for productivity and selectivity, even when a manufacturer has previously been found to be offering satisfactory media.

With regard to the fundamental concept of replacing lactose by glucose, the data in Table 4 substantiate earlier findings (11) that this may lead to higher counts. However, this is not always the case, as the data in Table 3 show - for the simple reason that lactose-negative types of *Enterobacteriaceae* do not always occur in large numbers. Actually the types of *Enterobacteriaceae* colonizing proteinaceous staples may vary greatly, as the data in Table 13 and those of other research teams (9,47) clearly indicate. But in no instance will glucose media lead to lower results, as this study and other investigations (8,16) demonstrated beyond any doubt. Therefore, use of glucose-containing MacConkey-type media will result in data of considerably better consistency than those reported for similar media based on lactose (2,48).

REFERENCES

1. Arpai, J. 1962. Nonlethal freezing injury to metabolism and motility of *Pseudomonas fluorescens* and *Escherichia coli*. *Appl. Microbiol.* 10:297-301.
2. Baird-Parker, A. C. 1976. Microbiological standards for foods? *Die Fleischwirtschaft* 56:96.
3. Banwart, G. J., and J. C. Ayres. 1953. Effect of various enrichment broths and selective agars upon the growth of several species of *Salmonella*. *Appl. Microbiol.* 1:296-301.
4. Breer, C., and W. Ruosch. 1978. Der quantitative Nachweis von Enterobacteriaceen im Rahmen der bakteriologischen Betriebshygienekontrolle. *Die Fleischwirtschaft* 58:1837-1838.
5. Bridson, E. Y. 1979. Culture media: problems in the standardization of selective agents for the isolation of *Enterobacteriaceae*. *Antonie van Leeuwenhoek* 45:(in press).
6. Brown, A. D., and J. F. Weideman. 1958. The taxonomy of the psychrophilic meat spoilage bacteria: a reassessment. *J. Appl. Bacteriol.* 21:11-17.
7. Burman, N. P. 1955. The standardization and selection of bile salt and peptone for culture media used in the bacteriological examination of water. *Proc. Soc. Water Treatm. Exam.* 4:10-20.
8. Cox, N. A., A. J. Mercuri, B. J. Juven, and J. E. Thomson. 1975. *Enterobacteriaceae* at various stages of poultry chilling. *J. Food Sci.* 40:44-46.
9. Cox, N. A., and A. J. Mercuri. 1978. Comparison of two minikits (API and R-B) for identification of *Enterobacteriaceae* isolated from poultry and meat products. *J. Food Prot.* 41:107-110.
10. Dahlberg, A. C. 1946. The relationship of the growth of all bacteria and coliform bacteria in pasteurized milk held at refrigeration temperatures. *J. Dairy Sci.* 29:651-655.
11. Drion, E. F., and D. A. A. Mossel. 1977. The reliability of the examination of foods, processed for safety, for enteric pathogens and *Enterobacteriaceae*: a mathematical and ecological study. *J. Hygiene* 78:301-324.
12. Eddy, B. P., and A. G. Kitchell. 1959. Cold-tolerant fermentative Gram-negative organisms from meat and other sources. *J. Appl. Bacteriol.* 22:57-63.
13. Fagerberg, D. J., J. A. Avens, and B. A. George. 1976. Importance of brands of dehydrated culture medium. *J. Milk Food Technol.* 39:351-352.
14. Galesloot, Th. E. 1953. Some aspects of the bacteriology of pasteurized milk. V. The deterioration of commercially pasteurized milk (H.T.S.T. process). *Netherl. Milk Dairy J.* 7:15-40.
15. Greene, V. W., and J. J. Jezeski. 1954. Influence of temperature on the development of several psychrophilic bacteria of dairy origin. *Appl. Microbiol.* 2:110-117.
16. Hamann, R., and H. Weber. 1978. Ein Beitrag zum quantitativen und qualitativen Nachweis von *Enterobacteriaceae* in Speiseeis. *Arch. Lebensm. Hyg.* 29:5-16.
17. Hamilton, M. A., and G. K. Bissonnette. 1975. Statistical inferences about injury and persistence of environmentally stressed bacteria. *J. Hygiene* 74:149-155.
18. Hartsell, S. E. 1951. The longevity and behavior of pathogenic bacteria in frozen foods: the influence of plating media. *Amer. J. Public Health* 41:1072-1077.
19. Hechelmann, H., E. Rossmann, M. Peric, and L. Leistner. 1973. Untersuchung zur Ermittlung der *Enterobacteriaceae*-Zahl bei Schlachtgeflügel. *Die Fleischwirtschaft* 53:107-113.
20. Janssen, D. W., and F. F. Busta. 1973. Repair of injury in *Salmonella anatum* cells after freezing and thawing in milk. *Cryobiology* 10:386-392.
21. Kirsch, R. H., F. E. Berry, C. L. Baldwin, and E. M. Foster. 1952. The bacteriology of refrigerated ground beef. *Food Res.* 17:495-503.
22. Kroninger, D. L., and G. J. Banwart. 1978. Effect of various selective agars on the growth of rare and common salmonellae. *J. Food Sci.* 43:1328-1329.
23. McCormack, W. M., W. E. De Witt, P. E. Bailey, G. K. Morris, P. Soeharjono, and E. J. Gangarosa. 1974. Evaluation of thiosulfate-citrate-bile salts-sucrose agar, a selective medium for the isolation of *Vibrio cholerae* and other pathogenic vibrios. *J. Infect. Diseases* 129:497-500.
24. Marth, E. H., and W. C. Frazier. 1957. Bacteriology of milk held at farm bulk cooling tank temperatures. III. Psychrophiles found and their growth. *J. Milk Food Technol.* 20:93-99.
25. Mercuri, A. J., N. A. Cox, M. O. Carson, and D. A. Tanner. 1978. Relation of *Enterobacteriaceae* counts to *Salmonella* contamination of market broilers. *J. Food Prot.* 41:427-428.
26. Moats, W. A., and J. A. Kinner. 1974. Factors affecting selectivity of brilliant green-phenol red agar for salmonellae. *Appl. Microbiol.* 27:118-123.
27. Mossel, D. A. A. 1957. The presumptive enumeration of lactose negative as well as lactose positive *Enterobacteriaceae* in foods. *Appl. Microbiol.* 5:379-381.
28. Mossel, D. A. A., and H. Zwart. 1960. The rapid tentative recognition of psychrotrophic types among *Enterobacteriaceae* isolated from foods. *J. Appl. Bacteriol.* 23:185-188.
29. Mossel, D. A. A., W. H. J. Mengerink, and H. H. Scholts. 1962. A McConkey type agar medium for the selective growth and enumeration of all *Enterobacteriaceae*. *J. Bacteriol.* 84:381.
30. Mossel, D. A. A., and M. A. Ratto. 1970. Rapid detection of sublethally impaired cells of *Enterobacteriaceae* in dried foods. *Appl. Microbiol.* 20:273-275.
31. Mossel, D. A. A. 1971. Microbiological culture media as ecosystems. Ecometric evaluation of media. *Miscellaneous Papers, Agricult. University, Wageningen* 9:29-39.
32. Mossel, D. A. A., and G. A. Harrewijn. 1972. Les défaillances dans certains cas des milieux d'isolement des *Enterobacteriaceae* des aliments et des médicaments secs. *Alimenta* 11:29-30.

33. Mossel, D. A. A., B. Krol, and P. C. Moerman. 1972. Bacteriological and quality perspectives of *Salmonella* radication of frozen boneless meats. *Alimenta* 11:51-59.
34. Mossel, D. A. A., G. A. Harrewijn, and C. F. M. Nesselrooy - van Zadelhoff. 1974. Standardization of the selective inhibitory effect of surface active compounds used in media for the detection of *Enterobacteriaceae* in foods and water. *Health Lab. Sci.* 11:260-267.
35. Mossel, D. A. A. 1977. Microbiology of foods. Occurrence, prevention and monitoring of hazards and deterioration. Utrecht, The University. p. 81 *et seq.*
36. Mossel, D. A. A., I. Eelderink, and J. P. Sutherland. 1977. Development and use of single 'polytropic' diagnostic tubes for the approximate taxonomic grouping of bacteria isolated from foods, water and medicinal preparations. *Zentralbl. Bakteriologie. Parasitenk. Abt. I, Orig.*, A 238:66-79.
37. Mossel, D. A. A., A. J. W. M. van Ekeren, and I. Eelderink. 1977. A simplified procedure for the examination of drinking water for bacteria of public health significance: the differential hydrobacteriogramme. *Zentralbl. Bakteriologie. Parasitenk. Abt. I, Orig.*, B 165:498-516.
38. Mossel, D. A. A. 1978. Index and indicator organisms - a current assessment of their usefulness and significance. *Food Technol. Australia* 30:212-219.
39. Mossel, D. A. A., I. Eelderink, M. Koopmans, and F. van Rossem. 1978. Optimisation of a MacConkey type medium for the enumeration of *Enterobacteriaceae*. *Lab. Practice* 27:1049-1050.
40. Naemura, L. G., and R. J. Seidler. 1978. Significance of low-temperature growth associated with the fecal coliform response, indole production and pectin liquefaction in *Klebsiella*. *Appl. Environ. Microbiol.* 35:392-396.
41. Nagel, C. W., K. L. Simpson, R. H. Vaughn, and G. F. Stewart. 1960. Micro-organisms associated with spoilage of refrigerated poultry. *Food Technol.* 14:21-23.
42. Newton, K. G., and C. O. Gill. 1978. The development of the anaerobic spoilage flora of meat stored at chill temperatures. *J. Appl. Bacteriol.* 44:91-95.
43. Nicholls, K. M., J. V. Lee, and T. J. Donovan. 1976. An evaluation of commercial thiosulphate citrate bile salt sucrose agar (TCBS). *J. Appl. Bacteriol.* 41:265-269.
44. Nouws, J. F. M., and P. G. Heymans. 1975. A microbiological classification system for minced meat. *Arch. Lebensm. Hyg.* 26:175-180.
45. Panes, J. J., and S. B. Thomas. 1959. The multiplication of coliaerogenes bacteria in milk stored at 3-5°C. *J. Appl. Bacteriol.* 22:272-277.
46. Parry, A. H., and G. H. Tee. 1958. Investigation and control of ropiness in milk at a pasteurising plant. *Monthly Bull. Min. Health London* 17:19-23.
47. Patterson, J. T., and P. A. Gibbs. 1977. Incidence and spoilage potential of isolates from vacuum-packaged meat of high pH value. *J. Appl. Bacteriol.* 43:25-38.
48. Pierson, C. J., B. S. Emswiler, and A. W. Kotula. 1978. Comparison of methods for estimation of coliforms, fecal coliforms and enterococci in retail ground beef. *J. Food Prot.* 41:263-266.
49. Raccach, M., and B. Juven. 1973. Studies of the recovery of frozen *Salmonella gallinarum*: the roles of cooling rate, initial cell concentrations and plating medium. *J. Food Technol.* 8:211-216.
50. Raccach, M., and B. J. Juven. 1976. Effect of suspending and plating media on the recovery of *Salmonella gallinarum* following freezing and thawing. *J. Food Technol.* 11:221-227.
51. Ray, B., and M. L. Speck. 1972. Repair of injury induced by freezing *Escherichia coli* as influenced by recovery medium. *Appl. Microbiol.* 24:258-263.
52. Ray, B., and M. L. Speck. 1972. Metabolic process during the repair of freeze-injury in *Escherichia coli*. *Appl. Microbiol.* 24:585-590.
53. Read, R. B., and A. L. Reyes. 1968. Variation in plating efficiency of *Salmonellae* on eight lots of brilliant green agar. *Appl. Microbiol.* 16:746-748.
54. Reusze, U. 1971. Erfahrungen über die Wirksamkeit von Fertignährböden bei der Salmonellen-Isolierung. *Arch. Lebensm. Hyg.* 22:11-15.
55. Reynolds, N., and P. C. Wood. 1956. Improved techniques for the bacteriological examination of molluscan shell-fish. *J. Appl. Bacteriol.* 19:20-25.
56. Schultze, W. D., and J. C. Olson, Jr. 1960. Studies on psychrophilic bacteria. I. Distribution in stored commercial dairy products. *J. Dairy Sci.* 43:346-350.
57. Taylor, W. I., and D. Schelhart. 1968. Isolation of *Shigellae*. VI. Performance of media with stool specimens. *Appl. Microbiol.* 16:1387-1393.
58. Taylor, W. I., and D. Schelhart. 1971. Isolation of *Shigellae*. VIII. Comparison of xylose lysine deoxycholate agar, Hektoen enteric agar, Salmonella-Shigella-agar and eosin methylene blue agar with stool specimens. *Appl. Microbiol.* 21:32-37.
59. Warseck, M., B. Ray, and M. L. Speck. 1973. Repair and enumeration of injured coliforms in frozen foods. *Appl. Microbiol.* 26:919-924.

Northolt, Van Egmond and Paulsch, *con't from p. 490*

29. Scott, P. M., W. van Walbeek, B. Kennedy, and D. Anyeti. 1972. Mycotoxins (ochratoxin A, citrinin, and sterigmatocystin) and toxigenic fungi in grains and other agricultural products. *J. Agric. Food Chem.* 20:1103-1109.
30. Scott, W. J. 1957. Water relations of food spoilage micro-organisms. *Adv. Food Res.* 7:83-127.
31. Shotwell, O. L., C. W. Hesseltine, E. E. Vandegraft, and M. L. Goulden. 1971. Survey of corn from different regions for aflatoxins, ochratoxin, and zearalenone. *Cereal Sci. Today* 16:226-273.
32. Shotwell, O. L., M. L. Goulden, and C. W. Hesseltine. 1976. Survey of U.S. wheat for ochratoxin and aflatoxin. *J. Assoc. Off. Anal. Chem.* 59:122-124.
33. Steyn, P. S. 1971. Ochratoxin and other dihydroisocoumarins. p. 179. In A. Ciegler, S. Kadis, and S. J. Ajl (eds.). *Microbiol toxins*. Vol. 6, Academic Press, New York and London.
34. Stoloff, L. 1976. Occurrence of mycotoxins in foods. p. 40. In J. V. Rodricks (ed.). *Mycotoxins and other related food problems*. Am. Chem. Soc., Washington, D.C.

Wood, *con't from p. 501*

- 14:1103.
65. VanEtten, H. D., and D. A. Smith. 1975. Accumulation of antifungal isoflavonoids and 1a-hydroxyphaseollone, a phaseollin metabolite, in bean tissue infected with *Fusarium solani* f. sp. *Phaseoli*. *Physiol. Plant Pathol.* 5:225.
66. Williams, M. C. 1970. Xanthotoxin and bergapten in spring parsley. *Weed Sci.* 18:479.
67. Wood, G. E. 1976. Stress metabolites of white potatoes. p. 369. In: J. V. Rodricks (ed.) *Mycotoxins and other fungal related food problems*. American Chemical Society, Washington, D.C.
68. Wyman, J. G., and H. D. VanEtten. 1978. Antibacterial activity of selected isoflavonoids. *Phytopathology* 68:583.