

Research Note

Content of Toxic and Essential Metals in Canned Mussels Commonly Consumed in Tenerife, Canary Islands, Spain

ANGEL GUTIÉRREZ,¹ GONZALO LOZANO,^{1*} ARTURO HARDISSON,² CARMEN RUBIO,² AND TOMÁS GONZÁLEZ³

¹Department of Animal Biology (Marine Sciences), University of La Laguna, Tenerife, Canary Islands, Spain; ²Department of Public Health, University of La Laguna, Tenerife, Canary Islands, Spain; and ³Canarian Public Health Service, Central Laboratory, Santa Cruz de Tenerife, Canary Islands, Spain

MS 03-376: Received 21 August 2003/Accepted 17 January 2004

ABSTRACT

Concentrations of three toxic heavy metals (Hg, Pb, Cd) and six essential heavy metals (Fe, Zn, Mn, Cr, Cu, Ni) were determined in mussel conserves (*Mytilus galloprovincialis*, *Bivalvia*, *Mollusca*) consumed habitually by individuals in Tenerife (Canary Islands, Spain). A total of 600 samples were analyzed, corresponding to six different commercial brands and four different processing types: pickled sauce (mixture of olive oil, vinegar, red pepper, laurel, and salt), coquille St. Jacques sauce (coquille St. Jacques broth), nature (water and salt), and bionature (water, salt, and soluble vegetal fiber). Samples were collected weekly from markets in Santa Cruz de Tenerife during a 12-month period. All values for toxic metals were lower than the permitted maximum for human consumption as proscribed in European Community Directive 2001/22/CE (1,000 µg/kg wet weight for Pb and Cd) and European Community Decision 93/351/EEC (500 µg/kg wet weight for Hg). For the six essential heavy metals, mussels are a very good source, contributing high percentages of the recommended daily allowance.

The tendency of mussels to accumulate heavy metals is related to their weight (8) and can reflect the local concentration of various metals and other toxins (4). Mussels are often used as bioindicators because they have the capacity to concentrate contaminants from seawater by several orders of magnitude.

However, these molluscs are an important component of aquaculture programs (Spain produces nearly 300,000 tons yearly) and make up part of the Spanish traditional diet. They can be used in their natural state or they can be commercially processed in many ways, the most popular of which is canning. Because of their popularity, mussels are a nutritionally important component of the Spanish diet.

In the Canary Islands (central-east Atlantic Ocean, 27 to 30°N and 13 to 19°W), there are no natural or cultured populations of mussels of the genus *Mytilus* (e.g., *Mytilus edulis* and *Mytilus galloprovincialis*), which are typical to the Iberian Peninsula. However, because demand for these molluscs is high they are imported and are one of the most common marine foods in the island diet. The material chosen for study was canned mussels of Galician origin (north-western Spain), which are always available in Santa Cruz de Tenerife supply markets.

Heavy metals can accumulate in aquatic organisms from exposure to toxic substances in water and sediments or in the food chain (38). Because consumption of contaminated foodstuffs is the main route of exposure to trace met-

als for most humans, information about dietary intake is important to assess risks to human health (21). In this study, we analyzed concentrations of toxic metals (Hg, Pb, and Cd) and essential metals (Fe, Zn, Mn, Cr, Cu, and Ni), which can become toxic at high concentrations.

Mercury is well known for its neurotoxicity, but it also affects the gastrointestinal tract and renal function (31). In aquatic environments, inorganic Hg is microbiologically transformed into the lipophilic organic compound methylmercury. This transformation makes Hg more prone to biomagnification in aquatic food chains (17). The presence of metals in marine sediments is a consequence of both natural and anthropogenic contamination, but the presence of toxic metals in some species of fish and their products indicates extreme contamination of the aquatic environment by those metals (29). Consequently, human populations with a traditionally high dietary intake of food originating from either freshwater or marine environments have the highest potential for exposure to methylmercury (17).

Cadmium in the human diet constitutes a potential chronic health hazard. In the nonsmoking general population, diet is the major source of Cd exposure. In 1968, the Japanese Ministry of Health and Welfare admitted that "Itai-itai" disease was caused primarily by Cd. The WHO International Programme on Chemical Safety established that important sources of Cd contamination include phosphate fertilizer, sewage sludge, plated or galvanized equipment, enamels, and glazes. Although molluscs and crustaceans accumulate Cd, fishes generally have Cd levels of <0.2 µg/kg (15).

* Author for correspondence. Tel: 34-22-318385; Fax: 34-22-318311; E-mail: glozano@ull.es.

TABLE 1. Reference materials for recuperation study

Metal	Reference material	Mean concentrations ($\mu\text{g}/\text{kg}$ wet weight) ^a	Certified values ($\mu\text{g}/\text{kg}$ wet weight)	Method ^b
Pb	NBS SRH 1577a bovine liver	135 \pm 15	135 \pm 15	ET-AAS
Pb	BCR-278 R mussel tissue	2,000 \pm 40	1,970 \pm 70	ET-AAS
Cd	BCR-278 R mussel tissue	348 \pm 7	347 \pm 8	FT-AAS
Cd	BCR CRM 184 bovine muscle	12 \pm 1	13 \pm 2	ET-AAS
Hg	BCR-278 R mussel tissue	196 \pm 9	195 \pm 10	CV-AAS
Cr	BCR-278 R mussel tissue	780 \pm 60	760 \pm 40	ICP-AES
Cu	BCR-278 R mussel tissue	9,450 \pm 130	9,390 \pm 100	ICP-AES
Cu	BCR-442 cod muscle	1,050 \pm 70	1,050 \pm 80	ICP-AES
Fe	BCR-278 R mussel tissue	5,460 \pm 300	5,460 \pm 300	ICP-AES
Mn	BCR-278 R mussel tissue	7,690 \pm 230	7,690 \pm 220	ICP-AES
Zn	BCR-278 R mussel tissue	8,310 \pm 1,700	8,280 \pm 1,000	ICP-AES
Ni	BCR-414 plankton	18,800 \pm 800	18,800 \pm 600	ICP-AES

^a $n = 11$.

^b ET-AAS, electrothermal atomic absorption spectrophotometry; CV-AAS, cold vapor atomic absorption spectrophotometry; ICP-AES, inductively coupled plasma atomic emission spectrophotometry.

Lead is a nonessential heavy metal that can occur in high concentrations in animals, especially mammals (32). High concentrations of Pb have been linked to human health problems, including nervous system dysfunction in fetuses and infants and hematotoxicity, reproductive dysfunction, and Alzheimer's disease in adults. Even low Pb concentrations can be associated with asthma, anemia, and learning disabilities (16, 20).

Iron and zinc are essential minerals for humans. Nevertheless, high Fe concentrations can produce toxic cardiovascular, neurological, and gastrointestinal effects (3). Zinc concentrations in bivalve molluscs usually exceed the 50% lethal dose during embryogenesis (33). Manganese is the least toxic of the metals analyzed in this study, but it can be toxic to the nervous system, resulting in Parkinsonian signs (32). Chromium compounds are often environmental pollutants and are known for their mutagenic, carcinogenic,

and toxic effects (28). Copper concentrations of 63.6 $\mu\text{g}/\text{liter}$ reduce the growth of bivalve molluscs and can be fatal at $>86.5 \mu\text{g}/\text{liter}$ (19). Aquatic Ni concentrations are usually approximately 6 $\mu\text{g}/\text{liter}$, but higher concentrations have been reported (14).

The main goal of this research was to assess heavy metal contamination in canned mussels normally consumed in Tenerife and to use these data to evaluate nutritional patterns and toxicological risk.

MATERIALS AND METHODS

Samples for study (five cans per week) were collected over the course of 12 months from various stores in Santa Cruz de Tenerife. A total of 600 samples were analyzed, corresponding to six different commercial brands (A through F) and four different processing types: pickled sauce (mixture of olive oil, vinegar, red pepper, laurel, and salt), coquille St. Jacques sauce (coquille St.

TABLE 2. General composition of mussels by brand

Brand	n	Dry weight (g)	% moisture	Ash weight (g)
A	150	3.25 \pm 0.04	67.51 \pm 0.34	0.17 \pm 0.01
B	90	3.30 \pm 0.001	67.03 \pm 0.15	0.19 \pm 0.01
C	51	3.21 \pm 0.04	67.92 \pm 0.27	0.19 \pm 0.02
D	129	3.16 \pm 0.07	67.95 \pm 0.17	0.22 \pm 0.02
E	93	3.23 \pm 0.04	67.68 \pm 0.24	0.18 \pm 0.02
F	57	3.35 \pm 0.03	66.53 \pm 0.22	0.21 \pm 0.02

TABLE 3. General composition of mussels by processing type

Process	n	Dry weight (g)	% moisture	Ash weight (g)
Coquille St. Jacques sauce	213	3.28 ± 0.04	66.8 ± 0.40	0.25 ± 0.01
Pickled sauce	240	3.30 ± 0.02	67.02 ± 0.15	0.18 ± 0.007
Nature	75	2.99 ± 0.08	69.80 ± 0.40	0.16 ± 0.02
Bionature	41	3.14 ± 0.05	68.62 ± 0.47	0.22 ± 0.03

TABLE 4. Heavy metal concentration (µg/kg wet weight) with respect to brand and processing type

Metal	Brand	Coquille St. Jacques sauce	Pickled sauce	Nature	Bionature	Mean ± SD
Pb	A	4.47 ± 0.68	7.75 ± 0.68	7.73 ± 0.82	8.16 ± 1.10	6.96 ± 0.48
	B	7.12 ± 0.86	7.87 ± 0.75			7.57 ± 0.25
	C		7.94 ± 1.09			7.94 ± 1.09
	D	5.28 ± 0.84	5.90 ± 0.92	6.38 ± 0.36		
	E	6.21 ± 0.80	7.96 ± 0.65	7.17 ± 0.20		
	F		8.09 ± 0.85	8.09 ± 0.85		
Cd	A	5.13 ± 0.46	4.49 ± 0.18	4.52 ± 0.32	4.35 ± 0.24	4.65 ± 0.08
	B	4.74 ± 0.35	4.94 ± 0.44			4.86 ± 0.11
	C		5.71 ± 0.87			5.71 ± 0.87
	D	6.00 ± 0.59	5.55 ± 0.29	5.01 ± 0.09		
	E	4.33 ± 0.36	3.98 ± 0.15	4.14 ± 0.10		
	F		4.60 ± 0.34	4.60 ± 0.34		
Cu	A	1,380 ± 70	1,870 ± 110	1,880 ± 110	1,580 ± 100	1,760 ± 70
	B	1,170 ± 120	1,560 ± 90			1,400 ± 80
	C		1,580 ± 130			1,580 ± 130
	D	1,120 ± 110	1,790 ± 190	1,460 ± 80		
	E	1,600 ± 150	1,790 ± 120	1,710 ± 90		
	F		1,550 ± 90	1,550 ± 90		
Ni	A	5,110 ± 270	5,570 ± 250	4,920 ± 400	8,870 ± 350	5,220 ± 190
	B	7,920 ± 100	6,360 ± 710			6,980 ± 250
	C		7,760 ± 260			7,760 ± 260
	D	8,300 ± 850	6,780 ± 290	8,030 ± 350		
	E	4,580 ± 270	4,950 ± 290	4,780 ± 200		
	F		5,830 ± 250	5,830 ± 250		
Fe	A	18,310 ± 2,920	42,180 ± 5,960	33,270 ± 2,240	35,520 ± 3,510	33,240 ± 3,010
	B	32,350 ± 3,070	26,650 ± 1,700			28,330 ± 1,730
	C		44,620 ± 4,700			44,620 ± 4,700
	D	28,260 ± 3,140	31,870 ± 3,800	32,070 ± 4,220		
	E	33,320 ± 5,130	42,550 ± 3,990	38,380 ± 3,310		
	F		26,220 ± 2,350	26,220 ± 2,350		
Zn	A	51,530 ± 2,530	62,370 ± 3,430	74,620 ± 4,400	58,890 ± 2,730	64,180 ± 2,510
	B	63,040 ± 2,630	56,680 ± 2,950			59,230 ± 2,150
	C		69,830 ± 2,810			69,830 ± 2,810
	D	60,880 ± 3,080	59,430 ± 3,600	71,060 ± 7,230		
	E	60,550 ± 3,620	53,770 ± 2,040	56,680 ± 2,080		
	F		61,190 ± 2,870	61,190 ± 2,870		
Cr	A	1,420 ± 100	1,480 ± 70	1,290 ± 80	1,750 ± 90	1,400 ± 50
	B	1,770 ± 80	1,740 ± 80			1,750 ± 40
	C		2,120 ± 100			2,120 ± 100
	D	1,760 ± 100	1,800 ± 100	1,390 ± 140		
	E	1,250 ± 90	1,270 ± 60	1,260 ± 50		
	F		1,690 ± 100	1,690 ± 100		
Mn	A	1,610 ± 180	1,740 ± 70	1,350 ± 110	1,800 ± 150	1,570 ± 80
	B	2,040 ± 180	1,370 ± 70			1,630 ± 110
	C		1,800 ± 190			1,800 ± 190
	D	1,820 ± 180	1,490 ± 170	1,510 ± 160		
	E	1,950 ± 160	1,640 ± 100	1,780 ± 100		
	F		1,430 ± 470	1,430 ± 470		

TABLE 5. Daily consumption of mussels necessary to exceed the admissible daily intake of Pb, with respect to brands and processing types

Brand	Amount (g wet weight) by processing type			
	Coquille St. Jacques sauce	Pickled sauce	Nature	Bionature
A	194.933	222.717	221.238	
B	210.970	202.429		
C		175.131		
D	166.666	180.180	205.338	229.885
E	230.946	251.256		
F		217.391		

Jacques broth), nature (water and salt), and bionature (water, salt, and soluble vegetal fiber). Whatever processed products were available at the time were collected. The individual mussels in each can (all the same size and coming from the same place) were homogenized to obtain the required amount of total sample (≈31.5 g) for analysis. Taking into account the peculiarities of Hg compared with the other metals analyzed, two different sample preparation protocols were followed. For Hg analysis, three 0.5-g subsamples (each weighed in Teflon reactors) were taken from the homogenized portions, 10 ml of concentrated HNO₃ was added, and the mixture was heated in a microwave oven long enough for total digestion to take place (≈20 min). The mixture was allowed to cool to room temperature, and the volume was brought to 50 ml by the addition of double-distilled deionized water.

For the remaining metals, three 10-g subsamples were weighed into porcelain crucibles and heated on a hot plate to 80 to 105°C until a constant dry weight was obtained. The crucibles were then placed in an oven furnace and the temperature was raised gradually (≈50°C/h) until 450 ± 25°C was reached. Approximately 18 to 24 h elapsed from the beginning of incineration until the formation of white ash. Crucibles were then left to cool to room temperature, and ash was dissolved with 1 ml of 0.6% HNO₃. Sample volume was brought to 50 ml by the addition of double-distilled deionized water.

For toxic heavy metal analysis, an atomic absorption spectrophotometer (4100 ZL Zeeman, Perkin Elmer, Foster City, Calif.) was used with a graphite 4100 ZL oven (Perkin Elmer) and an automatic AS 70 sampler for Pb and Cd (29) and with a hydrures generator, FIAS 400 injection system, and AS 90 automatic sampler (Perkin Elmer) for Hg (19, 28). For essential heavy metal analysis, an atomic emission spectrophotometer (Atomscan 25, Thermo Jarell Ash, Ventura, Calif.) of inductively coupled plasma was used. Table 1 lists the values for the recuperation study carried out using the corresponding reference materials to demonstrate the performance of the method.

Results were analyzed for normality with the Kolmogorov-Smirnov model (36) and for homogeneity of variance with a Levene test (27). When data were not normally distributed, the non-parametric Kruskal-Wallis test and Mann-Whitney U test (11) were used. Normally distributed data were analyzed using an analysis of variance and post hoc tests (24).

RESULTS AND DISCUSSION

Results regarding the general composition of the mussels are presented in Tables 2 and 3, and results from the heavy metal analyses are presented in Table 4. Dry weight, moisture percentage, and ash weight obtained were not sig-

TABLE 6. Daily consumption of mussels necessary to exceed the admissible daily intake of Cd, with respect to brands and processing types

Brand	Amount (g wet weight) by processing type			
	Coquille St. Jacques sauce	Pickled sauce	Nature	Bionature
A	212.765	129.032	129.366	
B	140.449	127.064		
C		125.944		
D	189.393	169.491	179.211	122.549
E	161.030	125.628		
F		123.609		

nificantly different among the different brands and processing types, mainly because the different brands market the same type of mussel in cans. The moisture percentage in the samples was less than that obtained by Moreiras et al. (22). In their study, 85.4% moisture was obtained, which is 20% higher than that obtained in the present study, but this higher moisture content could be due to the comparison between fresh and canned food; despite being immersed in liquid, canned mussels have been boiled and thus have lost part of their water content.

The Pb concentration found in samples was always much less than the maximum allowed by European Community Directive 2001/22/CE (13) (1,000 µg/kg wet weight; all concentrations reported here are on a wet weight basis unless otherwise indicated). The highest Pb concentrations were found in brand F mussels (8.09 ± 0.85 µg/kg) and in mussels processed with pickled sauce (8.16 ± 1.10 µg/kg) (Fig. 1). Other authors have found higher concentrations of Pb in other bivalves; for oysters, 1,250 to 14,000 µg/kg was found in areas of urban contamination (2) and 440 to 640 µg/kg (7) was found in zones considered normal.

The Cd concentration was also always much less than that permitted by Directive 2001/22/CE (1,000 µg/kg wet weight). The highest concentrations were found in brand C (5.71 ± 0.87 µg/kg) and in mussels processed with coquille St. Jacques sauce (6.0 ± 0.59 µg/kg) (Fig. 2). Several authors have studied Cd content in molluscs, obtaining concentrations in *Mytilus* spp. of 1,780 µg/kg in the Looe estuary (Cornwall, UK) and concentrations in *M. galloprovincialis* of 4,000 ± 1,500 µg/kg in Cape Ghir and 800 ±

TABLE 7. Nutritional importance of 150 g of canned mussels (approximately one can) in reference to heavy metals with recommended daily allowance (RDA)

Metal	Metal concentration		
	RDA (mg/day)	(mg/150 g mussels)	% RDA
Mn	2–5.0	0.27	13.5
Fe	15	6.60	44
Cu	1.3–3	0.24	18.46
Zn	15 or 12	10.47	15




FIGURE 1. Pb concentration in canned mussels ($\mu\text{g}/\text{kg}$ wet weight) with respect to brand (A) and processing type (B).

300 $\mu\text{g}/\text{kg}$ in Anza (Saharian coast) (18, 23). For *Venus verrucosa*, $3,930 \pm 2,840$ $\mu\text{g}/\text{kg}$ was obtained in Cansado Bay and $11,860 \pm 4,820$ $\mu\text{g}/\text{kg}$ was obtained in Levrier Bay (30). These concentrations are higher than those found by us because these researchers carried out their analyses in areas susceptible to patent contamination.

All sample concentrations of Hg were below the limit of detection for the analysis equipment, which was 5 $\mu\text{g}/\text{kg}$. Because of the small amount of sample processed (0.5 g maximum), the signal was never higher than background, so we could not determine the exact concentration of this metal in these samples. This limit of detection is much lower than the maximum concentration set by European Community Decision 93/351/EEC (12), which is 500 $\mu\text{g}/\text{kg}$ wet weight. Other authors have obtained Hg concentrations ranging from 10 to 210 $\mu\text{g}/\text{kg}$ (mean, 60 $\mu\text{g}/\text{kg}$) in fish, including prawns, and from 10 to 130 $\mu\text{g}/\text{kg}$ in shellfish, with means of 40 $\mu\text{g}/\text{kg}$ (25, 26) and 54 $\mu\text{g}/\text{kg}$ (37).

Brand C mussels contained the highest concentrations of Cr ($2,120 \pm 100$ $\mu\text{g}/\text{kg}$), Fe ($44,620 \pm 4,700$ $\mu\text{g}/\text{kg}$), Mn ($1,800 \pm 190$ $\mu\text{g}/\text{kg}$), and Zn ($69,830 \pm 2,810$ $\mu\text{g}/\text{kg}$) (Fig. 3). Zinc concentrations between 38,000 and 61,700




FIGURE 2. Cd concentration in canned mussels ($\mu\text{g}/\text{kg}$ wet weight) with respect to brand (A) and processing type (B).

$\mu\text{g}/\text{kg}$ have been reported in mussels and oysters (18) and concentrations between 35,100 and 69,400 $\mu\text{g}/\text{kg}$ have been found in *Homarus americanus* digestive glands (10).

Brand C mussels were small, with a mean size between 18 and 24 mm, whereas in cans of the other brands the mean sizes were larger, between 24.5 and 42.4 mm. This finding indicates that the assimilation of essential metals is higher in less mature organisms, as has been hypothesized by others (6, 34). Brand D mussels had the highest concentrations of Ni ($8,030 \pm 350$ $\mu\text{g}/\text{kg}$). Others have reported Ni concentrations ranging from 200 $\mu\text{g}/\text{kg}$ (2) to $7,080 \pm 1,570$ $\mu\text{g}/\text{kg}$ (1), in general agreement with the results of the present study.

Brand A mussels had the highest concentrations of Cu ($1,760 \pm 70$ $\mu\text{g}/\text{kg}$). Cu concentrations are highly variable in bivalve molluscs, with reports of 3,900 to 49,800 $\mu\text{g}/\text{kg}$ in mussels and oysters (18) and 5 and 20 $\mu\text{g}/\text{kg}$ for *M. galloprovincialis* (5). Data collected from brand C mussels confirm that Pb, Cu, Zn, and Fe concentrations in *Mytilus* decrease with increasing mussel weight, whereas Ni and Cd concentrations remain constant (8, 9).

Tables 5 and 6 present the daily consumption of mussels (expressed in grams) necessary to exceed the admis-




FIGURE 3. Concentrations of essential heavy metals Zn and Fe (A) and Cr, Mn, Ni, and Cu (B) with respect to brand of canned mussels.

sible daily allowance of Pb and Cd in humans. Because the amounts of mussels normally consumed are much lower than these thresholds, the mussel brands and preparations studied can be considered safe to consume.

We were unable to accurately assess Hg concentrations with the procedures used in this study because all concentrations were below the detection limits of the equipment. Thus, we concluded that Hg concentrations in these products are minimal with respect to the maximum legal limit for human consumption.

The nutritional significance of mussels, based on consumption of 150 g of canned mussels (approximately one can) for an adult, for the essential heavy metals that have recommended daily allowances (Mn, Fe, Cu, and Zn) is presented in Table 7. These data confirm the suitability of this food for human consumption. Because of low concentrations of toxic metals and high concentrations of essential metals, mussels have an important place among healthy high-quality foods according to recommendations of the World Health Organization published in 1988 (35).

REFERENCES

- Al-Madfa, H., M. A. R. Abdel-Moati, and F. H. Al-Gimaly. 1998. *Pinctata radiata* (pearl oyster): a bioindicator for metal pollution monitoring in the Quatari waters (Arabian Gulf). *Bull. Environ. Contam. Toxicol.* 60:245–251.
- Al-Sayed, H. A., A. M. Mahasaneh, and J. Al-Saad. 1994. Variations of trace metals concentrations in seawater and pearl oyster *Pinctata radiata* from Bahrain (Arabian Gulf). *Mar. Pollut. Bull.* 28:370–374.
- Baruthio, F. 1991. Toxicologie des éléments trace essentiels, p. 214–310. In *Les Oligoéléments en Médecine et Biologie*. Lavoisier-Tec Doc, Paris.
- Bayne, B. L. 1989. The biological effects of marine pollutants, p. 131–152. In J. Albaiges (ed.), *Marine pollution*. Hemisphere Publishing, New York.
- Bigas, M., E. Sagristá, G. Bozzo, M. Dufort, and M. Poquet. 2000. Occurrence of heavy metals and protozoan parasites in the mussel, *Mytilus galloprovincialis*, collected in the western Mediterranean. *Ecotoxicol. Environ. Restor.* 3:16–21.
- Blueweis, L., H. Fox, V. Kudzma, D. Nakashima, R. Peters, and S. Sams. 1978. Relationships between body size and sometime life history parameters. *Oecologia* 37:257–272.
- Bou-Olyan, A. H., S. Al-Mater, S. Al-Yakoob, and S. Al-Hazeens. 1995. Accumulation of lead, cadmium, copper and nickel by pearl oyster *Pinctata radiata* from Kuwait marine environment. *Mar. Pollut. Bull.* 30:211–214.
- Boyden, C. R. 1974. Trace elements and body size in molluscs. *Nature* 251:311–314.
- Bryan, G. W. 1976. Some aspects of heavy metals tolerance in aquatic organisms, p. 7–34. In P. M. Locwood (ed.), *Effects of pollutants on aquatic organisms*. Society of Experimental Biology Seminar Series 2. Cambridge University Press, New York.
- Chou, C. L., L. A. Paon, J. D. Moffatt, and B. Zwicker. 2000. Copper contamination and cadmium, silver and zinc concentrations in the digestive glands of American lobster (*Homarus americanus*) from the inner Bay of Fundy, Atlantic, Canada. *Bull. Environ. Contam. Toxicol.* 65:470–477.
- Choy, E. H. S., D. L. Scott, G. H. Kingsley, S. Thomas, A. G. U. Murphy, N. Staimos, and G. S. Panayi. 2001. Control of rheumatoid arthritis by oral tolerance. *Arthritis Rheum.* 44:1993–1997.
- European Community. 1993. 93/351/EEC: Commission decision of 19 May 1993 determining analysis methods, sampling plans and maximum limits for mercury in fishery products. Available at: <http://www.useu.be/agri/seafood2.html>.
- European Community. 2001. Directive 2001/22/CE. Available at: http://admi.net/eur/loi/leg_euro/fr_301L0022.html.
- Food and Agriculture Organization. 1972. *Marine pollution and sea life*, p. 343–345, 584–593. Food and Agriculture Organization, Rome.
- González-Soto, E., V. González-Rodríguez, C. López-Suárez, J. M. Castro-Romero, J. Pérez-Iglesias, and J. M. Fernández-Solís. 2000. Migration of lead and cadmium from ceramic materials used in food preparation. *Bull. Environ. Contam. Toxicol.* 65:598–603.
- Halliwell, D., N. Turoczy, and F. Stagnitti. 2000. Lead concentrations in *Eucalyptus* sp. in a small coastal town. *Bull. Environ. Contam. Toxicol.* 65:583–590.
- Hansen, J. C., and G. Danscher. 1997. Organic mercury: an environmental threat to the health of dietary-exposed societies. *Rev. Environ. Health* 12:107–116.
- Hardisson, A. 1984. Niveles de concentración de metales pesados (Hg, Zn, Cd, Pb, Cu, Fe) en especies marinas de interés bromatológico. Ph.D. thesis. University of La Laguna, Tenerife, Canary Islands, Spain.
- His, E., R. Beiras, and N. W. L. Seaman. 2000. The assessment of marine pollution—bioassays with bivalve embryos and larvae. *Mar. Biol.* 37:3–139.
- Kurasaki, M., D. I. Hartoto, T. Saito, M. Suzuki-Kurasaki, and T. Iwakuma. 2000. Metals in water in the Central Kalimantan, Indonesia. *Bull. Environ. Contam. Toxicol.* 65:591–597.
- Llobet, J. M., S. Granero, M. Schuhmacher, J. Corbella, and J. L. Domingo. 1998. Biological monitoring of environmental pollution

- and human exposure to metals in Tarragona, Spain. IV. Estimation of the dietary intake. *Trace Elem. Electrolytes* 15:136–141.
22. Moreiras, O., A. Carbajal, L. Cabrera, and C. Cuadrado. 1999. Tablas de composición de alimentos. Pirámide, Madrid.
 23. Moukrim, A., A. Kaaya, S. Najimi, M. Romeo, M. Gnassia-Barelli, and J. F. Narbone. 2000. Assessment of the trace metal levels in two species of mussels from the Agadir Marine Bay, south of Morocco. *Bull. Environ. Contam. Toxicol.* 65:478–485.
 24. Mycroft, R. H., D. C. Mitchell, and J. Kay. 2000. An evaluation of statistical procedures for comparing an individual's performance with that of a group of controls. *Cognit. Neuropsychol.* 19:291–299.
 25. Nixon, E., A. Rowe, and D. McLaughlin. 1994. Mercury concentrations in fish from Irish waters in 1993. Marine Environmental Series 2/94. Fisheries Research Centre, Abbotstown, Dublin.
 26. Nixon, E., A. Rowe, and D. McLaughlin. 1995. Mercury concentrations in fish from Irish waters in 1994. Marine Environmental Series 2/95. Fisheries Research Centre, Abbotstown, Dublin.
 27. Pan, G. 2002. Confidence intervals for comparing two scale parameters based on Levene's statistics. *J. Nonparametr. Stat.* 4:459–476.
 28. Parlak, H., B. Katalay, and B. Büyükişik. 1999. Accumulation and loss of chromium by mussels (*Mytilus galloprovincialis*). *Bull. Environ. Contam. Toxicol.* 62:286–292.
 29. Pokorny, B., and C. Ribaric-Lasnik. 2000. Lead, cadmium and zinc in tissues of roe deer (*Capreolus capreolus*) near the lead smelter in the Koroska Region (northern Slovenia). *Bull. Environ. Contam. Toxicol.* 64:20–26.
 30. Roméo, M., Z. Sidomou, and M. Gnassia-Barelli. 2000. Heavy metals in various molluscs from the Mauritanian coast. *Bull. Environ. Contam. Toxicol.* 65:269–276.
 31. Sierra, A., and A. Hardisson. 1991. La contaminación química de los alimentos. Aditivos alimentarios, p. 293–303. In G. Piédrola, J. del Rey, M. Domínguez, P. Cortina, R. Gálvez, A. Sierra, M. C. Sáenz, L. I. Gómez, J. Fernández-Crehuet, L. Salleras, A. Cueto, and J. J. Gestal (ed.), *Medicina preventiva y salud pública*, 9th ed. Salvat, Barcelona.
 32. Soria, M. L., G. Repetto, and M. Repetto. 1995. Revisión general de la toxicología de los metales, p. 293–359. In M. Repetto (ed.), *Toxicología avanzada*. Ediciones Diaz de Santos, Madrid.
 33. Spangber, J. V., and G. N. Cher. 1996. Developmental effects of barium exposure in a marine bivalve (*Mytilus californianus*). *Environ. Toxicol. Chem.* 15:1769–1774.
 34. Williamson, P. 1980. Variables affecting body burdens of lead, zinc and cadmium in a roadside population of the snail *Cepaca hortensis* Müller. *Oecologia* 44:213–220.
 35. World Health Organization (WHO). 1988. Dietary intake of Hg, Cd and Pb, assessment of chemical contaminants in food. Report on the results of the UNEP/FAO/WHO programme on health-related environmental monitoring (prepared in cooperation with MARC, London). WHO, Geneva.
 36. Xu, P., S. Huang, R. Zhue, X. Han, and H. Zhou. 2002. Phenotypic polymorphism of CYP2A6 activity in a Chinese population. *Eur. J. Clin. Pharmacol.* 58:333–337.
 37. Ysart, G., P. Miller, H. Crews, P. Robb, M. Baxter, C. De L'Argy, S. Lofthouse, C. Sargent, and N. Harrison. 1999. Dietary exposure estimates of 30 elements from the UK Total Diet Study. *Food Addit. Contam.* 16:391–403.
 38. Zyadah, M. A., and T. E. Adbel-Baky. 2000. Toxicity and bioaccumulation of copper, zinc and cadmium in some aquatic organisms. *Bull. Environ. Contam. Toxicol.* 64:740–747.