

Review

Outbreaks Where Food Workers Have Been Implicated in the Spread of Foodborne Disease. Part 5. Sources of Contamination and Pathogen Excretion from Infected Persons

EWEN C. D. TODD,^{1*} JUDY D. GREIG,² CHARLES A. BARTLESON,³ AND BARRY S. MICHAELS⁴

¹Department of Advertising Public Relations and Retailing, Michigan State University, East Lansing, Michigan 48824, USA; ²Public Health Agency of Canada, Laboratory for Foodborne Zoonoses, 160 Research Lane, Unit 206, Guelph, Ontario, Canada N1G 5B2; ³Bartleson Food Safety Consultants, P.O. Box 11983, Olympia, Washington 98508-1983, USA; and ⁴The B. Michaels Group, Inc., 487 West River Road, Palatka, Florida 32177, USA

MS 08-295: Received 17 June 2008/Accepted 18 July 2008

ABSTRACT

In this article, the fifth in a series reviewing the role of food workers in foodborne outbreaks, background information on the routes of infection for food workers is considered. Contamination most frequently occurs via the fecal-oral route, when pathogens are present in the feces of ill, convalescent, or otherwise colonized persons. It is difficult for managers of food operations to identify food workers who may be excreting pathogens, even when these workers report their illnesses, because workers can shed pathogens during the prodrome phase of illness or can be long-term excretors or asymptomatic carriers. Some convalescing individuals excreted *Salmonella* for 102 days. Exclusion policies based on stool testing have been evaluated but currently are not considered effective for reducing the risk of enteric disease. A worker may exhibit obvious signs of illness, such as vomiting, but even if the ill worker immediately leaves the work environment, residual vomitus can contaminate food, contact surfaces, and fellow workers unless the clean-up process is meticulous. Skin infections and nasopharyngeal or oropharyngeal staphylococcal or streptococcal secretions also have been linked frequently to worker-associated outbreaks. Dermatitis, rashes, and painful hand lesions may cause workers to reduce or avoid hand washing. Regardless of the origin of the contamination, pathogens are most likely to be transmitted through the hands touching a variety of surfaces, highlighting the need for effective hand hygiene and the use of barriers throughout the work shift.

In this article, the fifth in a series on food worker-associated outbreaks, the ways that pathogens can enter the food environment are discussed. In previous articles, the different outbreaks, the factors contributing to them, and the infective doses and carriage of the implicated pathogens have been described (46, 119–121). Pathogens that can infect food workers have multiple sources, and infected workers in turn become potential sources of contamination in food processing and preparation facilities. The pathogens then can become a part of the transient or resident flora. According to Snyder (116), transient pathogen sources include (i) fecal contamination on hands that remains after using the toilet, changing diapers, or cleaning up after pets; (ii) raw products (e.g., meat, poultry, fish, or unwashed fruits and vegetables); and (iii) infected cuts and boils that are touched or picked, or an infected fingernail. Transient pathogens are excreted in feces and various body fluids or tissues by persons infected or colonized by these pathogens. When diarrhea results in many liquid stools per day, hands easily become contaminated because billions of pathogen cells are present (6). However, many colonized individuals may be asymptomatic, or symptoms may be so mild that they are considered part of a normal digestive upset. Al-

though usually there are fewer pathogen cells per gram of solid stool than of liquid stool, the colonized individuals may erroneously think they are not a threat during food preparation and may become careless in their hygiene habits, allowing fecal contamination of food and the environment, including food contact surfaces. For viral pathogens, especially noroviruses and rotaviruses, vomitus can contain a concentration of particles similar to or greater than that in liquid diarrhea (24), and transmission and subsequent illness has resulted from cleanup procedures. Urine has occasionally been a means of pathogen transmission, typically only for organisms that invade the organs and blood supply, such as hepatitis A virus (HAV).

Resident sources of pathogens are less common and include permanent inhabitants on the epidermal skin layer, often protected in cracks and crevices where they cannot be removed easily through normal washing procedures. Although most microflora do not cause foodborne illness, *Staphylococcus aureus* can be a resident skin pathogen, multiplying in moist, warm areas such as the groin and frequently residing in the nasopharynx, from where it contaminates the skin on a regular basis through hand contact. This organism can be released through perspiration, aerosols from sneezing, and saliva onto food or food contact surfaces, cutlery, etc. Food workers also frequently develop

* Author for correspondence. Tel: 517-355-8371; Fax: 517-432-2589; E-mail: todde@msu.edu.

small cuts or burns in food preparation settings, and these wounds can become infected. Infected areas typically contain millions of *S. aureus* cells or group A streptococci, which can cause heavy contamination of foods handled by these workers or can cause impetigo. Staphylococcal nasal carriage seems to predispose people, particularly children, to the development of impetigo (71). Long-term or permanent carriers of pathogens such as *Salmonella* Typhi and some parasites also may be considered resident pathogen sources because shedding can occur on an intermittent basis over months or years. Injuries and infections reduce the inclination of workers to wash and dry their hands thoroughly and frequently. Quantitative data, where available, accentuate the importance of potential pathogen transmission through contaminated excretions and are discussed in detail in the following sections.

FECES AS A SOURCE OF CONTAMINATION FROM INFECTED AND COLONIZED WORKERS

Fecal sources from ill persons. Digested food is mainly liquid before reaching the colon, where water and nutrients are absorbed, leaving a semisolid stool. If the colon is damaged by enteric infections or toxins, the absorption of water is less efficient, resulting in watery stools with a moisture content of 50 to 90%. Bouts of diarrhea make it more likely that moist feces will penetrate toilet paper, contaminating hands and making effective cleanup more difficult. Levels of various pathogens in stool are listed in Table 1. *Salmonella* at up to 10^7 CFU/g can be found in feces of ill or early convalescent persons, who continue excreting for days or even weeks after the illness has resolved (126); infants can excrete this organism for much longer periods (6 to 8 months). Up to 10^9 HAV particles per ml or g of feces can occur, with concentrations highest before symptoms are apparent (95). Norovirus levels have ranged from 10^4 to 10^{10} DNA copies per g of feces from symptomatic patients (20). However, shedding can occur before the main symptoms are apparent, as demonstrated by Graham et al. (44) in dosed volunteers. Ozawa et al. (98) analyzed data from 55 norovirus outbreaks and 35 sporadic cases in Japan and found that genotype 4 (GII/4) strains were predominant and occurred with the highest viral load of all genotypes (mean values, 7.96×10^9 copies per g of stool). These authors also found that a large number of symptomatic food workers were infected with this strain. However, symptomatic and asymptomatic individuals for all genotypes had about the same viral load, 10^7 to 10^8 copies per g of stool, which confirms the importance of transmission by individuals who are infected but not ill. The magnitude of seroconversion was highest in individuals who vomited. Ward et al. (128, 129) found that up to 10^{11} viral particles were excreted from persons infected with rotavirus but only 10^6 to 10^7 particles were infectious, as demonstrated by cell culture. The concentration of protozoan parasites *Giardia* or *Cryptosporidium* in feces of infected persons can range from 10^5 to 10^7 cells per g; numbers are even higher for HAV (10^8 particles) and rotavirus (10^7 to 10^{12} particles) (41). Because 100 to 200 g of feces are produced per individual per day, Gerba (41) determined

that there could be as many as 10^{14} enteric pathogen cells in a single bowel movement of 100 g from an infected person. Because pathogens can be present in feces at levels of 10^4 to 10^{11} units per g (6), 0.1 mg of fecal material on the skin, which is barely noticeable, might contain up to 10^6 infectious bacterial cells, parasitic oocysts, or viral particles under diarrheic conditions. Gibson et al. (42) stated that the average American excretes 100 to 500 g of feces each day, and 0.1 g of residual fecal material remains on undergarments on a regular basis. This continual daily source of fecal contamination must be removed or excluded before workers touch food or food contact surfaces.

Fecal contamination of fingers and fingernails occurs more often in individuals with diarrhea, and Lin et al. (76), using ground beef or artificial feces as fecal matter surrogates, found that decontamination of the fingernail region was very difficult. Hand washing and alcohol gel had little effect, but use of a fingernail brush provided appreciable reductions. With $10^{-4.7}$ g of fecal contamination per hand and a pathogen load of 10^9 to 10^{11} units per g, as under diarrheal conditions, a 2- to 3-log reduction by washing and sanitizing would be insufficient to prevent a food worker from transmitting the pathogen to surfaces. Although fewer organisms probably are excreted when workers are asymptomatic or recovering from infection, any laxness in hand hygiene, such as not cleaning nails or only rinsing the hands, could lead to contamination of the food environment.

Sattar et al. (112) found that environmental surfaces remained contaminated with norovirus for 7 days after institutional outbreaks. Transfer of enteric bacteria from well individuals is illustrated by a study of postdefecation cleaning practices during which left and right hand fecal coliform counts were assessed for 90 individuals in Bangladesh (55). No toilet paper was used for cleaning; instead, traditional usage of the left hand and a small amount of water was used. Fecal coliform counts of the left hand averaged 3.93 log CFU, and those for the right hand counts averaged 2.99 log CFU. This degree of contamination on hands is related to the sanitation of latrines, which was critical for child survival (54). The authors speculated that the sitting area provided an opportunity for pathogens to transfer to hands and then into the mouth and was associated with the lack of water to clean the latrines. In rural communities in northern India, *Campylobacter* infection was significantly higher in persons who did not wash their hands with soap after perianal cleansing following defecation (10.2 versus 4.3%) (59).

In Santiago, Chile, where typhoid fever is endemic, family members of ill children belonging to a lower socioeconomic group were more likely to be carriers of bacterial enteric pathogens, such as enteropathogenic *Escherichia coli*, *Shigella* spp., *Salmonella*, and *Campylobacter* (13.8%), than were those individuals in a higher socioeconomic group (8.3%) (3). In homes with frequent carriers of these pathogens, unhygienic habits such as poor toilet care, inadequate hand washing, and children eating unwrapped candies were noted. The women who prepared the meals often were the carriers, and 39.5% of individuals who

TABLE 1. Levels of pathogens in clinical specimens and body excretions

Pathogen	Source of contamination	Contamination levels (per g or ml)	Reference
<i>Salmonella</i>	Feces while ill or during early convalescence	10^5 – 10^7 CFU	126
	Feces in late excretion period (infants excrete longer than do adults)	10^0 – 10^3 CFU	
	Feces during convalescence	6×10^3 CFU 15 days after illness 5×10^2 to 6×10^7 CFU (median, 6.0×10^6 CFU) <10 days after illness; 1.3×10^2 to 1.6×10^9 CFU (median, 1.0×10^5 CFU) 10–19 days after illness; 0×10^6 to 3.5×10^6 CFU (median, 2.5×10^4 CFU) 20–25 days after illness; 7.0×10^1 to 1.8×10^5 CFU (median, 1.4×10^2 CFU) 6–35 days after illness; 2.0×10^0 to 3.5×10^4 CFU (median, 5.5×10^3 CFU) 42–50 days after illness; 0×10^4 to 6×10^4 CFU (median, 2.5×10^4 CFU) 69–102 days after illness	85 103
	Pus in infected lesions	10^7 – 10^8 CFU (median) for intra-abdominal and anorectal and soft tissue infections (one sample with almost exclusively <i>S. aureus</i> and two with beta hemolytic streptococci)	70
<i>S. aureus</i> and <i>Streptococcus pyogenes</i>	Saliva in a sneeze from carriers	Typical person infected with streptococci: saliva, 10^0 – 10^6 CFU; <100 CFU/154 cm ² 1.5–9.5 ft (0.5–2.9 m) from sneeze source. One carrier sneezed twice (days 1 and 6): saliva, 3.2×10^6 to 7.5×10^6 CFU; 23–500 CFU/154 cm ² 1.5–9.5 ft from sneeze source	48
<i>S. pyogenes</i>	Saliva in a cough from carriers	10^3 – 10^6 CFU (1 of 20 persons infected with streptococci coughed 6 CFU/154 cm ² 9.5 ft [2.9 m] from cough source; most of the other 19 persons did not cough any streptococci)	48
Enteroviruses (e.g., coxsackievirus, echovirus, poliovirus)		10^3 – $10^{7.5}$ infectious particles; $10^{8.2}$ infectious particles	41
Hepatitis A virus	Feces, highest numbers before symptoms begin	10^9 virions	95
Norovirus Group G-I	Feces while ill	10^8 infectious particles	41
Norovirus Group G-II		2.2×10^4 to 2.9×10^{10} copies/g of feces; median, 8.4×10^5 copies/g	20
Norovirus Group G-I		2.5×10^4 to 7.7×10^{10} copies/g of feces; median, 3.0×10^8 copies/g	20
Norovirus Group G-I/4		GI 2.79×10^7 copies/g of feces	98
Norovirus Group G-II		GI/4 2.02×10^8 copies/g of feces	98
Norovirus Group G-II/4		GII, 3.81×10^8 copies/g of feces	98
Rotavirus	Feces and vomitus while ill	GII/4 7.96×10^9 copies/g of feces	98
		10^{11} particles excreted but only 10^6 – 10^7 infectious	129
		100 times more virus in vomitus than in feces	24
		8×10^9 to 10×10^9 infectious particles	4
		$>10^{12}$ infectious particles	8
		10^{10} – 10^{12} infectious particles in feces	41
<i>Cryptosporidium</i> spp.		10^8 – 10^9 oocysts in a single bowel movement	16
		10^6 – 10^7 oocysts; 3×10^9 oocysts/day	41
<i>Giardia lamblia</i> or <i>G. intestinalis</i>		< 10^9 cysts daily in stools	16
		1×10^6 to 5×10^6 cysts	41

cooked in families with children suffering from typhoid fever had *E. coli* on their hands compared with 16.1% for those who had no children ill with *Salmonella* Typhi infection. Khan et al. (64) found that the risk of children developing diarrhea from *Shigella* within these families was high whether the members were ill or not. The above data in-

dicate the levels of contamination that can result from a lack of toilet paper and clean water and how easily one hand can contaminate the other, increasing the risk of both symptomatic and asymptomatic infections.

The use of toilet paper, however, does not guarantee there will be no fecal contamination. Hutchinson (57) found

that when solid stools containing "scant growth" of *Shigella sonnei* from a convalescent patient were touched by fingers wrapped with a double thickness of toilet paper, the pathogen reached the bare fingertips in four of five attempts. When the experiment was repeated with loose stools with heavier growth from patients with acute or mild shigellosis, contamination of fingertips occurred on every occasion. Even more revealing is the work of Michaels et al. (89), who collected data from volunteers on four consecutive days employing coprostanol as a marker for fecal contamination. Coprostanol is formed by the microbial breakdown of cholesterol in the human intestine and has been used as a human fecal biomarker (107). Volunteers were instructed to use six stacks of toilet paper, each with six sheet layers, to clean after each defecation during each day of the study. The amount of coprostanol in each layer of toilet paper was measured as was the level of coprostanol per gram of feces for that test day. The last sheet in each stack closest to the hand frequently had coprostanol contamination, indicative of finger contamination (15 of 24 individuals, 63%), despite the fact that penetration through the entire stack of six sheets occurred much less frequently (7 of 24 individuals, 29%). This finding indicates that fingers tend to become contaminated by both poke through the toilet paper and, more importantly, circumvention of the barrier protection afforded by the paper, which becomes contaminated, and transfer of the fecal matter to the last toilet sheet. The fecal mass on the paper from the fingers, based on the amount of coprostanol detected, was calculated, and the geometric mean quantities of fecal material detected on hands was estimated at approximately $10^{-5.6}$ g. However, on one occasion the fecal mass was found to be much higher at >1 mg. From the work of Feachem et al. (35) linking enteric bacterial counts to fecal mass, the results of Michaels et al. (89) indicate that $10^{3.4}$ to $10^{4.4}$ fecal coliforms could be present on hands when initially contaminated. Previous investigators had shown lower levels of hand contamination ranging from 5 to 2,000 CFU (fecal coliforms, *Enterobacteriaceae*, or *E. coli*) (25, 26, 49, 55, 68). Nevertheless, any fecal contamination on the hands has the potential to harbor pathogens if they are present in the gut, and these studies indicate the limited effectiveness of toilet paper and the quantity and frequency of hand contamination during toilet paper use.

Carriage rates before, during, and after illness and in asymptomatic individuals are important to determine. Unfortunately, information is lacking for several of these parameters for pathogens potentially associated with food workers (121). Carriage rates range between <1 and 36%, and shedding can occur many days before symptoms appear, making exclusion of excreting employees from the food environment very difficult. This difficulty is compounded by the fact that many pathogens survive well in the environment, and reinfection can occur in endemic regions.

Asymptomatic fecal excretion. Worker health status is not necessarily linked to risk of contamination. Medus et al. (85) noted that 64 (53%) of 121 *Salmonella*-positive

food workers in 19 foodborne outbreaks in Minnesota reported not having recent gastrointestinal symptoms, and Sattar et al. (113) stated that possibly $>90\%$ of those individuals infected with viruses remain asymptomatic while discharging infective particles into their surroundings. Graham et al. (44) found that after inoculation of 50 volunteers with norovirus in buffered saline with bicarbonate to neutralize stomach acidity, 82% became infected, 68% were symptomatic (vomiting and/or watery diarrhea), and 32% were asymptomatic; the overall infection rate was higher than previously recognized, with a high rate of subclinical infection, prolonged virus shedding, and a significant correlation of magnitude of antibody responses with severity of symptoms. Although early studies indicated that norovirus shedding began with onset of symptoms and did not extend beyond 72 h, Graham et al. found that after inoculation of the 50 volunteers, virus first appeared in the stool at 15 h, and excretion peaked at 3 days and continued for at least 7 days.

In outbreak scenarios, those involving asymptomatic workers are as frequent as those involving workers who were ill (46, 119, 120), and excretion of pathogens can occur both before the illness and after the patient has recovered. During HAV infection, as many as 10^9 viral particles per gram of feces may be shed from >24 h to 2 weeks before symptoms develop (43, 95). Peak fecal excretion occurs before the onset of jaundice and other symptoms and then declines after jaundice appears. HAV excretion, however, can continue for many weeks after recovery and for up to several months in children. A food worker was still shedding Norwalk-like virus 10 days after resolution of illness following an outbreak, indicating that the recommendations of exclusion of workers for 48 to 72 h after recovery may not be sufficient to prevent transmission (99). This finding was supported by a study of 99 individuals infected with Norwalk-like (noro) viruses (110); the virus was detected in 26% of the patients up to 3 weeks after the onset of symptoms; the highest proportion of long-term shedders (38%) were children less than 1 year of age. Long-term shedding was not associated with increased severity of disease or prolonged duration of clinical signs. Children have been identified as long-term excretors of enteric viruses, and these infected children may infect parents in the home, as demonstrated in a recent survey (46, 119, 120).

Hutchinson (57) found *S. sonnei* on the hands of 49% of young children in nursery schools in infected communities after these children had used the toilet, and over one-third of these children touched their faces and mouths or sucked their fingers. She also found that toilet seats in schools, nurseries, and private homes in these same communities were a means of transmission; 32% of these seats were contaminated with *S. sonnei*. Mothers in developing countries frequently carry *E. coli* as a result of caring for infants (9, 84, 87). In Thailand, enterotoxigenic *E. coli* was found on the hands of 6 of 42 mothers and of 37 of 50 children (30). These infected women typically prepared the meals for the family. Pether and Scott (103) examined stools of patients convalescing from *Salmonella* infections

from 6 to 102 days after symptoms ceased. As expected, stool counts tended to decrease over time but not in a consistent way. Median values for days after illness were 6.0×10^6 CFU at < 10 days, 1.0×10^5 CFU at 10 to 19 days, 2.5×10^4 CFU at 20 to 25 days, 1.4×10^2 CFU at 26 to 35 days, 5.5×10^3 CFU at 42 to 50 days, and 2.5×10^4 CFU at 69 to 102 days. The specimens that remained positive between 42 and 102 days had higher median counts than did some earlier specimens, indicating that an on-going colonization within the intestines may occur, allowing a continual supply of organisms for excretion, although all patients were asymptomatic. However, the counts of all patients diminished over time. The authors discounted concern about these numbers, assuming that the infectious dose was relatively high and that hand washing would remove any organisms present on hands after defecation. However, the large number of reports of salmonellosis outbreaks associated with food workers call this reasoning into question (46).

Many factors may influence whether pathogen ingestion will lead to symptomatic or asymptomatic infections. Noda et al. (97) conducted a statistical analysis of norovirus outbreaks in Japan and found that the attack rate in oyster-associated outbreaks was significantly higher than that in food worker-associated outbreaks, but the number of cases was greater in the latter (median, 40 versus 17 for the oyster-linked outbreaks). Most of the facilities involved in the food worker-associated outbreaks were large restaurants such as hotels that served food for parties and schools. The authors concluded that the attack rate in foodborne outbreaks of norovirus infection may be influenced by differences in implicated foods (sewage-contaminated oysters versus food worker-contaminated restaurant items), susceptibility of the host for norovirus infection, and pathogenicity of the strains. They also noted that a single norovirus genotype was responsible for most of the food worker-associated outbreaks (typically GII/4 or GII/3), whereas multiple norovirus genotypes were frequently involved in the oyster-associated outbreaks (accumulated over time in the oyster beds). The attack rate in 27 outbreaks associated only with GII/4 was lower than that in 136 other outbreaks, suggesting that GII/4 noroviruses cause asymptomatic infection more frequently than do other norovirus genotypes.

Well-investigated norovirus outbreaks sometimes demonstrate the complicated nature of infection spread, which can depend on prolonged asymptomatic shedding by some infected persons, the environmental stability of the implicated strains, and the lack of lasting immunity in persons who have been infected previously (15). One norovirus outbreak that occurred at a family reunion in West Virginia in 2006 involved attendees from Florida, Maryland, New York, Pennsylvania, Virginia, and West Virginia (18). The investigation indicated that a combination of person-to-person and foodborne transmission of two strains of norovirus, likely introduced by persons from two different states and subsequently into at least two food items, was the probable cause of these illnesses, highlighting the challenge of investigating and controlling norovirus outbreaks.

The carriage rate for *Salmonella* in England was de-

termined by testing more than 2,800 stool specimens; 5% of ill individuals who presented to general practitioners were carriers compared with 1.1% of those individuals in the community that had loose stools or vomiting and 0.4% of those with no apparent illness (123). To determine the carriage rate for *Salmonella* in food workers, 331,644 fecal specimens were collected from workers in food factories, hotels, restaurants, and supermarkets in one region in Japan (94); 0.032% of these samples contained *Salmonella*, and the most common serovars were Infantis, Corvallis, Enteritidis, and Agona. There were only two distinct pulsed-field gel electrophoresis (PFGE) profiles among the 16 *Salmonella* Corvallis isolates, indicating that the sources of infection for the many different types of food workers were not extensive. In Japan, these same serovars have been associated with broiler chickens and eggs. In the same community, a survey of diarrheal stools from clinical patients revealed a higher rate of infection (1.25%) but not the same serovar distribution (mainly *Salmonella* Enteritidis). This research indicates that relatively few asymptomatic food workers may carry *Salmonella*, but the reservoir is probably animal sources such as poultry or eggs, which workers contact while preparing food. In an outbreak in Jordan, *Salmonella* Enteritidis infection was traced to one food worker in a hospital (65); 183 of 619 hospital employees and patients met the case definition after eating mashed potatoes for lunch. No diarrhea was reported by kitchen employees up to 3 months before the outbreak. However, the investigation revealed that 11 kitchen employees were positive for the same *Salmonella* serovar, but only 1 of them was responsible for preparing the mashed potatoes on the day of the lunch. Although the potatoes were mechanically mashed, this worker had mixed the potatoes with milk by hand and wore no gloves. He was apparently asymptomatic at the time but was the first patient to present to the emergency room about 16 h after the meal. Because the potatoes were prepared on the morning of the luncheon, there would have been only a few hours for *Salmonella* growth after the mashing.

Three percent of typhoid fever survivors become permanent carriers, harboring the organisms in the gallbladder, biliary tract, or rarely the intestine or urinary tract (12). Horwood and Minch (56) found *E. coli* on the hands of 13 of 34 food workers from foodservice establishments; those individuals without *E. coli* were waitresses and others not intimately handling food. de Wit and Kampelmacher (25) proposed that *E. coli* and fecal coliform counts increase as human involvement with moist foods of animal origin increases, because bacterial counts on the hands of more than 250 food workers and non-food workers revealed that hand contamination was more related to the type and nature of raw animal products they handled than to toilet hygiene. A survey of the hands of food processing plant workers revealed *E. coli* at >20 CFU per hand sample taken before and after defecation and hand washing (4 and 25%, respectively) (26). However, the average level of *E. coli* before and after collection of positive samples was about the same: 2.30 log CFU per sample. These quantitative data indicate that toilet use itself is not the only source of *E.*

coli for workers. The authors concluded that asymptomatic *Salmonella* excretors have only a low risk of transferring *Salmonella* from their stools by their hands to food. Contamination of hands with *Salmonella* is more likely to result from inconsistent hand washing during preparation of meat (69).

VOMIT AND URINE AS SOURCES OF CONTAMINATION FROM INFECTED AND COLONIZED WORKERS

Vomitus. Vomiting is the forceful expulsion of stomach contents through the mouth and is a symptom of a number of intestinal diseases caused by emetic toxins produced by infectious or toxigenic agents acting on the vomiting center of the brain. Nausea usually precedes vomiting. The onset of vomiting may be rapid and uncontrollable, often occurring during or shortly after meal time, which is indicative of a short incubation period for an emetic toxin in the food or the end of the prodrome stage for noroviruses. Noroviruses frequently cause projectile vomiting, and electron microscopic examination has revealed that a minimum of 10^6 virus particles per ml may be present in vomitus (51). Cotterill et al. (24) reported that rotavirus concentration in vomitus can be 100 times that in feces and that Norwalk virus is also found in vomitus. These findings suggest why vomiting is an important means of transmission for enteric viruses. The causative agent may be isolated from gastric contents, but acidic gastric secretions may destroy the pathogen before it can be isolated. Environmental cleaning and disinfection are vital for preventing disease transmission.

Vomitus can contaminate the environment, rest rooms, hand washing stations, and food worker clothing (22, 31, 77), and contamination of food may occur directly via aerosolized particles or hands. Noroviruses also can be transmitted to other individuals via aerosols produced during vomiting (14, 51, 81). Aerosol transmission resulted in infection of individuals that had simply walked through an emergency department while a vomiting patient was being examined (114). If an individual vomits within the kitchen area, fellow workers may become infected through inhalation of the agent or by hand-to-mouth transfer from contaminated environmental surfaces. The ease of airborne transmission of norovirus also was demonstrated in an outbreak at a restaurant where no food source was implicated but analysis of the attack rate revealed an inverse correlation with the distance from a person who had vomited (81). Although vomiting usually precedes unexpected bowel movements, vomiting and diarrhea may occur at the same time, making it difficult to determine which body fluid resulted in the transmission. This difficulty was demonstrated when a worker at a catering company frosted many cakes during a 6-h shift, although he had five episodes of diarrhea and two of vomiting at the beginning of his shift (73). Residual vomitus or feces on his hands after cleaning up from an attack was the likely source of the norovirus as he mixed the frosting with his hands. His actions infected nine other employees, who in turn caused another 3,000 illnesses from consumption of the cakes they prepared. In another scenar-

io, a concert attendee vomited four times in an auditorium waste bin, on the carpet, and in the adjacent men's toilet. Before the next day's performance, the area was cleaned with a vacuum and a spillage compound without disinfectant. The next day, school children who sat on the same level of the auditorium where the infected attendee had been seated contracted norovirus infections from the viral particles still remaining on the seats and other areas of the auditorium (31). The staff who had worked on the upper tiers that same evening or had helped clean up after the vomiting and the family members who accompanied the infected attendee also were infected. The most likely cause of the 300-case outbreak was inadequate cleaning and disinfection of hard surfaces, carpets, and soft furnishings contaminated by vomit, leading to aerosols. In another outbreak, a kitchen assistant vomited into a kitchen sink; he tried to disinfect it with a chlorine-based sanitizing powder but failed to inform his supervisor (101). Boiled potatoes for a potato salad were cooled in the same sink the next day by another staff member, and the contaminated salad infected 56 wedding guests.

Another more recent example illustrates the importance of preventing vomiting while working in the kitchen environment and the importance of appropriate cleanup if vomiting occurs. In Michigan in 2006, a norovirus infection outbreak occurred in late January and early February 2006 (19). At least 364 restaurant patrons became ill with gastroenteritis after dining at a restaurant where employees had reported to work while ill. Low-level transmission occurred in the week before 28 January, with seven patrons who dined at the restaurant between 21 and 27 January meeting the case definition. From 21 January to 3 February, exposure to the virus likely occurred by contact with contaminated surfaces and objects. On 28 January, a line cook vomited at home before reporting to work at 11:00 a.m. and then vomited again into a waste bin beside the frontline workstation at approximately 2:00 p.m. while preparing antipasti platters, pizzas, and salads. After vomiting, he remained on site (but off the cooking line) and left work at 4:15 p.m. This person also reported to work on 29 January from 11:00 a.m. to 4:30 p.m. while still experiencing loose stools. Because of the open physical layout of the restaurant, no barrier impeded airborne spread of the virus from the kitchen to the main dining area. Direct foodborne transmission also was likely, as demonstrated by Bidawid et al. (7), who found that 46, 18, and 13% of surrogate feline calicivirus was transferred from contaminated fingerpads to ham, lettuce, and metal disks, respectively. The antipasti platter was one of many dishes that the infected line cook prepared, and statistical analysis revealed a significant association between the platter and the ill persons. Attack rates increased after the cook vomited and among employees who worked on 28 January, with a higher percentage of line cooks becoming ill compared with servers. Other environmental contamination probably contributed to transmission. The investigation revealed deficiencies in employee hand washing practices, cleaning and sanitizing of food and nonfood contact surfaces, temperature monitoring and maintenance of potentially hazardous food (even though

this issue would not impact the multiplication of norovirus directly), and maintenance of hand-sink stations for easy accessibility and proper use. A quaternary ammonium-based sanitizer that was ineffective against norovirus was used to clean the restaurant. Management was advised by the local health department to disinfect the restrooms and all surface areas within at least a 25-ft (7.6 m) radius of the vomiting site with a bleach solution. These actions were effective for preventing further illnesses.

Urine. Urine has been occasionally reported as a source of some pathogens. An HAV infection outbreak at a U.S. Navy base occurred when salad was intentionally contaminated by a cook with a personality disorder. As a mark of insubordination, he deliberately urinated into the salad during its preparation while he was still asymptomatic and transmitted his infectious HAV particles to the men eating the salad (62). *Salmonella* Typhi and *Salmonella* Paratyphi also can be detected in urine during systemic infection (11), highlighting the necessity of effective hand hygiene even for food workers who use the toilet only for urination.

SKIN AS A SOURCE OF CONTAMINATION FROM INFECTED AND COLONIZED WORKERS

Skin infections and secretions. Skin, the largest organ of the body, may have a large exposed area for abrasions, cuts, and punctures to occur, and these wounds can develop into infected boils and lesions. Boils and other deep lesions are typically caused by invasive infections by *S. aureus* carrying Panton-Valentine leucocidin genes (53). Extensive involvement of hands and arms is difficult to avoid during food preparation and food service, and this involvement has led to contamination, mainly by *S. aureus*, and subsequent outbreaks. Work in the kitchen environment can result in rough skin and small cuts, which may become infected. Boils also can occur through rubbing of skin by clothing or removal of hair follicles. These boils tend to occur on parts of the body other than the hands but act as reservoirs for *S. aureus* hand contamination through touching. Although workers tend to ignore these minor irritations, if the sores or boils are on the hands or arms, affected workers often reduce or stop hand washing to reduce the pain and inconvenience of removing bandages. As the infection progresses, pus can spread infectious organisms over skin surfaces and increase the risk of food contamination.

Open cuts, sores, and eczematous or excoriated lesions were associated with 25 outbreaks (37% caused by staphylococci and 24% caused by streptococci) reviewed by Greig et al. (46). The pathogens were carried by food workers who in turn contaminated custards, ham and cheese sandwiches, and potato, macaroni, egg, rice, tuna, and chicken salads. A home-catered social in Saskatchewan in 1988 resulted in 49 cases of *S. aureus* infection. The same phage type was isolated from potato salad, six patients, and six food workers; the same strain was isolated from an infected cut of a worker who mixed the salad with bare hands (118). In Georgia in 2000, PFGE analysis revealed that *S. aureus* from the food, ill patrons, and a plastic bandage

from a worker's finger were indistinguishable (74). An outbreak of group A streptococcal pharyngitis at a school banquet was traced to contaminated macaroni and cheese. The same strain of *Streptococcus pyogenes* was isolated from affected individuals and a food worker's hand wound obtained in a fistfight 2 weeks earlier (34). Culture-positive purulent lesions can contain different pathogens, including enterococci, at up to 2×10^8 CFU/ml of fluid (70) (Table 1). *S. aureus* counts from persons with infected skin lesions may reach up to 10^5 CFU/cm² (96), but direct contact with food alone is insufficient for an infective dose or for toxin production in food, and a subsequent growth period is required to reach the high numbers necessary to produce toxin. However, the presence of infected open cuts and sores is not a prerequisite for contamination of food (52). Symptomless carriage of *S. aureus* and *S. pyogenes* in the nares and on the skin is common, and small numbers of these pathogens can continually contaminate surfaces, including food. Typically, temperature abuse of the food during storage is an additional requirement, and Bryan (13) found that such abuse was a contributory factor in 40% of foodborne outbreaks of *S. aureus* infection, where the storage time was sufficient for the organism to produce a heat-stable enterotoxin.

Skin infections on hands, however, do not necessarily prove that these lesions are the source of an outbreak. Phage typing of the skin may indicate a different strain from that in the implicated food. In one scenario, both *S. aureus* and *Salmonella* were found in leftover turkey but could not be traced to skin infections or colonized anterior nares of food workers (86). In another outbreak, 36 persons rapidly became ill with *S. aureus* intoxication after eating vanilla slices (thick vanilla-flavored custard in puff pastry) from a bakery (36). One of these persons developed five *S. aureus*-infected skin lesions on her legs 1 day after she ate the contaminated food. Investigators surmised that her legs became contaminated with diarrhea, and the pathogen subsequently invaded the skin. In another situation, although a food worker had severe cellulitis of his hands, he could not be linked to an outbreak of *Streptococcus* pharyngitis (33).

Molecular and toxin typing can help determine the cause of an outbreak, even when multiple bacterial strains have been isolated. After a staphylococcal food poisoning outbreak affecting 10 of 356 students who attended a high school breakfast, 27 *S. aureus* isolates producing different enterotoxin types were recovered from 7 patients, 2 food workers, and food leftovers (130). Based on genotyping results and enterotoxin typing, three distinct strains were identified. Two of these strains were found in specimens from patients and a hand lesion of a food worker, indicating the precise source of the contamination. However, not all outbreaks involving infected cuts are caused by staphylococci or streptococci. A food handler who had painful skin lesions on his hands was associated with an outbreak of HAV infection lasting several weeks (127). He prepared bread products and sandwiches for an English village shop while wearing visibly soiled bandages on his hands, which prevented him from washing properly. Viral particles were

likely excreted onto his hands during his mild illness while he continued to work preparing the bakery items.

Dermatitis and other skin diseases. Impetigo can be caused by either *S. aureus* or *Streptococcus* spp. In streptococcal cases, *S. pyogenes* colonizes the skin and can be transferred from one lesion to another and from one mucous membrane surface to another (28). The lesions eventually develop a characteristic honey-colored crust and heal without scarring. These lesions most commonly occur on the face or extremities. Colonization of the mucous membranes of the nose or throat occurs about 2 to 3 weeks after *S. pyogenes* appears on the skin, and the bacteria can survive an average of 10 days on the skin before any infected lesions appear. Occupational skin diseases (contact dermatitis being most frequently reported) account for approximately 40% of all occupational illnesses according to the U.S. Bureau of Labor Statistics, and contact dermatitis accounts for more than 90% of workers' compensation claims for occupational skin disease (82, 83). Occupations where hands are frequently immersed in water, as in food industries, have an increased potential for skin problems (75, 82). Age, sex, and skin allergy history also are important for food workers.

Women are more likely to have an atopic history and/or become sensitized to allergens than are men. Older individuals generally have dry, thin skin that is prone to cracking, resulting in more frequent cases of irritation and pain, which discourage hand washing. Atopic dermatitis (an allergic hypersensitivity such as eczema that affects various parts of the body) results in increased risk for sensitization and irritation by hand sanitizers, disinfectants, and metals such as chromium and nickel in stainless steel watches or jewelry. Atopy, which tends to have a strong hereditary component, also can be associated with food allergens, antimicrobial ingredients in soaps, allergens associated with gloves and antimicrobials, alcohol in instant hand sanitizers or hand antiseptics (10, 38, 39, 125, 136), glove powders, and pyrogens from sterile latex gloves (72, 131, 133).

Animal testing has revealed that several components are needed for sensitization of the skin leading to atopic diseases, including skin damage, an allergen potentiator, and occlusion of the skin (66). In the food processing and service environment, known allergens are present, and the skin can be damaged with brushes, cut, scratched, scalded, or burned or can come into contact with caustic chemicals. When hands are placed in gloves, the occlusion provided is conducive to the development of allergic contact dermatitis (105, 106). These issues highlight the importance of skin health in the overall control of transmission of human pathogens from food workers (88).

Sweat. Sites rich in sweat glands are well-recognized secondary niches of *S. aureus* (29). Although contamination of food by sweat has not been identified directly in an outbreak report, perspiration can allow normal skin microflora, including *S. aureus*, to be transferred to food. Staphylococci can migrate through perspiration to contaminate hands, jewelry, and work surfaces (1, 2, 58, 90). Moist hands facilitate the transfer of microorganisms, and sweat

can transport potential pathogens colonizing the skin (100). In a hot kitchen environment, droplets of sweat containing *S. aureus* or *S. pyogenes* from the face and nose may directly contaminate food being prepared or indirectly contaminate food after perspiration is wiped away with the hands. If temperature abuse occurs, foodborne illness can result (21).

NASOPHARYNGEAL AND OROPHARYNGEAL SECRETIONS AS SOURCES OF CONTAMINATION FROM INFECTED AND COLONIZED WORKERS

S. aureus. Enterotoxigenic staphylococci have been isolated from about 30% of the population and 12 to 58% of food workers in different parts of the world (2, 102, 104). An estimated 25 to 50% of food workers are cutaneous or nasopharyngeal carriers of *S. aureus*, and 15 to 20% of these strains are enterotoxigenic (50, 91, 109). After an outbreak, these percentages may be higher (27). Food workers can harbor strains that produce both high amounts of toxin and more than one enterotoxin (108). Reina et al. (109) found that strains isolated from the skin were more enterotoxigenic than those from the respiratory tract or other clinical specimens; 61.7% of these strains produced one or more enterotoxins (mostly enterotoxins C and B). The evidence pointing to food workers as the source of many outbreaks of staphylococcal infection is compelling. Matching staphylococci phage types were obtained from the food, the affected individual, or the handler in 27% of outbreaks from 1977 to 1981 in the United States, and 67% of food workers were asymptomatic carriers of the same phage type as that in the implicated food (52). In a study of 358 outbreaks and sporadic cases of staphylococcal food poisoning in the United Kingdom between 1969 and 1990, strains from 79% of the incidents produced staphylococcal enterotoxin A or enterotoxin A plus another enterotoxin (132). The level of *S. aureus* in the implicated food ranged from none detected to 1.5×10^{10} CFU/g (median, 3.0×10^7 CFU/g).

Hatakka et al. (50) studied the carriage rates of *S. aureus* in airline catering staff in Finland and found prevalences in noses and on hands of 29 and 9%, respectively. Forty-six percent of the strains isolated from workers were enterotoxigenic; 6% from their hands and 12% from their nasal cavities. PFGE results indicated that the same strain tended to be found in noses and hands, but detection of carriers is best done through nasal cavity sampling (50). The most prevalent enterotoxin producer was type A, which accounted for 34.9% of all the enterotoxigenic strains. The level of toxin production was somewhat lower than that found by Reali (108), who found that 76 to 80% of strains from nasal swabs of healthy carriers and from fecal and/or urine specimens and 100% of isolates from infected wounds and cutaneous and urogenital infections were toxin producers. However, the level of toxin production was similar to that in a more recent study of food workers in Santiago conducted by Figueroa et al. (37), who found that 54% of the strains were enterotoxigenic, with producers of enterotoxin A dominating. In another survey of food workers in

Botswana, 57.5% were positive for *S. aureus* (78). Of the 204 *S. aureus* strains isolated, 30.9, 44.6, and 24.5% were from the hand, nasal cavity, and face, respectively, and 21% of the isolates were enterotoxigenic, a lower percentage than in the Chilean and Finnish studies. These data indicate that many food workers are carriers of enterotoxigenic *S. aureus*, and screening of these workers and administration of antibiotics or removal of these workers from the food preparation area is impractical.

***S. pyogenes* (group A streptococci and beta-hemolytic streptococci).** The nasopharynx (area between the nose and throat) and the oropharynx (oral part of the pharynx reaching from the soft palate to the level of the hyoid bone) are potential breeding areas for microbial pathogens. *S. aureus* and *S. pyogenes* may reside in these areas without causing any indication of colonization, and coughs, sneezes, and wiping of the nose or mouth can transfer these bacteria to the hands. Even if cultures of these areas are positive for *S. pyogenes* or *S. aureus*, it is difficult to determine during outbreak investigations exactly how food has been contaminated. The organisms may originate from sputum or respiratory droplets or by direct hand contact by food workers. Persons whose nasal passages are colonized by beta-hemolytic streptococci are more likely to transmit infection than those with positive throat but negative nasal cultures, and such carriers disseminate the streptococci in highly contaminated nasal secretions that reach the environment, chiefly via the hands when blowing noses but also through contaminated handkerchiefs, clothing, and bedding (48). One chronic carrier had 2.4×10^7 CFU from a nose blow into a handkerchief compared with 3.8×10^4 CFU in his saliva (48). Bar-Dayan et al. (5) elucidated the differences between the symptoms of patients with endemic airborne streptococcal pharyngitis and those of patients with epidemic foodborne streptococcal pharyngitis. The patients with foodborne streptococcal pharyngitis had a significantly higher frequency of sore throat, fever, pharyngeal erythema, tonsillar enlargement, and submandibular lymphadenopathy and a lower frequency of coryza and cough compared with the patients with endemic airborne streptococcal pharyngitis. Although both foodborne and airborne streptococcal infection caused upper respiratory tract infection, the clinical manifestation of foodborne streptococcal pharyngitis was more severe and more confined to the pharynx than that of the more endemic airborne disease. Involvement of the nasal mucosa and bronchial tree was more common in cases of airborne streptococcal pharyngitis than in cases of the foodborne disease.

Experimental observations revealed that saliva in a cough of one carrier contained *S. pyogenes* at 10^3 to 10^6 CFU/ml and delivered 6 CFU/154 cm² at a distance of 9.5 ft (2.9 m), but most coughs from and loud talking by carriers spread few streptococci into the environment (48). Although sneezing is generally recognized as a risk factor and sneeze guards are put around food displays, few instances have been reported in which sneezing has been the direct cause of foodborne illness, and no reported foodborne illnesses have been associated with coughing (117). However,

sneezes did allow the beta-hemolytic streptococci from 7 of 20 carriers to reach as far as 9.5 ft, and the organisms were persistent enough in the air to be cultured up to 16 min later (47). High numbers of *S. pyogenes* cells can be expelled in large droplets during a sneeze, settling in a zone of up to 1.5 ft (0.5 m) from the carrier source. A single sneeze by a nasal carrier of *S. pyogenes* is capable of expelling up to 5.0×10^7 CFU, whereas a sneeze by a throat carrier can expel around 1×10^5 CFU and a cough may expel 40,000 to 50,000 CFU (48). When outbreaks have occurred, they typically have been associated with streptococcal infection of a food worker, as demonstrated in the following two episodes. Egg salad served at a charity luncheon in Baltimore in 1957 infected 600 people, who contracted streptococcal pharyngitis (33). It was assumed that a woman who cooked and peeled the boiled eggs, at least in part, contaminated them with *S. pyogenes* by sneezing, although sneezing was not observed. A similar situation occurred in 1990, when egg salad served for lunch at a military base caused 61 cases of streptococcal infection (79). The same strain of *S. pyogenes* was isolated from the salad and the worker who had prepared the food while ill. The egg salad was stored in a refrigerator with a door that did not close properly, resulting in inadequate cooling of the salad. Sneezing or throat secretions were associated with the egg salad contamination. Following initial contamination, temperature abuse allows exponential growth of organisms. In their review of 18 streptococcal foodborne outbreaks, Katzenell et al. (63) found that cold salads were the main vehicle of infection, and the outbreaks occurred either in warm climates or during summer months in more equitable regions where there were opportunities for rapid growth. Almost all of these salads were prepared 24 h in advance of the meal time, and some of the foods were kept out of the refrigerator for several hours before they were served, providing opportunities for growth. In one experiment, streptococci isolated from throats of patients with pharyngitis were grown at room temperature on a medium containing eggs; these bacteria multiplied by 8 log units in 40 h. In outbreaks for which the data have been published, food workers typically had sore throats and/or pharyngitis either before or during the meal preparation; one of the workers had an infected lesion. Asymptomatic carriage was rare. Although the risk of food contamination is high when food is touched by colonized food workers (13), it is not always easy to discover how a particular worker becomes infected and how that worker then contaminated the food, causing illness. An effective approach to preventing transmission of staphylococcal and streptococcal infections is training food employees to recognize infected lesions and utilize double waterproof barriers, such as a bandage and finger cot to cover an infected lesion on a finger, and to follow good personal hygiene practices (124). Community-acquired methicillin-resistant *S. aureus* (MRSA) is spread by direct contact, in aerosols, and on fomites. However, occasionally MRSA is linked to food. In one serious hospital outbreak in The Netherlands, 21 hematology patients (77.8%) developed clinical disease, and 5 died (67). Subsequently, MRSA was detected in food and in the throat of

one health care worker who prepared food for the hematology patients. Food contaminated by this worker most likely caused the first case of MRSA septicemia, and the outbreak strain probably was transmitted to the surgical unit by a colonized nurse. Another outbreak investigation of MRSA infections revealed that a worker at a convenience store was associated with three cases of MRSA infection that occurred after consumption of pork and coleslaw (61). The worker apparently became colonized while visiting an elderly relative and remained culture positive without showing signs of illness for more than 8 months.

Work exclusion criteria. Postsymptomatic workers may continue to excrete pathogens, but at lower rates, for days, weeks, and occasionally years. Although shedding duration can be measured only by a regular stool-testing regimen, this approach often is ineffective, costly, and not recommended, especially for individuals hospitalized for >3 days (55, 92, 111, 115). An outbreak involving an airline caused 290 cases of salmonellosis during a period of 6 days (135). Unfortunately, the infected employee was not identified during regular inspection and testing of the establishment. Stool testing and exclusion of workers has been an issue for many decades, and recommendations differ among jurisdictions. Winter (134) recognized the limitations of annual testing of food workers for pathogens but understood that some form of certification for workers within the European Economic Community was necessary. A better approach would be for employees in large companies to be examined by a nurse or physician and certified to be free from infectious agents such as *Salmonella* Typhi, HAV, *S. pyogenes*, and *Mycobacterium tuberculosis*. Winter (134) found that potential employees may be rejected (5 to 50% of them) because of decayed teeth, infested hair, dirty body and clothes, and dirty or bitten fingernails. Pether and Scott (103) examined the stools of convalescent carriers of different *Salmonella* serovars and suggested that although some of their fingers were contaminated, the clinically well food worker with formed stools should be allowed back at work without further examination of fecal specimens.

Generally, pathogen-negative stool samples, either pre-employment or from an employee recovering from a diarrheal illness, are not necessary conditions of employment or return to work, with the exception of typhoid and paratyphoid infections and infection caused by verocytotoxin-producing *E. coli* (VTEC) (32). However, typhoid and paratyphoid fevers and HAV infection need special consideration because of the severity of the illnesses and because individuals can continue to carry and excrete the etiologic agents for long periods after recovery, with a consequent risk of food contamination. There is no consistent approach among different jurisdictions and associations for stool testing and worker exclusions. Typically, three negative stool specimens are required before an infected food employee can return to work. When VTEC infection, including *E. coli* O157:H7 infection, is identified in a food worker, the worker should be excluded from work until the bowel habit has been normal for 48 h and two negative fecal samples taken 48 h apart have been obtained. Symptomless contacts of a

person with HAV infection can continue food handling, but workers who have symptoms of hepatitis, have been in an outbreak or have been associated with family members suffering from HAV infection, or have traveled to regions where HAV is endemic would be excluded from work until they have a medical release based on laboratory testing. The Food Marketing Institute (40) has published similar recommendations but also stated that the single most critical way that food workers can prevent the spread of HAV is to wash hands thoroughly and often upon arriving at work, after using the toilet, after breaks, and at many other times during food preparation. Disorders such as Crohn's disease or ulcerative colitis are not a barrier to employment as a food worker, even though such disorders may result in diarrhea. Such workers must be made aware of the need to seek medical advice and notify their managers if any change from their normal bowel habit occurs, because such a change must be assumed to be infectious until proven otherwise (32). According to the Food Code (124), an employee is required to report *Salmonella*, *Shigella*, *E. coli* O157:H7, and HAV infections or illnesses such as diarrhea, vomiting, fever, jaundice, or sore throat with fever. Employees are required to provide written medical documentation that they are free from the four specified infectious agents via stool testing (17). After an outbreak in Michigan, new guidelines were produced recommending that foodservice workers with suspected norovirus not return to work until they are asymptomatic for 48 to 72 h, longer than the previously recommended 24 h (17). Thus, as new data from investigations of outbreaks are generated, they may influence exclusion policies.

CONCLUSION

Enteric organisms from fecal sources are excreted during an infection, whether the individual is symptomatic or not, but the infection exists over a limited time period (usually hours to weeks). Many of these enteric pathogens are of concern to food workers, e.g., *Salmonella*, *Shigella*, and noroviruses. These pathogens can contaminate the hands after defecation or from touching fecally soiled clothing or surfaces. Changing messy diapers, cleaning after episodes of vomiting or diarrhea by family members, and contact with sick or healthy pets also are well-established risk factors. Michaels et al. (89) found that fecal matter as detected by coprostanol can reach the fingers despite the use of up to six layers of toilet paper under conditions of normal perianal cleaning. Thus, the use of toilet paper is no guarantee of uncontaminated hands, and the situation is worse for individuals with diarrhea, especially when the individual has long or artificial fingernails, which are notoriously difficult to clean. Infected workers that are excreting a pathogen also risk infecting their colleagues during shared operational activities. Contamination levels may be higher when workers handle raw foods of animal origin than after toileting. Handling of raw foods can occur in a meat department of a supermarket or in a restaurant, where *Campylobacter*, *Salmonella*, and *E. coli* O157:H7 can be common contaminants.

Similar scenarios may occur through handling crops,

foods, or food ingredients contaminated with animal manure, animal intestines, sewage, nonpotable water, etc., which may occur during harvesting, hunting, or butchering. Lues and Van Tonder (80) found that both aprons and hands of food workers may be vehicles of cross-contamination, but there was no statistical correlation between the organisms on hands and those on aprons, indicating other sources of contamination for aprons. The impact of environmental sources in food environments, including raw foods of animal origin, were further discussed by Todd et al. (122).

Parasites tend to be overlooked as a source of foodborne infections, particularly in developed countries where they rarely cause outbreaks. However, *Cryptosporidium*, *Giardia*, and *Cyclospora* all have caused foodborne outbreaks, although the role of the food preparer in transmitting these agents has not always been clear. Carriage rates of both diarrheic and healthy populations can be >5% (123), providing many opportunities for fecal contamination of foods and contact surfaces. No known outbreaks of *Entamoeba histolytica* infection have been reported, but this organism is found worldwide and especially where hygiene is poor. Other parasites that can be spread in food are the obligate parasitic eukaryotic microsporidia. Fresh produce such as berries, sprouts, and leafy greens can contain as high as 10^3 spores of *Enterocytozoon* spp. (60). These organisms probably come from contaminated irrigation water.

Carriers of bacterial or viral pathogens were originally infected after they were exposed to a pathogen source, such as during a foodborne or waterborne outbreak or through other infected persons such as convalescent patients and other carriers. From 1979 to 1995, at least 450,000 persons ≥ 20 years of age were hospitalized for gastroenteritis in the United States (93). This number excludes ill children, from whom siblings, parents, friends, caregivers, and teachers can acquire the infective agent and in turn become asymptomatic excretors.

As an indication of the likelihood of spread of pathogens by food workers, Clayton and Griffith (23) found that on average workers in the United Kingdom washed their hands adequately on only 9% of occasions after they had touched their faces or hair and on only 25% of those instances in which they touched potentially contaminated objects. In the United States, Green et al. (45) observed that workers made hand washing attempts (i.e., removed gloves if worn and placed hands in running water, used soap, and dried hands) after only 27% of activities such coughing, sneezing, working with raw animal products, or handling dirty equipment, utensils, or cloths. These data are in contrast to the confidence expressed by Pether and Scott (103), who considered convalescent *Salmonella* carriers a minimal risk for contaminating food. More recent data have indicated that worker hygiene is lax at times and that newer agents such as noroviruses have lower infectious doses than do salmonellae. These data give us reasons to consider the risks that occur in food preparation venues where there are opportunities for fecal and nasal contamination of hands on a regular basis and a lack of adequate hand washing. The next article in this series on food worker infections and

hygiene (122) will cover the transmission and survival of pathogens in the food processing and preparation environment.

ACKNOWLEDGMENT

The authors appreciate the critical review by Shirley B. Bohm (Consumer Safety Officer, Retail Food Protection, Center for Food Safety and Applied Nutrition, U.S. Food and Drug Administration, College Park, Md.), who offered helpful comments and new information to improve the manuscript.

REFERENCES

- Adesiyun, A. A., L. A. Webb, and H. T. Romain. 1997. Relatedness of *Staphylococcus aureus* strains isolated from milk and human handlers in dairy farms in Trinidad. *Zentbl. Vetmed. B* 44:551–560.
- Al bustan, M. A., E. E. Udo, and T. D. Chugh. 1996. Nasal carriage of enterotoxin-producing *Staphylococcus aureus* among restaurant workers in Kuwait City. *Epidemiol. Infect.* 116:319–322.
- Alvarez, M. L., F. Wurgaft, J. Espinoza, M. Araya, and G. Figuero. 1992. Hygiene habits and carriers in families with a child who has had typhoid fever. *Rev. Saude Publica* 26:75–81.
- Anderson, E. J., and S. G. Weber. 2004. Rotavirus infection in adults. *Lancet Infect. Dis.* 4:91–99.
- Bar-Dayan, Y., J. Shemer, and Y. Bar-Dayan. 1996. Food-borne and air-borne streptococcal pharyngitis—a clinical comparison. *Infection* 25:12–15.
- Barker, J., and M. V. Jones. 2005. The potential spread of infection caused by aerosol contamination of surfaces after flushing a domestic toilet. *J. Appl. Microbiol.* 99:339–347.
- Bidawid, S., N. Malik, O. Adegunrin, S. A. Sattar, and J. M. Farber. 2004. Norovirus cross-contamination during food handling and interruption of virus transfer by hand antiseptics: experiments with feline calicivirus as a surrogate. *J. Food Prot.* 67:103–109.
- Bishop, R. F. 1996. Natural history of human rotavirus infection. *Arch. Virol.* 12(Suppl.):119–128.
- Black, R. E., G. Lopez de Romana, K. H. Brown, N. Bravo, O. G. Bazalar, and H. C. Kanashiro. 1989. The incidence and etiology of infantile diarrhea and major routes of transmission in Huascar, Peru. *Am. J. Epidemiol.* 189:785–799.
- Borgatta, L., M. Fisher, and N. Robbins. 1989. Hand protection and protection from hands: hand-washing, germicides and gloves. *Women Health* 15:77–92.
- Braddick, M. R., B. J. Crump, and M. L. Yee. 1991. How long should patients with *Salmonella typhi* or *Salmonella paratyphi* be followed-up? A comparison of published guidelines. *J. Public Health Med.* 13:101–107.
- Brooks, G. F., and K. C. Carroll. 2004. Enteric gram-negative rods (*Enterobacteriaceae*), chap. 16. In G. F. Brooks, K. C. Carroll, J. S. Butel, and S. A. Morse (ed.), *Jawetz, Melnick, & Adelberg's medical microbiology*, 24th ed. Available at: <http://www.accessmedicine.com/content.aspx?aID=2758073>. Accessed 5 May 2008.
- Bryan, F. L. 1988. Risks of practices, procedures and processes that lead to outbreaks of foodborne diseases. *J. Food Prot.* 51:663–673.
- Caul, E. O. 1994. Small round structured viruses: airborne transmission and hospital control. *Lancet* 343:1240–1242.
- Centers for Disease Control and Prevention. 2001. “Norwalk-like viruses”: public health consequences and outbreak management. *Morb. Mortal. Wkly. Rep.* 50(RR09):1–18.
- Centers for Disease Control and Prevention. 2005. Giardiasis surveillance—United States, 1998–2002. In *Surveillance summaries*, January 28, 2005. *Morb. Mortal. Wkly. Rep.* 54(SS-1):9–16.
- Centers for Disease Control and Prevention. 2006. Multisite outbreak of norovirus associated with a franchise restaurant—Kent County, Michigan, May 2005. *Morb. Mortal. Wkly. Rep.* 55:395–397.
- Centers for Disease Control and Prevention. 2007. Multistate outbreak of norovirus gastroenteritis among attendees at a family re-

- union—Grant County, West Virginia, October 2006. *Morb. Mortal. Wkly. Rep.* 56:673–678.
19. Centers for Disease Control and Prevention. 2007. Norovirus outbreak associated with ill food-service workers—Michigan, January–February 2006. *Morb. Mortal. Wkly. Rep.* 56:1212–1216.
 20. Chan, M. C., J. J. Sung, R. K. Lam, P. K. Chan, N. L. Lee, R. W. Lai, and W. K. Leung. 2006. Fecal viral load and norovirus-associated gastroenteritis. *Emerg. Infect. Dis.* 12:1278–1280.
 21. Charles, R. H. G. 1985. Food handlers and food poisoning. *Health Hyg.* 6:2–7.
 22. Cheesebrough, J. S., J. Green, C. L. Gallimore, P. A. Wright, and D. W. Brown. 2000. Widespread environmental contamination with Norwalk-like viruses (NLV) detected in a prolonged hotel outbreak of gastroenteritis. *Epidemiol. Infect.* 125:93–98.
 23. Clayton, D., and C. Griffith. 2004. Observations of food safety practices in catering using notational analysis. *Br. Food J.* 106:211–227.
 24. Cotterill, H., A. Curry, and T. Riordan. 1988. Rotavirus in vomit. *J. Infect.* 16:206–207.
 25. de Wit, J. C., and E. H. Kampelmacher. 1981. Some aspects of microbial contamination of hands of workers in food industries. *Zentbl. Bakteriologie Mikrobiologie Hygiene B* 172:390–400.
 26. de Wit, J. C., and F. M. Rombouts. 1992. Faecal micro-organisms on the hands of carriers: *Escherichia coli* as model for Salmonella. *Zentbl. Hygiene Umweltmed.* 193:230–236.
 27. Do Carmo, L. S., C. Cummings, V. R. Linardi, R. S. Dias, J. M. De Souza, M. J. De Sena, D. A. Dos Santos, J. W. Shupp, R. K. P. Pereira, and M. Jett. 2004. A case study of a massive staphylococcal food poisoning incident. *Foodborne Pathog. Dis.* 1:241–246.
 28. The Doctor's Doctor. 2004. Impetigo. Available at: <http://www.thedoctorsdoctor.com/diseases/impetigo.htm>. Accessed 30 January 2008.
 29. Eady, E. A., and J. H. Cove. 2003. Skin and soft tissue infection. *Curr. Opin. Infect. Dis.* 16:103–124.
 30. Echeverria, P., D. N. Taylor, J. Seriwatana, U. Leksomboon, W. Chaicumpa, C. Tirapat, and B. Rowe. 1987. Potential sources of enterotoxigenic *Escherichia coli* in homes of children in Thailand. *Bull. W.H.O.* 65:207–215.
 31. Evans, M. R., R. Meldrum, W. Lane, D. Gardner, C. D. Ribeiro, C. I. Gallimore, and D. Westmoreland. 2002. An outbreak of viral gastroenteritis following environmental contamination at a concert hall. *Epidemiol. Infect.* 129:355–360.
 32. Expert Working Group. 1995. Food handlers, fitness to work, guidelines for food business managers. UK Department of Health. Available at: <http://www.tendringdc.gov.uk/NR/rdonlyres/23E8B031-6293-43B4-B916-2AD90630A0A1/0/FoodHandlersFitnessstoWork.pdf>. Accessed 4 April 2008.
 33. Farber, R. E., and F. A. Korff. 1958. Foodborne epidemic of group A beta hemolytic *Streptococcus*. *Public Health Rep.* 73:203–209.
 34. Farley, T. A., D. R. Johnson, E. L. Kaplan, K. Y. Kelso, F. Mahoney, and S. A. Wilson. 1993. Direct inoculation of food as the cause of group A streptococcal pharyngitis. *J. Infect. Dis.* 167:1232–1235.
 35. Feachem, R. G., D. J. Bradley, H. Garelick, and D. D. Mara. 1983. Sanitation and disease: health aspects of excreta and wastewater management. John Wiley & Sons, Chichester, UK.
 36. Fenton, P. A., K. W. Dobson, A. Eyre, and M. W. McKendrick. 1984. Unusually severe food poisoning from vanilla slices. *J. Hyg.* 93:377–388.
 37. Figueroa, G., P. Navarrete, M. Caro, M. Troncoso, and G. Faundez. 2002. [Carriage of enterotoxigenic *Staphylococcus aureus* in food handlers]. *Rev. Med. Chile* 130:859–864.
 38. Fisher, A. A. 1986. Contact dermatitis. Lea & Febiger, Philadelphia.
 39. Fisher, L. B., and H. I. Maibach. 1975. Effect of some irritants on human epidermal mitosis. *Contact Dermatitis* 1:273–276.
 40. Food Marketing Institute. 2004. Hepatitis A information guide. Food Marketing Institute, Washington, D.C. Available at: <http://www.fmi.org/foodsafety/HepatitisAGuide.pdf>. Accessed 4 April 2008.
 41. Gerba, C. 2000. Assessment of enteric pathogen shedding by bathers during recreational activity and its impact on water quality. *Quant. Microbiol.* 2:55–64.
 42. Gibson, L. L., J. B. Rose, and C. N. Haas. 1999. Use of quantitative microbial risk assessment for evaluation of the benefits of laundry sanitation. *Am. J. Infect. Control* 27:S34–S39.
 43. Goller, J. L., A. Dimitriadis, A. Tan, H. Kelly, and J. A. Marshall. 2004. Long-term features of norovirus gastroenteritis in the elderly. *J. Hosp. Infect.* 58:286–291.
 44. Graham, D. Y., X. Jiang, T. Tanaka, A. R. Opekun, H. P. Madore, and M. K. Estes. 1994. Norwalk virus infection of volunteers: new insights based on improved assays. *J. Infect. Dis.* 170:34–43.
 45. Green, L. R., C. A. Selman, V. Radke, D. Ripley, J. C. Mack, D. W. Reimann, T. Stigger, M. Motsinger, and L. Bushnell. 2006. Food worker hand washing practices: an observation study. *J. Food Prot.* 69:2417–2423.
 46. Greig, J. D., E. C. D. Todd, C. A. Bartleson, and B. Michaels. 2007. Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 1. Description of the problem, methods and agents involved. *J. Food Prot.* 70:1752–1761.
 47. Hamburger, M., M. J. Green, and G. V. Hamburger. 1945. Problem of “dangerous carrier” of hemolytic streptococci. 2. Spread of infection by individuals with strongly positive nose cultures who expelled large numbers of hemolytic streptococci. *J. Infect. Dis.* 77:96–108.
 48. Hamburger, M., Jr., and O. H. Robertson. 1948. Expulsion of group A hemolytic streptococci in droplet nuclei by sneezing, coughing, and talking. *Am. J. Med.* 4:690–701.
 49. Han, A. M., K. N. Do, T. Aye, and T. Hlaing. 1986. Personal toilet after defaecation and the degree of hand contamination according to different methods used. *J. Trop. Med. Hyg.* 89:237–241.
 50. Hatakka, M., K. J. Bjorkroth, K. Asplund, N. Maki-Petays, and H. J. Korkeala. 2000. Genotypes and enterotoxigenicity of *Staphylococcus aureus* isolated from the hands and nasal cavities of flight-catering employees. *J. Food Prot.* 63:1487–1491.
 51. Hedberg, C. W., and M. T. Osterholm. 1993. Outbreaks of foodborne and waterborne viral gastroenteritis. *Clin. Microbiol. Rev.* 6:199–210.
 52. Holmberg, S. D., and P. A. Blake. 1984. Staphylococcal food poisoning in the United States. New facts and old misconceptions. *JAMA* 251:487–489.
 53. Holmes, A., M. Ganner, S. McGuane, T. L. Pitt, B. D. Cookson, and A. M. Kearns. 2005. *Staphylococcus aureus* isolates carrying Panton-Valentine leucocidin genes in England and Wales: frequency, characterization, and association with clinical disease. *J. Clin. Microbiol.* 43:2384–2390.
 54. Hoque, B. A., J. Chakraborty, J. T. A. Chowdhury, U. K. Chowdhury, M. Ali, S. El Arifeen, and R. B. Sack. 1999. Effects of environmental factors on child survival in Bangladesh: a case control study. *Public Health* 113:57–64.
 55. Hoque, B. A., D. Mahalanabis, M. J. Alam, and M. S. Islam. 1995. Post-defecation handwashing in Bangladesh: practice and efficiency perspectives. *Public Health* 109:15–24.
 56. Horwood, M. P., and V. A. Minch. 1951. The numbers and types of bacteria found on the hands of food handlers. *Food Res.* 16:133–136.
 57. Hutchinson, R. I. 1956. Some observations on the method and spread of Sonne dysentery. *Mon. Bull. Minst. Health Public Health Lab. Serv.* 15:110–118.
 58. Ifesan, B. O., F. C. Adetuyi, A. K. Onifade, and O. O. Agarry. 2004. Microorganisms associated with human jewellery. *J. Food Agric. Environ.* 2:14–16.
 59. Jain, D., S. Sinha, K. N. Prasad, and C. M. Pandey. 2005. *Campylobacter* species and drug resistance in a north Indian rural community. *Trans. R. Soc. Trop. Med. Hyg.* 99:207–214.
 60. Jedrzejewski, S., T. K. Graczyk, A. Słodkiewicz-Kowalska, L. Tamang, and A. C. Majewska. 2007. Quantitative assessment of contamination of fresh food produce of various retail types by human virulent microsporidian spores. *Appl. Environ. Microbiol.* 73:4071–4073.
 61. Jones, T. F., M. E. Kellum, S. S. Porter, M. Bell, and W. Schaffner.

2002. An outbreak of community-acquired foodborne illness caused by methicillin-resistant *Staphylococcus aureus*. *Emerg. Infect. Dis.* 8:82–84.
62. Joseph, P. R., J. D. Millar, and D. A. Henderson. 1965. An outbreak of hepatitis traced to food contamination. *N. Engl. J. Med.* 273:188–194.
 63. Katzenell, U., J. Shemer, and Y. Bar-Dayana. 2001. Streptococcal contamination of food: an unusual cause of epidemic pharyngitis. *Epidemiol. Infect.* 127:179–184.
 64. Khan, A., K. A. Talukder, S. Huq, D. Mondal, M. A. Malek, D. K. Dutta, G. B. Nair, and A. S. Faruque. 2006. Detection of intra-familial transmission of *Shigella* infection using conventional serotyping and pulsed-field gel electrophoresis. *Epidemiol. Infect.* 134:605–611.
 65. Khuri-Bulos, N. A., M. A. Khalaf, A. Shenhabi, and K. Shami. 1994. Foodhandler-associated *Salmonella* outbreak in a university hospital despite routine surveillance cultures of kitchen employees. *Infect. Control Hosp. Epidemiol.* 15:311–314.
 66. Kligman, A. M. 1966. The SLS provocative patch test in allergic contact sensitization. *J. Invest. Dermatol.* 46:573–583.
 67. Kluytmans, J., W. van Leeuwen, W. Goessens, R. Hollis, S. Messer, L. Herwaldt, H. Bruining, M. Heck, J. Rost, N. van Leeuwen, A. van Belkum, and H. Verbrugh. 1995. Food-initiated outbreak of methicillin-resistant *Staphylococcus aureus* analyzed by pheno- and genotyping. *J. Clin. Microbiol.* 33:1121–1128.
 68. Knights, B., K. Redway, and V. Edwards. 1997. A study of hand washing habits in public toilets and the bacterial contamination of the hands before and after washing. Applied Ecology Research Group, School of Biosciences, University of Westminster. Available at: <http://www.wmin.ac.uk/~redwayk/research/toilet.htm>. Accessed 15 July 2008.
 69. Kohl, K. S., K. Rietberg, S. Wilson, and T. A. Farley. 2002. Relationship between home food-handling practices and sporadic salmonellosis in adults in Louisiana, United States. *Epidemiol. Infect.* 129:267–276.
 70. Konig, C., J. Blaser, and H. P. Simmen. 1998. Bacterial concentrators in pus and infected peritoneal fluid—implications for bactericidal activity of antibiotics. *J. Antimicrob. Chemother.* 42:227–232.
 71. Koning, S., A. van Belkum, S. Snijders, W. van Leeuwen, H. Verbrugh, J. Nouwen, M. Op't Veld, L. W. van Suijlekom-Smit, J. C. van der Wouden, and C. Verduin. 2003. Severity of nonbullous *Staphylococcus aureus* impetigo in children is associated with strains harboring genetic markers for exfoliative toxin B, Pantone-Valentine leukocidin, and the multidrug resistance plasmid pSK41. *J. Clin. Microbiol.* 41:3017–3021.
 72. Kure, R., H. Grendahl, and J. Paulssen. 1982. Pyrogens from surgeons' sterile latex gloves. *Acta Pathol. Microbiol. Immunol. Scand. B* 90:85–88.
 73. Kuritsky, J. N., M. T. Osterholm, H. B. Greenberg, J. A. Korlath, J. R. Godes, C. W. Hedberg, J. C. Forfang, A. Z. Kapikian, J. C. McCullough, and K. E. White. 1984. Norwalk gastroenteritis: a community outbreak associated with bakery product consumption. *Ann. Intern. Med.* 100:519–521.
 74. Lane, K. M., M. Scarborough, and M. Park. 2001. Outbreaks of foodborne disease in Georgia, 2000. *Ga. Epidemiol. Rep.* 17:1–3.
 75. Larson, E., J. J. Leyden, K. J. McGinley, G. L. Grove, and G. H. Talbot. 1986. Physiologic and microbiologic changes in skin related to frequent handwashing. *Infect. Control* 7:59–63.
 76. Lin, C. M., M. P. Doyle, B. S. Michaels, K. A. Williams, and F. M. Wu. 2003. A comparison of handwashing techniques to remove *Escherichia coli* and calicivirus under natural or artificial fingernails. *J. Food Prot.* 66:2296–2301.
 77. Lo, S. V., A. M. Connolly, D. Joynson, S. R. Palmer, P. D. Thomas, and D. Wright. 1994. The role of the pre-symptomatic food handler in a common source outbreak of food-borne SRSV gastroenteritis in a group of hospitals. *Epidemiol. Infect.* 113:513–521.
 78. Loeto, D., M. I. Matsheka, and B. A. Gashe. 2007. Enterotoxigenic and antibiotic resistance determination of *Staphylococcus aureus* strains isolated from food handlers in Gaborone, Botswana. *J. Food Prot.* 70:2764–2768.
 79. Lossos, I. S., I. Felsenstein, R. Breuer, and D. Engelhard. 1992. Food-borne outbreak of group A beta-hemolytic streptococcal pharyngitis. *Arch. Intern. Med.* 152:853–855.
 80. Lues, J. F. R., and I. Van Tonder. 2007. The occurrence of indicator bacteria on hands and aprons of food handlers in the delicatessen sections of a retail group. *Food Control* 18:326–332.
 81. Marks, P. J., I. B. Vipond, D. Carlisle, D. Deakin, R. Fey, and E. O. Caul. 2000. Evidence for airborne transmission of Norwalk-like virus (NLV) in a hotel restaurant. *Epidemiol. Infect.* 124:481–487.
 82. Mathias, C. G. 1990. Prevention of occupational contact dermatitis. *J. Am. Acad. Dermatol.* 23:742–748.
 83. Mathias, C. G. T., and J. H. Morrison. 1988. Occupational skin diseases, United States: results from the Bureau of Labor Statistics Annual Survey of Occupational Injuries and Illnesses, 1973 through 1984. *Arch. Dermatol.* 124:1519–1524.
 84. Mathur, R., and V. Reddy. 1983. Bacterial contamination of infant foods. *Indian J. Med. Res.* 77:342–346.
 85. Medus, C., K. E. Smith, J. B. Bender, J. M. Besser and C. W. Hedberg. 2006. *Salmonella* outbreaks in restaurants in Minnesota, 1995 through 2003: evaluation of the role of infected foodworkers. *J. Food Prot.* 69:1870–1878.
 86. Meehan, P. J., T. Atkeson, D. E. Kepner, and M. Melton. 1992. A foodborne outbreak of gastroenteritis involving two different pathogens. *Am. J. Epidemiol.* 136:611–616.
 87. Mensah, P., D. Yeboah-Manu, K. Owusu-Darko, and A. Ablordey. 2002. Street foods in Accra, Ghana: how safe are they? *Bull. W.H.O.* 80:546–554.
 88. Michaels, B., and T. Ayers. 2000. Hazard analysis of the personal hygiene process, p. 191–200. In Proceedings of the 2nd National Sanitation Foundation International Conference on Food Safety, Savannah, Ga.
 89. Michaels, B., M. Celis, T. Ayers, and V. Gangar. 2001. A microbial survey of toilet paper and associated performance variables related to its role in reducing communicable disease transmission. *J. Food Prot.* 64(Suppl. A):92.
 90. Mills, S. J. C., D. J. Holland, and A. E. Hardy. 2000. Operative field contamination by the sweating surgeon. *Aust. N.Z. J. Surg.* 70:837–839.
 91. Minor, T. E., and E. H. Marth. 1976. Staphylococci and their significance in food. Elsevier Scientific Publishing Company, Amsterdam.
 92. Morris, A. J., P. R. Murray, and L. B. Reller. 1996. Contemporary testing for enteric pathogens: the potential for cost, time, and health care savings. *J. Clin. Microbiol.* 34:1776–1778.
 93. Mounts, A. W., R. C. Holman, M. J. Clarke, J. S. Bresee, and R. I. Glass. 1999. Trends in hospitalizations associated with gastroenteritis among adults in the United States, 1979–1995. *Epidemiol. Infect.* 123:1–8.
 94. Murakami, K., T. Ishihara, K. Horikawa, and T. Oda. 2007. Features of *Salmonella* serovars among food handlers in Kyushu, Japan. *New Microbiol.* 30:155–159.
 95. Nainan, O. V., G. Xia, G. Vaughan, and H. S. Margolis. 2006. Diagnosis of hepatitis A virus infection: a molecular approach. *Clin. Microbiol. Rev.* 19:63–79.
 96. Nilsson, E., C. Henning, and M. L. Hjörleifsson. 1986. Density of the microflora in hand eczema before and after topical treatment with a potent corticosteroid. *J. Am. Acad. Dermatol.* 15:192–197.
 97. Noda, M., S. Fukuda, and O. Nishio. 2008. Statistical analysis of attack rate in norovirus foodborne outbreaks. *Int. J. Food Microbiol.* 122:216–220.
 98. Ozawa, K., T. Oka, N. Takeda, and G. S. Hansman. 2007. Norovirus infections in symptomatic and asymptomatic food handlers in Japan. *J. Clin. Microbiol.* 45:3996–4005.
 99. Parashar, U. D., L. Dow, R. L. Fankhauser, C. D. Humphrey, J. Miller, T. Ando, K. S. Williams, C. R. Eddy, J. S. Noel, T. Ingram, J. S. Bresee, S. S. Monroe, and R. I. Glass. 1998. An outbreak of viral gastroenteritis associated with consumption of sandwiches: implications for the control of transmission by food handlers. *Epidemiol. Infect.* 121:615–621.
 100. Patrick, D. R., G. Findon, and T. E. Miller. 1997. Residual moisture

- determines the level of touch-contact-associated bacterial transfer following hand washing. *Epidemiol. Infect.* 119:19–25.
101. Patterson, W., P. T. Fryers, J. Green, and P. Haswell. 1997. Outbreak of small round structured virus gastroenteritis arose after kitchen assistant vomited. *Commun. Dis. Rep.* 27:R101–R103.
 102. Pereira, M. L., L. S. do Carmo, E. J. dos Santos, and M. S. Bergdoll. 1994. Staphylococcal food poisoning from cream-filled cake in a metropolitan area of south-eastern Brazil. *Rev. Saude Publica* 28:406–409.
 103. Pether, J. V. S., and R. J. D. Scott. 1983. *Salmonella* carriers; are they dangerous? A study to identify finger contamination with salmonellae by convalescent carriers. *J. Infect.* 5:81–88.
 104. Polledo, J. J. F., M. L. Garcia, B. Moreno, and I. Menes. 1985. Importance of food handlers as a source of enterotoxigenic. *Zentbl. Bakteriol. Mikrobiol. Hyg. B* 181:364–373.
 105. Ramsing, D. W., and T. Agner. 1996. Effect of glove occlusion on human skin. (I). Short-term experimental exposure. *Contact Dermatitis* 34:1–5.
 106. Ramsing, D. W., and T. Agner. 1996. Effect of glove occlusion on human skin (II). Long-term experimental exposure. *Contact Dermatitis* 34:258–262.
 107. Rayner, M., P. Nichols, D. Nevins, and D. C. White. 1996. Rapid method for extraction of the fecal sterol coprostanol using supercritical fluid carbon dioxide. *Div. Environ. Chem. Preprints Ext. Abstr.* 36:273–276.
 108. Reali, D. 1982. Enterotoxin A and B production in strains of *Staphylococcus aureus* isolated from human beings and foods. *J. Hyg. (Lond.)* 88:103–106.
 109. Reina, J., F. Salva, and P. Alomar. 1989. Enterotoxin production by strains of *Staphylococcus aureus* isolated from clinical specimens. *J. Clin. Pathol.* 42:217–218.
 110. Rockx, B., M. de Wit, H. Vennema, J. Vinjé, E. de Bruin, Y. van Duynhoven, and M. Koopmans. 2002. Natural history of human *Calicivirus* infection: a prospective cohort study. *Clin. Infect. Dis.* 35:246–253.
 111. Rohner, P., D. Pittet, B. Pepey, T. Nije-Kinge, and R. Auckenthaler. 1997. Etiological agents of infectious diarrhea: implications for requests for microbial culture. *J. Clin. Microbiol.* 35:1427–1432.
 112. Sattar, S. A., N. Lloyd-Evans, V. S. Springthorpe, and R. C. Nair. 1986. Institutional outbreaks of rotavirus diarrhoea: potential role of fomites and environmental surfaces as vehicles for virus transmission. *J. Hyg. (Lond.)* 96:277–289.
 113. Sattar, S. A., V. S. Springthorpe, J. Tetro, R. Vashon, and B. Keswick. 2002. Hygienic hand antiseptics: should they not have activity and label claims against viruses? *Am. J. Infect. Control* 30:355–372.
 114. Sawyer, L. A., J. J. Murphy, J. E. Kaplan, P. F. Pinsky, D. Chacon, S. Walmsley, L. B. Schonberger, A. Phillips, K. Forward, C. Goldman, J. Brunton, R. A. Fralick, A. O. Carter, W. G. Gary, Jr., R. I. Glass, and D. E. Low. 1988. 25- to 30-nm virus particle associated with a hospital outbreak of acute gastroenteritis with evidence for airborne transmission. *Am. J. Epidemiol.* 127:1261–1271.
 115. Siegel, D. L., P. H. Edelstein, and I. Nachamkin. 1990. Inappropriate testing for diarrheal diseases in the hospital. *JAMA* 263:979–982.
 116. Snyder, O. P. 2008. A “safe hands” hand wash program for retail food operations. Hospitality Institute of Technology and Management, St. Paul, Minn. Available at: <http://www.hi-tm.com/Documents/Safehands.html>. Accessed 25 February 2008.
 117. Stryker, W. S., R. R. Facklam, and D. W. Fraser. 1982. Foodborne outbreak of group G streptococcal pharyngitis. *Am. J. Epidemiol.* 116:533–540.
 118. Todd, E. C. D., and P. Chapman. 1996. Foodborne and waterborne disease in Canada. Annual summaries: 1988 and 1989. Health Protection Branch, Health Canada, Polyscience Publications, Morin Heights, Quebec, Canada.
 119. Todd, E. C. D., J. D. Greig, C. A. Bartleson, and B. S. Michaels. 2007. Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 2. Description of outbreaks by size, severity, and settings. *J. Food Prot.* 70:1975–1993.
 120. Todd, E. C. D., J. D. Greig, C. A. Bartleson, and B. S. Michaels. 2007. Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 3. Factors contributing to outbreaks and description of outbreak categories. *J. Food Prot.* 70:2199–2217.
 121. Todd, E. C. D., J. D. Greig, C. A. Bartleson, and B. S. Michaels. 2008. Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 4. Infective doses and pathogen carriage. *J. Food Prot.* 71:2339–2373.
 122. Todd, E. C. D., J. D. Greig, C. A. Bartleson, and B. S. Michaels. Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 6. Transmission and survival of pathogens in the food processing and preparation environment. *J. Food Prot.*, in press.
 123. Tompkins, D. S., M. J. Hudson, H. R. Smith, R. P. Eglin, J. G. Wheeler, M. M. Brett, R. J. Owen, J. S. Brazier, P. Cumberland, V. King, and P. E. Cook. 1999. A study of infectious intestinal disease in England: microbiological findings in cases and controls. *Commun. Dis. Public Health* 2:108–113. [Erratum in *Commun. Dis. Public Health* 1999, 2:222.]
 124. U.S. Food and Drug Administration. 2005. FDA Food Code and 2007 Supplement to the Food Code. Available at: <http://www.cfsan.fda.gov/~dms/foodcode.html>. Accessed 28 April 2008.
 125. van der Meeren, H. L., and P. E. van Erp. 1986. Life-threatening contact urticaria from glove powder. *Contact Dermatitis* 14:190–191.
 126. van Schothorst, M., and H. J. Beckers. 1978. Persistent excretion of salmonellas. *Br. Med. J.* 2:1301.
 127. Warburton, A. R. E., R. Buttery, J. V. Parry, K. R. Perry, S. Rampling, K. N. Ward, and T. G. Wreghitt. 1991. Hepatitis A outbreak involving bread. *Epidemiol. Infect.* 106:199–202.
 128. Ward, R. L., D. I. Bernstein, D. R. Knowlton, J. R. Sherwood, E. C. Young, T. M. Cusack, J. R. Rubino, and G. M. Schiff. 1991. Prevention of surface-to-human transmission of rotaviruses by treatment with disinfectant spray. *J. Clin. Microbiol.* 29:1991–1996.
 129. Ward, R. L., D. R. Knowlton, and M. J. Pierce. 1984. Efficiency of rotavirus propagation in cell culture. *J. Clin. Microbiol.* 19:748–753.
 130. Wei, H. L., and C. S. Chiou. 2002. Molecular subtyping of *Staphylococcus aureus* from an outbreak associated with a food handler. *Epidemiol. Infect.* 128:15–20.
 131. White, N. 1990. Endotoxins on surgical gloves. *J. Hosp. Infect.* 16:390.
 132. Wieneke, A. A., D. Roberts, and R. J. Gilbert. 1993. Staphylococcal food poisoning in the United Kingdom, 1969–90. *Epidemiol. Infect.* 110:519–531.
 133. Williams, P. B., and J. F. Halsey. 1997. Endotoxin as a factor in adverse reactions to latex gloves. *Ann. Allergy Asthma Immunol.* 79:303–310.
 134. Winter, J. 1983. Certification of food handlers: a system to fulfill EEC requirements. *J. Soc. Occup. Med.* 33:93–97.
 135. World Health Organization. 1976. Foodborne *Salmonella* infections contracted on aircraft. *W.H.O. Wkly. Epidemiol. Rec.* 51:265–266.
 136. Zuger, C. 1981. Allergy to zinc dimethyldithiocarbamate in rubber gloves. *Contact Dermatitis* 7:337–338.

Journal of Food Protection Indices to Volume 71

AUTHOR INDEX

- ABDULMAWJOOD, A., see Schöenbrücher, 2059
 ABECIA, E. C. G., see Boesche Tomazelli, 2442
 ABEE, T., see Van Boeijen, 2007
 ABOLLO, E., see Pascual, 2316
 ABONG'O, B. O., Prevalence of *E. coli* O157:H7 in vegetables, 816
 ABOU-TALEB, M., Shelf life of coated and semifried tuna, 770
 ABRIOUEL, H., see Cobo Molinos, 2460
 ABU AL-HASAN, A. S., see Osaili, 934
 ABU BAKAR, F., see Chai, 2448
 ACHESON, D. W. K., see Klontz, 1277
 ACUFF, G. R., see Niebuhr, 714; Sawyer, 405
 ADAMS, K. D., see Miller, 2078
 AILES, E. C., Factors for produce contamination, 2389
 AKBAS, M. Y., see Emer, 914
 AKDEMIR EVRENDILEK, G., High-pressure processing of Turkish white cheese, 102
 AKINS, E. D., Microflora on cantaloupes, 46
 AL-ADHAMI, B. H., SSCP analysis for rapid discrimination of *Salmonella* isolates, 1960
 ALBILLOS, S., see Jackson, 445
 ALBRECHT, J. A., see Yarbava, 77
 ALLAIN, R., see Blais, 2301
 ALLEN, V. M., *Campylobacter* in thinning of broiler flocks, 264
 ALLENDE, A., Impact of water reuse in fresh-cut washing, 2514
 ALLENDE, A., see Selma, 286
 ALMOND, G. W., see Mason, 584
 AL-NABULSI, A. A., see Osaili, 934
 ALONSO, E., see Chicón, 1453
 AL-TAHER, F. M., see Jackson, 445
 ALTHAUS, R., see Zorraquino, 1193
 ALTIERI, C., see Bevilacqua, 1366; Cibelli, 2129
 ALVARADO, C. Z., see Gailunas, 629
 ALVARADO, M., see Brooks, 293
 ALVAREZ-OJEDA, G., see Gallegos-Robles, 2217
 AMERATUNGA, R., Latex glove use by food handlers, 2334
 AMERATUNGA, S., see Ameratunga, 2334
 AMÉZQUETA, S., OTA removal procedure for cocoa shells, 1422
 AMINLARI, M., see Amiri, 411
 AMIRI, S., Antibacterial activity of dextran-conjugated lysozyme in cheese curd, 411
 AMMOR, M. S., Role of quorum sensing in food spoilage, 1510
 ANADÓN, A., Acute toxicity of rosemary extracts, 790
 ANANTHESWARAN, R. C., see Hayman, 426
 ANDERSON, N., see Keller, 775
 ANDERSON, R. C., see Edrington, 27
 ANDO, M., Mercury in bluefin tuna, 595
 ANDREWS, K., see Crippen, 1981
 ANGELINI, E., OTA determinations in grapes, 2488
 ANGIIONI, A., see Schirra, 967
 AOUNI, R., see Beneduce, 2082
 AOYAMA, K., see Sakai, 1855
 APPEL, B., see Scherer, 699
 ARAGÃO, G. M. F., see Salomão, 643
 ARAGON-ALEGRO, L. C., *B. cereus*: enterotoxigenic and genetic profile, 2115
 ARES, I., see Anadón, 790
 ARIAS, C. R., see Limthammahisorn, 157
 ARLORIO, M., see Bonetta, Silvia, 205
 ARNOLD, G., see Pointon, 1123
 ARNOLD, J. W., see Sanders, 271
 ARROYO-LÓPEZ, F. N., see Bautista-Gallego, 1412
 ARTHUR, T. M., *E. coli* O157:H7 and *Salmonella* in lairages, 1752; Effects of interventions on MDR *Salmonella*, 2177; *Salmonella* prevalence in bovine lymph nodes, 1685
 ARTHUR, T. M., see Kalchayanand, 621
 ARUSCAVAGE, D., Survival of *E. coli* O157 on lettuce, 2384
 ARVANITTOYANNIS, I., see Hadjichristodoulou, 1632
 ASAI, Y., Incidence of *Salmonella* in imported seafood, 1460
 ASAO, T., see Sakai, 1855
 ASSÉRÉ, A., see Lagendijk, 1898
 ATANASSOVA, V., *Arcobacter* spp. in poultry processing, 2533; Microbiological quality of sushi, 860
 AUTIO, W. R., see Rodriguez, 170
 ÁVILA, M., see Morales, 811
 AWAKOWICZ, P., see Deilmann, 2119
 AYYASH, M. M., see Osaili, 934
 AZCARATE, M. P., *Alternaria* toxins in wheat, 1262
 AZEREDO, J., see Silva, 1379
 AZNAR, R., see Elizaguivel, 2110
 BAALHAM, T., see Williams, 835
 BACCARIN, A., see Tomazelli, 2442
 BACH, S. J., see Niu, 691
 BACKER, L. C., see Ailes, 2389
 BAERT, L., Survival of murine norovirus 1 on onions and spinach, 1590
 BAHK, G.-J., see Hong, 1248
 BAKER-AUSTIN, C., Antibiotic resistance in *V. parahaemolyticus*, 2552
 BALASUBRAMANIAM, V. M., see Akdemir Evrendilek, 102
 BALDAUF, N. A., see Männig, 2249
 BALTRUWEIT, I., see Röder, 2263
 BANG, J., see Kim, 940
 BANG, W., Detection of *S. aureus*, 191
 BARBUTI, S., see Ravyts, 1817
 BARI, M. L., Hot water treatment decontamination of mung bean seeds, 830
 BARIONI JÚNIOR, W., see Tomazelli, 2442
 BARMPALIA-DAVIS, I. M., Survival of *L. monocytogenes* in a gastrointestinal model, 2014
 BARNETT, J. D., see Yancy, 210
 BARRETT, J. B., see McGowan-Spicer, 760
 BARRETT, N. J., see Little, 309
 BARRY-RYAN, C., see Gutierrez, 1846
 BARTLESON, C. A., see Todd, 2339, 2582
 BATTLE, R., see López, 1889
 BAUER, M. L., see Tabe, 539
 BAUERMEISTER, L. J., Peracetic acid mixture as an antimicrobial treatment in poultry, 1119
 BAUMGARTNER, A., see Templer, 2100
 BAUTISTA-GALLEGO, J., Effects of chloride salts on microbial growth, 1412
 BAYNES, R. E., see Mason, 584
 BAZZO, I., see Angelini, 2488
 BEAUFORT, A., see Midelet-Bourdin, 2198
 BEDIN, J. M., see Tomazelli, 2442
 BEDNAR, C., see Kwon, 1651
 BEDNARCZYK, R., see Nelson, 365
 BEER, M., see Rhoades, 2272
 BEHNKE, A., see Kuhn, 993
 BELESSI, C.-I. A., Survival of *L. innocua* in dairy products, 742
 BELK, K. E., see Bowling, 83; Byelashov, 728; Carlson, 1343, 2223; Dewell, 573, 1114, 2228; Mukherjee, 1349
 BELLETTI, N., see Sado Kamdem, 2572; Torriani, 93

- BELLINGER, G. R., see Carlson, 2223
 BELLOQUE, J., see Chicón, 1453
 BENDER, J. B., see Smith, 2153
 BENEDUCE, L., Occurrence of STEC in meat products in Morocco, 2082
 BENEDESI, A., see Di Cagno, 1491
 BEN OMAR, N., see Cobo Molinos, 2460
 BERA, F., see Yao, 431
 BERENDS, B. R., see Wijnker, 1199
 BERGIN, D., see Gonzales Barron, 1330
 BERRANG, M. E., *L. monocytogenes* in drain pipes, 66
 BERRY, E. D., Determining *E. coli* O157:H7 levels in manure and soil, 2233
 BERTOLINI, P., see Gregori, 1626
 BESSER, T. E., see Cobbold, 1023
 BEUCHAT, L. R., *Salmonella* in tomatoes, 1572
 BEUCHAT, L. R., see Kim, 940; Vojdani, 356
 BEUMER, R. R., see Verhoeff-Bakkenes, 1018
 BEVILACQUA, A., Metabiotic effects of *Fusarium*, 1366
 BEVILACQUA, A., see Cibelli, 1219
 BHADURI, S., see Wesley, 1162
 BIANCHINI, A., see Voss, 2036
 BIBINOV, N., see Deilmann, 2119
 BIDAWID, S., see Pagotto, 1434
 BIGLER, L., see Miescher Schwenninger, 2481
 BJÖRKROTH, J., see Tolvanen, 1550
 BJÖRKROTH, K. J., see Vihavainen, 2312
 BJORNSDOTTIR, K., see Kreske, 2404
 BLACK, D. G., Efficacy of household products against *Bacillus* spores, 473
 BLACK, D. G., see Zhong, 1015
 BLAIR, I. S., see Giotis, 1481
 BLAIS, B. W., Detection of *Salmonella* Enteritidis by polymixin-ELISA, 392; Distinguishable control strains for food microbiology testing, 2301
 BLAIS, B. W., see Al-Adhami, 1960
 BLANA, V. A., see Paramithiotis, 1244
 BLANCO, D., see Venturini, 1701
 BLANDING, S., see Ailes, 2389
 BOA, A. C., see Chrystal, 2526
 BOARDMAN, G. D., see Sheldon, 345
 BOLLAERTS, K., see Delhalle, 1320
 BOLLINGER, L. M., see Hussein, 927
 BOLTON, F. J., see Little, 19
 BONETTA, SARA, see Bonetta, Silvia, 205
 BONETTA, SILVIA, Decarboxylating strains in a typical Italian cheese, 205
 BONGAERTS, R. J., see Crippen, 1981
 BONO, J. L., see Arthur, 2177
 BOOR, K. J., see Chan, 420
 BOOREN, A. M., see Tuntivanich, 2548; Warsaw, 153
 BORGER, A. C., see Seman, 1386
 BORGETTI, C., see Paulsen, 376
 BORGO, M., see Angelini, 2488
 BORSA, J., see Turgis, 516, 1237
 BORSETTI, F., see Gregori, 1626
 BOSILEVAC, J. M., Effects of reduced enrichment volumes, 1768
 BOSILEVAC, J. M., see Arthur, 1685, 1752, 2177; Kalchayanand, 621
 BOURKE, P., see Gutierrez, 1846
 BOVER-CID, S., see Torriani, 93
 BOWEN, B. M., see Chan, 420
 BOWERS, J. W. J., see Bauermeister, 1119
 BOWLING, M. B., Cross-contamination of CNS tissue due to beef carcass splitting, 83
 BOWLING, R. A., see Carlson, 2223
 BOXRUD, D. J., see Smith, 2153
 BOYER, R. R., see Gailunas, 629
 BRACKETT, R. E., see Klontz, 1277
 BRADY, Y. J., see Limthammahisorn, 157
 BRANDL, M. T., Decontamination of almonds with infrared heat, 897
 BRANDL, M. T., see Danyluk, 397
 BRASHEARS, M. M., see Brooks, 293; Miller, 2078
 BRAYMEN, C., see Smith, 2153
 BREIDT, F., JR., see Kreske, 2404
 BRICHTA-HARHAY, D. M., see Arthur, 1685, 1752, 2177; Kalchayanand, 621
 BROOKS, J. C., Spoilage and safety characteristics of ground beef, 293
 BROOKS, J. C., see Miller, 2078
 BROWN, C. A., Antimicrobial activity of lactoferrin, 319
 BROWN, E. W., see Xi, 2067
 BROWN, J. C., Antibiotic-resistant bacteria in herbal products, 1486
 BROŽKOVÁ, I., see Pejchalová, 719
 BULLERMAN, L. B., see Hassan, 2213; Obritsch, 1401; Voss, 2036
 BÜLTE, M., see Schönenbrücher, 2059
 BUONOCORE, G. G., see Conte, 119
 BURILLO, J., see Carramiñana, 502
 BUSH, E., see Wesley, 1162
 BUTLER, F., see Gonzales Barron, 1330
 BUUR, J. L., see Mason, 584
 BYELASHOV, O. A., Control of *L. monocytogenes* on frankfurters, 728
 CABEDO, L., *Salmonella* and *L. monocytogenes* in ready-to-eat food, 855
 CABEZA, M. C., see Hoz, 2001
 CABRAS, P., see Schirra, 967
 CABRERA-DIAZ, E., see Neal, 2415; Sawyer, 405
 CALL, J. E., see Luchansky, 2190; Orozco R., 676; Porto-Fett, 918
 CALLAWAY, T. R., see Edrington, 27
 CALO-MATA, P., see Miranda, 2537
 CALZADA, J., see Morales, 811
 CAMA, V., see Ortega, 2410
 CAMBERO, M. I., see Hoz, 2001
 CANNESON, P., see Shabala, 1169
 CARDILLO, D., see Bevilacqua, 1366
 CARDINAL, M., see Midelet-Bourdin, 2198
 CARLSON, B. A., Reduction of hide microbial contamination, 2223; Reduction of microbial contamination on cattle hides, 1343
 CARPENTIER, B., see Lagendijk, 1898
 CARRAMIÑANA, J. J., Antimicrobial essential oils against *L. monocytogenes* in pork, 502
 CARRARO, E., see Bonetta, Silvia, 205
 CARRASCOSA, A. V., see Nunez, 1496
 CARRILLO, C. D., see Medeiros, 2087
 CARROLL, T., see Robertson, 1186
 CARTER, J. M., see He, 2053
 CARVER, D. K., see Wright, 1791
 CASADO, E. M., see Sánchez, 2497
 CASON, J. A., see Hinton, 1707
 CASSONE, A., see Di Cagno, 1491
 CASTELL, V., see Martí-Cid, 2148
 CASTILLO, A., see Neal, 2415
 CATIGNANI, G. L., see Clare, 126
 CENTRELLA, B., see Carlson, 2223
 CEPEDA, A., see Miranda, 2537
 CEPEDA, M., see Soni, 134
 CERVENKA, L., Susceptibility of *Arcobacter* spp. to aromatic compounds, 165
 CHAI, L.-C., Contamination of kitchen by *C. jejuni* in raw vegetables, 2448
 CHAN, C.-H., see Little, 19
 CHAN, Y. C., *L. monocytogenes* growth at low temperature, 420
 CHANG, Y.-C., Multiplex PCR detection of enterotoxin genes in *Aeromonas* spp., 2094
 CHAPLEAU, N., see Ritz, 1612
 CHAPMAN, A., see Pointon, 1123
 CHATELLIER, S., see Healy, 1372
 CHATURONGAKUL, S., see Chan, 420
 CHEKROUN, A., see Kaddouri, 1270
 CHEMALY, M., *L. monocytogenes* in poultry flocks, 1996
 CHEN, F.-C., see Kilonzo-Nthenge, 608
 CHEN, G., see Medeiros, 1666
 CHEN, G.-H., Expression of goat lactoferricin from *E. coli*, 2523
 CHEN, H., see Kingsley, 1598

- CHEN, J., *Salmonella* Enteritidis in eggs, 2073
 CHEN, Y., see Lomonaco, 2559
 CHENG, C.-M., Detection of *Salmonella* by PCR in foods, 2436
 CHENG, C.-P., see Chuang, 1108
 CHENG, L. W., see He, 2053
 CHEW-M., Y., see Gallegos-Robles, 2217
 CHIA, T. W. R., Physicochemical properties of *Salmonella*, 2421
 CHIANG, I.-H., see Chen, 2523
 CHIANG, M.-L., Protein expression in stressed *V. parahaemolyticus*, 2589
 CHICÓN, R., Hydrolysates of β -lactoglobulin with low IgE binding, 1453
 CHILTON, N. B., see Al-Adhami, 1960
 CHINA, B., see Ghafir, 35
 CHIRTEL, S., see Fu, 888; Keller, 775
 CHO, A.-Y., Enzyme immunoassay for detection for amygdalin, 2048
 CHOI, J.-G., see Shim, 781
 CHOISEUL, J., see Lai, 578
 CHOU, C.-C., see Chiang, 2589
 CHRYSTAL, N. D., *Campylobacter* on retail broiler carcasses in New Zealand, 2526
 CHUANG, Y.-C., PCR detection of *Salmonella* by *stdA* primers, 1108
 CHUNG, D.-H., see Shim, 781
 CHUNG, J., see Cho, 2048
 CHUREY, J. J., see Salomão, 643
 CIBELLI, F., Proteolytic of molds, 1366
 CIBELLI, F., see Bevilacqua, 1366
 CICCARONE, C., see Cibelli, 2129
 CIFUENTES, A., see Mendiola, 2138
 CINER, F. L., see Zuckschwerdt, 1007
 CLARE, D. A., Antibacterial milkfat globule membrane fractions, 126
 CLAYTON, H. A., see Ailes, 2389
 CLOGHER, P., see Nelson, 365
 COBBOLD, R. N., Non-O157 STEC from humans, cattle, and food in the United States, 1023
 COBO MOLINOS, A., *L. monocytogenes* inactivation by AS-48, 2460
 COISSON, J. D., see Bonetta, Silvia, 205
 COLLIN, V., see Healy, 1372
 COMBS, B., see Pointon, 1123
 CONESA, M. A., see Selma, 286
 CONTE, A., Active packaging, 119
 COOKE, P., see Ukuku, 684
 COPIN, S., see Midelet-Bourdin, 2198
 CÓRDOBA, J. J., see Sánchez, 2497
 CORNEAU, N., see Pagotto, 1434
 COX, M., see Wesley, 545
 CRIPPEN, T. L., Fourteen-day-old chicken cecal microflora, 1981
 CRITZER, F. J., Transcription of *stx*₁, *marA*, and *eaeA* in *E. coli* O157:H7, 1469
 CROLEY, T. R., see Zuckschwerdt, 1007
 CRONIN, U. P., Physiological response of *B. cereus* to food processing treatments, 2168
 CROOKS, C., see Ameratunga, 2334
 CRUZ, A. G., see Nunes Filho, 1253
 CRUZ, J., Thermal and hydrostatic inactivation, 196
 CUI, S., see Koitabashi, 706
 CUTLER, S. A., see Patton, 2519

 D'AGOSTINO, R., see Conte, 119
 D'AMICO, D. J., *Listeria* spp. in farmstead cheese processing environments, 2239; Pathogen incidence in farmstead raw milk, 1580; Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
 D'ANGELIS, C. E. M., see Minei, 634
 DANIELS, A., see Nelson, 365
 DANIELSEN, M., see Huys, 339
 DANILA, R. N., see Smith, 2153
 DANYLUK, M. D., *Salmonella* migration into almonds, 397
 DANZEISEN, G. T., see Thorsness, 2295
 D'AQUINO, S., see Schirra, 967
 DARGATZ, D. A., see Dewell, 1114, 2228
 DA SILVA, C. M., see Tomazelli, 2442
 DATTA, A. R., see Klontz, 1277
 DAUBE, G., see Delhalle, 1320; Ghafir, 35
 DAVIDSON, P. M., see Black, 473; Zhong, 1015
 DAVIS, M. A., see Cobbold, 1023
 DAVOS, D., see Pointon, 1123
 DAY, J. B., Thermal resistance of *F. tularensis*, 2208
 DAYRE, M., see Thouand, 380
 DE ANGELIS, M., see Di Cagno, 1491
 DEBEVERE, J., see Baert, 1590
 DECASTELLI, L., see Lai, 578
 DE FREITAS, J. B., see Tomazelli, 2442
 DEILMANN, M., New sterilization process for aseptic packaging, 2119
 DE JESÚS, A. J., CO₂ influence on *L. monocytogenes* growth, 1915
 DE JONG, A. E. I., see Verhoeff-Bakkenes, 1018
 DE JONGE, R., see Verhoeff-Bakkenes, 1018
 DE LAMBALLERIE, M., see Ritz, 1612
 DE LAS HERAS, C., see Rodríguez-Mahillo, 2578; Solas, 1273
 DE LAS RIVAS, B., *M. morgani* ornithine decarboxylase, 657
 DELHALLE, L., Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
 DEL NOBILE, M. A., see Conte, 119
 DEL OLMO, A., Bactericidal effect of lactoferrin and its derivatives, 2468
 DE MARTINIS, E. C. P., see Minei, 634
 DE MARTINO, M., see Wesley, 545
 DEMİRBAŞ, N., Food safety for the Turkish dairy industry, 440
 DE PINNA, E., see Little, 19
 DERENS, E., see Legendijk, 1898
 DE SADELEER, L., see Delhalle, 1320
 DESTRO, M. T., see Aragon-Alegro, 2115
 DEVRIES, J. W., see Jackson, 445
 DE VUYST, L., see Ravyts, 1817
 DEWELL, G. A., Effect of transportation and lairage on risk of *S. enterica* in cattle, 2228; Impact of transportation and lairage on *E. coli* O157:H7 in cattle, 1114
 DEWELL, R. D., Potentially prohibited tissue removal, 573
 DEWELL, R. D., see Dewell, 1114, 2228
 DEWULF, J., see Delhalle, 1320
 DE ZUTTER, L., see Delhalle, 1320; Ghafir, 35; Rasschaert, 146
 D'HAENE, K., see Huys, 339
 DIARRA, M. S., see Lefebvre, 1774
 DI CAGNO, R., *Lactobacillus* for gluten and phytate hydrolysis in gluten-free bread, 1491
 DICKSON, J. S., see Niebuhr, 714; Patton, 2519
 DIERICK, K., see Ghafir, 35
 DIEZ-GONZALEZ, F., see Pal, 1806; Xu, 333
 DOETKOTT, C., see Nde, 386; Thorsness, 2295
 DOETKOTT, D., see Khaita, 1679
 DOETKOTT, D. K., see Tabe, 539
 DOMINGO, J. L., see Martí-Cid, 2148
 DOMINGUEZ, S. A., Modeling the growth of *Salmonella* in poultry, 2429
 DONNELLY, C. W., see D'Amico, 1563, 1580, 2239
 DONOGHUE, D. J., see Reyes-Herrera, 223
 DOORN, J., see Lenhart, 70
 DOS SANTOS, A., see Tomazelli, 2442
 DOSTALÍKOVÁ, E., see Pejchalová, 719
 DOWD, S. E., Toxin expression in O157 lineages, 1673
 DOWD, S. E., see Crippen, 1981
 DOWSETT, P., see Pointon, 1123
 DRAKE, M. A., see Bang, 191
 DROSINOS, E. H., see Belessi, 742; Paramithiotis, 1244
 DRUART, M. J., see D'Amico, 1563
 D'SOUZA, D. H., see Critzer, 1469
 DUARTE, D. A. M., see Tomazelli, 2442
 DUHÉ, J. M., see Gorski, 1556
 DUNAIF, G., see Jackson, 445
 DUNAMS, D., see Pritt, 2144
 DURÁN-QUINTANA, M. C., see Bautista-Gallego, 1412
 DYKES, G. A., see Chia, 2421; Eglezos, 613; Goulter, 1338
 DZIERŻANOWSKA, D., see Rożynek, 602
 DZIERŻANOWSKA-FANGRAT, K., see Rożynek, 602

- EDGE, K., see Nelson, 365
 EDGE, S., see Allen, 264
 EDRINGTON, T. S., Multidrug-resistant *Salmonella*, 27
 EGERVÁRN, M., see Huys, 339
 EGLEZOS, S., Bacteriology of raw chicken nuggets, 613; Microbiology of raw nuts, 402
 EIFERT, J., see Whitney, 820
 ELFERING, K., see Smith, 2153
 ELIZAQUÍVEL, P., DNA extraction for PCR detection of food pathogens, 2110
 ELLERBROEK, L., see Pappelbaum, 735
 EL-MECHERFI, K. E., see Kaddouri, 1270
 EL-SHAROUD, W. M., *S. aureus* and Domiati cheese, 2567
 ELSSEER, D., see Tolvanen, 1550
 EMER, Z., Decontamination of *E. coli* with ozone, 914
 EMERY, J., see Allen, 264
 ENGELBREKTON, A. L., see Tian, 1696
 EREMIN, S. A., see Shim, 781
 ESCARTIN, E. F., see Orozco, 60; Orozco R., 676
 ESTES, S., see Jackson, 445
 ETTINGER, M. R., see Pao, 1544
 EVANS, M., see Pritt, 2144
- FABBI, L. M., see Tomazelli, 2442
 FABIANSOON, S., see Pointon, 1123
 FAILS, A. D., see Dewell, 573
 FAJARDO, V., see Martín, 564
 FAJARDO BERNÁRDEZ, P., Effects of a probiotic culture on performance of weaned piglets, 1797
 FAKHR, M. K., see Nde, 386
 FANNING, S., see Healy, 1372
 FARBER, J. M., see Medeiros, 2087
 FARKAS, B. E., see Viazis, 109
 FARMER, D., see Cheng, 2436
 FARNIR, F., see Delhalle, 1320
 FARRÈ, R., see Fiore, 979
 FAVIA, P., see Conte, 119
 FAZIL, A., Application of MCDA to a food safety decision-making problem, 2323
 FEDERIGHI, M., see Ritz, 1612
 FEDORKA-CRAY, P. J., see McGowan-Spicer, 760; Riley, 807
 FEGAN, N., see Chia, 2421; Eglezos, 613
 FENTE, C. A., see Miranda, 2537
 FERNANDEZ, N., see Zorraquino, 1193
 FILES, K. D., see Chan, 420
 FILIPINI, T. A., see Tomazelli, 2442
 FINO, V. R., UV-light inactivation of viruses on fresh produce, 908
 FIORE, A., In vitro digestion and bioavailability of LDPS, 979
 FLAHERTY, D., see Gorski, 1556
 FLEISHMAN, G. J., see Keller, 775
 FLICK, G. J., see Sheldon, 345
 FOGLIANO, V., see Fiore, 979
 FOODNET EMERGING INFECTIONS PROGRAM WORKING GROUP, see Nelson, 365
 FORBES, L. B., Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
 FORMENTIN, K., see Rhoades, 2272
 FOX, J. T., *E. coli* O157:H7 in feces and on beef carcasses, 1761
 FRANCO, C. M., see Miranda, 2537
 FRANK, J. F., see Berrang, 66; Sanders, 271
 FRANZ, C., see Yao, 431
 FRANZ, E., Linking soil quality to human cadmium exposure, 2504
 FRATAMICO, P., see Gallegos-Robles, 2217
 FRATAMICO, P. M., see Orozco R., 676
 FRAVALO, P., see Chemaly, 1996
 FREZZA, A. L. C., see Tomazelli, 2442
 FRIEDMAN, M., see Juneja, 1604; Ravishankar, 1145
 FRISHMAN, D., see Grimm, 13
 FRUSTOLI, M. A., see Ravys, 1817
 FRY, F. S., see Yancy, 210
- FRYE, J. G., see McGowan-Spicer, 760
 FU, T.-J., Growth of *Salmonella* during sprouting, 888
 FU, T.-J., see Jackson, 445
 FUCIÑOS GONZÁLEZ, C., see Fajardo Bernárdez, 1797
 FUJII, T., see Kimura, 468; Takahashi, 839
 FUKAYA, T., see Kimura, 468
 FULLONE, M. R., see Fiore, 979
 FUNG, D. Y. C., see Lai, 865
 FURUKAWA, I., see Asai, 1460
 FURUNO, J. P., see Nelson, 365
- GAILUNAS, K. M., Bacterial reduction in UV irradiated chill brines, 629
 GAJADHAR, A. A., Al-Adhami, 1960; Forbes, 558
 GALLAGHER, D. L., see Sheldon, 345
 GALLEGOS-ROBLES, M. A., Identification of *Salmonella* serotypes by PCR-RFLP, 2217
 GALLO, M., see Fiore, 979
 GÁLVEZ, A., see Cobo Molinos, 2460
 GAMBLE, H. R., see Forbes, 558
 GARAU, A., see Schirra, 967
 GARBER, E. A. E., Detection of abrin, 1868; Detection of melamine, 590; Ricin and abrin in beverages, 1875
 GARCÍA, M. L., see Solas, 1273
 GARCÍA, T., see Martín, 564
 GARCÍA-RISCO, M. R., see Anadón, 790
 GARDINI, F., see Sado Kamdem, 2572; Torriani, 93
 GARLAND, J., see Vega, 522
 GARRIDO-FERNÁNDEZ, A., see Bautista-Gallego, 1412
 GAST, R. K., *Salmonella* Enteritidis growth on egg yolk membranes, 1905
 GATTO, V., see Torriani, 93
 GAUTHIER, M., see Blais, 2301
 GELLYNCK, X., Economics of *Campylobacter* control in poultry, 479
 GENDEL, S. M., see Jackson, 445
 GEORÑARAS, I., see Barmpalia-Davis, 2014; Carlson, 1343; Simpson, 494
 GEVEKE, D. J., see Ukuku, 684
 GHAFIR, Y., Hygiene indicator criteria for meat in Belgium, 35
 GHAZALI, F. M., see Chai, 2448
 GIBBS, P. S., see Tabe, 539
 GIBSON, G. R., see Rhoades, 2272
 GIL, M. I., see Allende, 2514; Selma, 286
 GILBERT, J., see Şenyuva, 1500
 GILL, A. O., Killing *E. coli* O157 with high pressure processing, 2182
 GILLESPIE, J., see Nelson, 365
 GIOBBE, S., see Schirra, 967
 GIOTIS, E. S., Contribution of sigma B factor in alkali tolerance in *L. monocytogenes*, 1481
 GIULIANI, G., see Di Cagno, 1491
 GJERDE, B., see Robertson, 959
 GOBBETTI, M., see Di Cagno, 1491
 GÖBEL, K. A., see Schönenbrücher, 2059
 GODARD, C., see Rasschaert, 146
 GODDARD, J. M., Rechargeable antimicrobial polyethylene, 2042
 GODWIN, S. L., see Kilonzo-Nthenge, 608
 GOLDEN, D. A., see Critzer, 1469; Zhong, 1015
 GOMES, A. P., see Kongo, 986
 GOMES, B. C., see Minei, 634
 GONZALES BARRON, U., Application of meta-analysis of food safety, 1330
 GONZALEZ, C., see Venturini, 1701
 GONZÁLEZ, I., see Martín, 564
 GONZÁLEZ, R., see De las Rivas, 657
 GONZÁLEZ-MUÑOZ, M., see Rodríguez-Mahillo, 2578; Solas, 1273
 GONZÁLEZ-PEÑAS, E., see Amézqueta, 1422
 GOOCH, J., see Baker-Austin, 2552
 GOODRIDGE, L. D., see Willford, 825
 GORSKI, L., *L. monocytogenes* sprout colonization and stress response, 1556
 GOULTER, R. M., Effect of different water temperatures on knife decontamination, 1338

- GRACE, D., Risk assessment for STEC in unpasteurized milk, 257
- GRAM, L., see Jensen, 1028
- GRANDIN, T., see Dewell, 1114, 2228
- GRANT, K., see Little, 309
- GRANT, M. A., *E. coli* O157:H7 enrichment in produce, 139
- GRAUMANN, G. H., *E. coli* O157:H7 inhibition in dry sausage by mustard powder, 486
- GRAY, J. T., see Riley, 807
- GREENWOOD, M., see Little, 19
- GREGORI, R., Potassium sorbate for controlling postharvest brown rot, 1626
- GREIG, J. D., see Todd, 2339, 2582
- GREINER, S. T., see Sawyer, 405
- GRGURINA, I., see Fiore, 979
- GRIF, K., see Pappelbaum, 735
- GRIFFITH, R., see Wesley, 545
- GRIFFITHS, M., see Klotz, 1884
- GRIFFITHS, M. W., see Morales-Rayas, 2543; Thouand, 380
- GRIGORYAN, G., see Loui, 200
- GRIJSPEERDT, K., see Gellynck, 479
- GRIMM, M., Cellulose production in *E. sakazakii* biofilms, 13
- GROVES, E., see D'Amico, 1580
- GRUYTERS, H., see Röder, 2263
- GUAMIS-LÓPEZ, B., see Velázquez-Estrada, 2283
- GUARD-BOULDIN, J., see Gast, 1905
- GUERINI, M. N., see Arthur, 1685; Kalchayanand, 621
- GUERZONI, M. E., see Sado Kamdem, 2572
- GÜLMEZ, M., see Oral, 218
- GURAYA, R., see Gast, 1905
- GUTIERREZ, J., Antimicrobial efficacy and sensory acceptability of plant essential oils, 1846
- GÜVEN, A., see Oral, 218
- HA, S.-D., see Shim, 781
- HADJICHRISTODOULOU, C., Risk factors for food premises unsatisfactory inspection results, 1632
- HAFNER, S., Detection of DRG in advanced meat recovery product, 2307
- HAJDOK, C., see Xie, 903
- HALE, D. S., see Sawyer, 405
- HALET, D., see Gellynck, 479
- HALFMANN, H., see Deilmann, 2119
- HALL, C., see Sasanya, 1205
- HALL, M. R., see Hussein, 927
- HAMAD, S. H., Microbial spoilage of rutab, 1406
- HAMAMA, A., see Beneduce, 2082
- HAMILL, S., Isolation of *Arcobacter* from beef, 850
- HAMILTON, D. R., see Mie, 1442
- HAN, J., see Turgis, 1237
- HANCOCK, D. D., see Cobbold, 1023
- HANNA, M. A., see Voss, 2036
- HANSMAN, G. S., Viral contamination in clams, 1689
- HANSON, D. J., see Bang, 191
- HAO, Y.-Y. D., see Day, 2208
- HARA-KUDO, Y., see Asai, 1460
- HARGRAVES, S. J., see Chrystal, 2526
- HARPER, J. W., see Akdemir Evrendilek, 102
- HARRIS, D. D., see Miller, 2078
- HARRIS, J. A., see Allen, 264
- HARRIS, L. J., see Danyluk, 397
- HARRISON, M. A., see Akins, 46; Kim, 252
- HARTNETT, E., see Gellynck, 479
- HASSAN, H., see Kreske, 2404
- HASSAN, H. M., see Clare, 126
- HASSAN, Y. I., Antifungal activity of *L. parcasei*, 2213
- HAVLIK, J., see Kokoska, 2475
- HAYMAN, M. M., Heat shock induces barotolerance in *L. monocytogenes*, 426
- HE, G.-Q., see Zhou, 1266
- HE, X., Detection of risin in foods, 2053
- HEALY, B., Evaluation of an automated rep-PCR system, 1372
- HEBERT, P. D. N., see Yancy, 210
- HEDBERG, C. W., see Smith, 2153
- HEIN, I., see Pappelbaum, 735
- HELLER, I., see Pappelbaum, 735
- HELLSTRÖM, S., see Tolvanen, 1550
- HENSEL, A., see Scherer, 699
- HENSHILWOOD, K., see Lowther, 1427
- HERNÁNDEZ, P. E., see Martín, 564
- HERNÁNDEZ-HERRERO, M. M., see Velázquez-Estrada, 2283
- HERRERA, A., see Carramiñana, 502
- HERRERO, A. M., see Hoz, 2001
- HETTIARACHCHY, N. S., see Sivarooban, 2105
- HIGA, L. R. G., see Tomazelli, 2442
- HILL, A., see Klotz, 1884
- HILL, D. E., see Forbes, 558
- HILL, J., see Hafner, 2307
- HILL, W. E., see Stopforth, 509; Yancy, 210
- HILLERS, V. N., see Medeiros, 1666
- HING, H. L., see Tang, 2453
- HINTON, A., JR., Poultry skin flora after successive washings, 1707
- HINTON, A., JR., see Musgrove, 1258
- HISA, K., see Izumi, 52, 530
- HO, W.-L., see Chiang, 2589
- HOFFMAN, T. W., see Dewell, 573
- HOLDS, G., see Pointon, 1123
- HOLLEY, R. A., see Graumann, 486; Osaili, 934; Palanichamy, 6
- HOLT, P. S., see Gast, 1905
- HOLZAPFEL, W., see Yao, 431
- HOLZHAUSER, T., see Röder, 2263
- HONG, C.-H., Aerobic plate counts as a measure of HACCP effectiveness, 1248
- HOTCHKISS, A. T., JR., see Rhoades, 2272
- HOTCHKISS, J. H., see Goddard, 2042
- HOUF, K., see Rasschaert, 146
- HOVINGH, E., see Van Kessel, 1967
- HOZ, L., E-beam radiation of dry-cured ham, 2001
- HRICOVA, D., Electrolyzed water in the food industry, 1934
- HSIEH, Y.-H. P., see Rao, 1000
- HU, Y., see Chan, 420
- HUANG, B., see Eglezos, 402, 613
- HUANG, H., see Loui, 200
- HUBY-CHILTON, F., see Al-Adhami, 1960
- HUCKLESBY, L., see Little, 19
- HUME, M. E., see Soni, 134
- HUMPHREY, T. J., see Williams, 835
- HURD, H. S., see Wesley, 545
- HURST, W., see Akins, 46
- HUSSEIN, H. S., Detection of Shiga toxin-producing *E. coli*, 927
- HUYNH, S., see Brandl, 897
- HUYS, G., Antimicrobial susceptibility of *L. parcasei*, 339
- HWANG, J.-K., see Lee, 1926
- HYATT, D. R., see Dewell, 1114, 2228
- IBACH, A., see Röder, 2263
- IBÁÑEZ, E., see Mendiola, 2138
- IGARASHI, H., see Sakai, 1855
- IHARA, H., see Sakai, 1855
- ILIC, S., Coliforms and foodborne pathogens on spinach in packing plants, 2398
- IMSILP, K., see Poapolathep, 870
- INATSU, Y., see Bari, 830
- INGRAM, A., see Nelson, 365
- INGRAM, K. D., see Musgrove, 1258
- INGRAVALLE, F., see Lai, 578
- IRONSIDE, C. J., see Chrystal, 2526
- IRVIN, E. A., see Jensen, 1028
- ISOBE, S., see Bari, 830
- ITURRIAGA, M. H., see Orozco R., 676
- IVERSEN, C., see Grimm, 13; Healy, 1372

- IZUMI, H., On-farm contamination of persimmons, 52; On-farm contamination of satsuma mandarin fruit, 530
- JACKSON, C. R., see McGowan-Spicer, 760
- JACKSON, L. S., Cleaning and validation to prevent allergen cross-contact, 445
- JACOBS-REITSMA, W. F., see van der Fels-Klerx, 1974
- JANES, M., see Melody, 1475
- JANISE, A., see Röder, 2263
- JANKOVSKA, I., see Kokoska, 2475
- JANS, C., see Miescher Schwenninger, 2481
- JAROSZ, M., see Rozynek, 602
- JAYAPRAKASHA, G. K., see Soni, 134
- JAYARAO, B. M., see Van Kessel, 1967
- JAYAS, D. S., see Palanichamy, 6
- JAYKUS, L. A., see Ailes, 2389; Melody, 1475; Viazis, 109
- JENSEN, A., Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- JENSON, I., see Phillips, 1232; Simmons, 373
- JEONG, E. Y., *T. vulgaris* and a stored food mite, 351
- JESUDHASAN, P., see Soni, 134
- JIANG, S.-T., see Chen, 2523
- JIANG, X., see Brown, 1486
- JIN, T., see Ukuku, 1988
- JITTINANDANA, S., see Kamireddy, 973
- JO, C., see Lim, 1505
- JOENS, L., see Ravishankar, 1145
- JOHNSON, M., see Wesley, 545
- JOHNSON, M. G., see Sivaroban, 2105
- JOHNSON, R. P., see Niu, 691
- JOHNSTON, L. M., see Ailes, 2389
- JONES, D. R., see Musgrove, 1258
- JONES, S. L., Foodborne disease outbreaks in the catering industry, 550; Operational practices associated with foodborne disease outbreaks, 1659
- JORDAN, D., see Phillips, 1232
- JØRGENSEN, F., see Williams, 835
- JØRGENSEN, J. S., see Lai, 578
- JUGIAU, F., see Ritz, 1612
- JULOTOK, M., see Giotis, 1481
- JUNEJA, V. J., Facilitated thermal inactivation of *E. coli* O157:H7, 1604
- JUNG, Y. S., see Kim, 1465
- KADDOURI, H., Immunoreactivity of irradiated milk and whey samples, 1270
- KADORA, K. M., see Osaili, 934
- KADOWAKI, K.-I., see Shinoda, 2257
- KAGKLI, D.-M., see Paramithiotis, 1244
- KAILASAPATHY, K., see Paturi, 796
- KALCHAYANAND, N., Antimicrobial interventions of bovine heads, 621
- KALCHAYANAND, N., see Arthur, 1685, 1752, 2177
- KAMIREDDY, N., Acidified sodium chlorite treatment of trout fillets, 973
- KAMRUZZAMAN, M., see Koitabashi, 706
- KANEKO, M., see Asai, 1460
- KANEKO, S., see Asai, 1460
- KANG, D.-H., see Mah, 953
- KANG'ETHE, E., see Grace, 257
- KAO, S.-C., see Chang, 2094
- KARAGÖZLÜ, C., see Demirbaş, 440
- KARNS, J. S., see Van Kessel, 1967
- KATAYAMA, Y., see Ando, 595
- KATHARIOU, S., see Wright, 1791
- KAWAI, Y., see Abou-Taleb, 770
- KAWAMOTO, S., see Bari, 830
- KAWASAKI, K.-I., see Ando, 595
- KEERATIPIBUL, S., Risk assessment of *Listeria* in ready-to-eat chicken production line, 946
- KEGODE, R., see Khaita, 1679
- KELLER, S. E., Modification of the submerged coil, 775
- KELLERMEIER, J. D., see Brooks, 293
- KELLEY, L. C., see Hafner, 2307
- KENDALL, P., see Lenhart, 70
- KENDALL, P. A., see Barmpalia-Davis, 2014; Byelashov, 728; Medeiros, 1666; Simpson, 494
- KENNEY, P. B., see Kamireddy, 973
- KENNON, L., see Kwon, 1651
- KERR, H. J., see Black, 473
- KESKINEN, L. A., *Listeria* transfer from knives to turkey breast, 176
- KESSEN, V., see Atanassova, 2533
- KHAITSA, M. L., Antimicrobial resistance and class 1 integrons in *E. coli*, 1679
- KHAITSA, M. L., see Tabe, 539
- KHALID, M. F., see Pao, 1544
- KHEROUA, O., see Kaddouri, 1270
- KIERMEIER, A., see Mie, 1442; Pointon, 1123
- KIKOKU, Y., Heat-resistant fungi from blueberries, 2030
- KILLNER, M., see Tomazelli, 2442
- KILONZO-NTHENGE, A., *Listeria* and *Enterobacteriaceae* in domestic refrigerators, 608
- KIM, C.-H., see Shim, 781
- KIM, H., Resistance of *E. sakazakii* to desiccation, 940
- KIM, H. G., see Jeong, 351
- KIM, J. K., Transfer of *E. coli* O157:H7 to lettuce from ice, 252
- KIM, J.-Y., see Shim, 781
- KIM, K.-S., see Shim, 781
- KIM, K.-Y., see Shim, 781
- KIM, M.-G., see Shim, 781
- KIM, T. J., Antimicrobial effect of muscadine seed extracts, 1465
- KIM, Y.-S., Growth inhibitory effects of kimchi against foodborne pathogens, 325
- KIMOTO-NIRA, H., Immunomodulatory activity of lactococci, 2124
- KIMURA, B., Proteolytic *C. botulinum* in rice at low pH, 468
- KIMURA, B., see Takahashi, 839
- KIMURA, R., see Kimura, 468
- KING, D. A., see Arthur, 1752
- KING, D. J., see Thomas, 1214
- KINGSLEY, D. H., HPP inactivation of FCV, 1598
- KINUTA, M., see Matsumoto, 1228
- KIŞLA, D., Microbiological quality of stuffed mussels, 617
- KITAGAWA, Y., see Noba, 1038
- KLANGKAEW, N., see Poapolathep, 870, 1931
- KLEIN, G., see Atanassova, 860, 2533; Kuhn, 993; Strotmann, 845; Wasenaar, 1734
- KLEINBAUM, D. G., see Ailes, 2389
- KLONTZ, K. C., U.S. FDA regulatory management of human listeriosis, 1277
- KLOTZ, V., Alkaline phosphatase in cow's, goat's, and sheep's milk, 1884
- KNABEL, S. J., see Hayman, 426; Lomonaco, 2559
- KNIEL, K. E., see Fino, 908
- KOBAYASHI, M., see Kimoto-Nira, 2124
- KOCA, N., see Akdemir Evrendilek, 102
- KOITABASHI, T., STEC and *E. coli* O157:H7 from retail meats in China, 706
- KOKOSKA, L., Influence of extraction methods on *N. sativa* essential oil, 2475
- KONG, M. G., see Perni, 302, 1619
- KONGO, J. M., Bacteriological safety in São Jorge cheese manufacture, 986
- KONIECZNY, P., see Rozynek, 602
- KOOHMARAIE, M., see Arthur, 1685, 1752, 2177; Bosilevac, 1768; Kalchayanand, 621
- KOOLMEES, P. A., see Wijnker, 1199
- KORKEALA, H., see Tolvanen, 1550
- KORSAK, D., see Rozynek, 602
- KORSAK, N., see Delhalle, 1320
- KOTTAPALLI, B., see Stopforth, 509, 625, 653
- KOZAKI, S., see Sakai, 1855
- KREMASTINO, J., see Hadjichristodoulou, 1632
- KRESKE, A. C., Survival of *E. coli* O157:H7 in acid, 2404
- KUHN, K., Detection of PAHs in smoked meat, 993
- KUMAGAI, S., see Poapolathep, 870, 1931
- KUSAMA, T., see Shinoda, 2257

- KUSUTJARIT, N., see Poapolathep, 870
 KWON, J., Consumer food safety knowledge and behaviors, 1651
- LABBE, R., see Rahmati, 1178
 LABUZA, T. P., see Pal, 1806; Xu, 333
 LACROIX, C., see Miescher Schwenninger, 2481
 LACROIX, M., see Turgis, 516, 1237
 LAGENDIJK, E., Domestic refrigeration practices, 1898
 LAI, J., Microscopic method for animal constituents detection in feed-stuffs, 578
 LAI, W. H., Screening clenbuterol by strip and ELISA, 865
 LAMACCHIA, F., see Beneduce, 2082
 LAMPEN, A., see Kuhn, 993
 LAMPILA, L. E., see Obritsch, 1401
 LANCIOTTI, R., see Sado Kamdem, 2572
 LANDETE, J. M., see De las Rivas, 657
 LANDGRAF, M., see Aragon-Alegro, 2115
 LAPARRA, J. M., see Fiore, 979
 LARKIN, J., see Keller, 775
 LARSEN, S., see Wesley, 545
 LAURY, A., see Niebuh, 714
 LAW, B., see Ravishankar, 1145
 LEANO, F., see Smith, 2153
 LEE, H. S., see Jeong, 351
 LEE, H.-Y., see Chai, 2448
 LEE, J. O., see Klontz, 1277
 LEE, K., see Aruscavage, 2384
 LEE, K.-H., see Shim, 781
 LEE, L. Y., Antibacterial activity of xanthorrhizol, 1926
 LEE, M., see Lim, 1505
 LEE, S. H., see Shabala, 1169
 LEES, D. N., see Lowther, 1427
 LEFEBVRE, B., Virulence genes in *E. coli* O157:H7 and non-O157 isolates, 1774
 LEHNER, A., see Grimm, 13
 LEJEUNE, J. T., see Aruscavage, 2384; Ilic, 2398
 LEKROENGSIK, S., see Keeratipibul, 946
 LELEU, G., see Midelet-Bourdin, 2198
 LENHART, J., Consumer assessment of safety/date labeling on RTE meat products, 70
 LE NÔTRE, Y., see Chemaly, 1996
 LEON, J. S., see Ailes, 2389
 LEROI, F., see Midelet-Bourdin, 2198
 LEROY, F., see Ravys, 1817
 LEWIS, M., see Allen, 264
 LEWIS IVEY, M. L., see Aruscavage, 2384
 LI, T.-C., see Hansman, 1689
 LIM, D. G., Comparison of detection methods for CNST, 1505
 LIM, J. H., see Jeong, 351
 LIMITONE, A., see Di Cagno, 1491
 LIMTHAMMAHISORN, S., In vitro cold shock gene expression in *V. vulnificus*, 157
 LIN, J.-H., see Chuang, 1108
 LIN, W., see Cheng, 2436
 LINDELL, A. H., see Baker-Austin, 2552
 LINZ, J. E., see Moorman, 182
 LITTLE, C. L., *Salmonella* contamination of food service eggs, 19; Sandwiches from health care establishments, 309
 LIU, D. W., see Perni, 302
 LIU, S., see Thouand, 380
 LIU, S.-H., see Wong, 416
 LIZARRAGA, T., see Amézqueta, 1422
 LLOBET, J. M., see Martí-Cid, 2148
 LOGUE, C. M., see Nde, 386; Thorsness, 2295
 LOMONACO, S., Epidemiologic relevance of MVLST, 2559
 LONERAGAN, G. H., see Miller, 2078; Riley, 807
 LONERGAN, S. M., see Patton, 2519
 LOPES, G. V., see Aragon-Alegro, 2115
 LOPEZ, L. C., see Conte, 119
 LÓPEZ, P., Role of fat content in trapping volatile compounds, 1889
- LÓPEZ DE CERAIN, A., see Amézqueta, 1422
 LÓPEZ-FANDIÑO, R., see Chicón, 1453
 LÓPEZ-GÁLVEZ, F., see Allende, 2514; Selma, 286
 LÓPEZ-PEDEMONTE, T., see Velázquez-Estrada, 2283
 LORIMER, M., see Pointon, 1123
 LOUI, C., 16S rDNA profiling of alfalfa sprouts-associated bacteria, 200
 LOWTHER, J. A., Detection of noroviruses in oysters, 1427
 LU, S., see He, 2053; Loui, 200
 LU, Y., see Kim, 1465
 LUCAS, R., see Cobo Molinos, 2460
 LUCHANSKY, J. B., Blade tenderization of beef containing *E. coli*, 2190
 LUCHANSKY, J. B., see Orozco R., 676; Porto-Fett, 918
 LUCIA, L. M., see Sawyer, 405
- MACOVEI, L., Enterococcal contamination of food by houseflies, 435
 MADDEN, R. H., see Hamill, 850
 MAFFEI, M., see Sado Kamdem, 2572
 MAH, J.-H., Heat tolerance of *C. sporogenes* spores, 953
 MAI, T., see Stopforth, 625
 MAILLER, S., see Healy, 1372
 MALAKAR, P. K., see Chai, 2448
 MALCATA, F. X., see Kongo, 986
 MALLE, P., see Midelet-Bourdin, 2198
 MALOUIN, F., see Lefebvre, 1774
 MANDERSON, K., see Rhoades, 2272
 MANDRELL, R. E., see Tian, 1696
 MANN, A., see Ortega, 2410
 MANN, D. A., see Beauchat, 1572
 MÄNNIG, A., *Salmonella* identification and strain differentiation by HGMP-IR, 2249
 MANZARDO, G. G. G., see Grimm, 13
 MARCHIS, D., see Lai, 578
 MARCY, J., see Whitney, 820
 MARI, M., see Gregori, 1626
 MARKS, B. P., see Stasiewicz, 279; Tuntivanich, 2548; Warsaw, 153
 MÁRQUEZ-GONZÁLEZ, M., see Neal, 2415
 MARTÍ-CID, R., Dietary intake of hexachlorobenzene, 2148
 MARTÍN, A., see Sánchez, 2497
 MARTÍN, I., Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
 MARTÍN, R., see Martín, 564
 MARTÍN-ÁLVAREZ, P. J., see Chicón, 1453
 MARTÍN-BELLOSO, O., see Raybaudi-Massilia, 1150
 MARTÍNEZ, M. A., see Anadón, 790
 MARTÍNEZ-LARRAÑAGA, M. R., see Anadón, 790
 MARTÍNEZ-PÉREZ, A., see Al-Adhami, 1960; Blais, 392, 2301
 MARTÍNEZ-RODRÍGUEZ, A. J., see Nunez, 1496
 MASON, J. A., see Yancy, 210
 MASON, S. E., Sulfamethazine water medication pharmacokinetics and contamination, 584
 MASSA, S., see Beneduce, 2082
 MATAK, K. E., see Gailunas, 629
 MATARAGAS, M., Survival of *L. monocytogenes*, 1835
 MATISSEK, R., see Röder, 2263
 MATSUMOTO, N., Isolation of *C. jejuni* from bovine bile, 1228
 MATTISON, K., see Pagotto, 1434
 MÄTTÖ, J., see Huys, 339
 MAXIM, J. E., see Neal, 2415
 McALLISTER, T. A., see Niu, 691; Stephens, 648
 McARTHUR, J. V., see Baker-Austin, 2552
 McCARTHY, P. V., see Klontz, 1277
 McCASKEY, P. C., see Hafner, 2307
 McDONALD, C., see Little, 19
 McDOWELL, D. A., see Giotis, 1481
 McEWEN, S., see Fazil, 2323
 McFEETERS, R. F., see Pérez-Díaz, 639
 MCGOWAN-SPICER, L. L., Antimicrobial resistance and virulence in enterococci, 760
 McHUGH, T. H., see Brandl, 897
 MCKAY, A. M., *mecA*-positive *Staphylococcus* spp. in raw milk, 186

- McKEAN, J. D., see Wesley, 545
 McKEE, S. R., see Bauermeister, 1119
 McKENZIE, H., see Pointon, 1123
 McLANDSBOROUGH, L. A., see Rodriguez, 170
 McLAUHLIN, J., see Little, 309
 McMEECHAN, A., see Williams, 835
 McMEEKIN, T. A., see Chia, 2421
 MECA, G., see Tafuri, 2133
 MEDEIROS, D. T., *Campylobacter* spp. in raw and ready-to-eat foods, 2087
 MEDEIROS, L., see Lenhart, 70
 MEDEIROS, L. C., Cancer patients and food safety, 1666
 MEDUS, C., see Smith, 2153
 MEILE, L., see Miescher Schwenninger, 2481
 MEINERSMANN, R. J., see Berrang, 66; McGowan-Spicer, 760
 MELDRUM, R., see Little, 19
 MELIS, R. A. M., see Schirra, 967
 MELODY, K., Icing to control *V. vulnificus* and *V. parahaemolyticus*, 1475
 MÉNDEZ BATÁN, J., see Fajardo Bernárdez, 1797
 MENDIOLA, J. A., Antimicrobial activity of *D. salina* extracts, 2138
 MENDIZÁBAL, Á., see Rodríguez-Mahillo, 2578
 MENDONCA, A., see Willford, 825
 MENG, J., see Xi, 2067
 MESSENS, W., see Gellynck, 479
 MEYER, J., see Strohbehn, 1641
 MEYER, J. D., see Seman, 1386
 MEYER, S. D., see Smith, 2153
 MICHAELIDIS, C., see Ammor, 1510
 MICHAELS, B. S., see Todd, 2339, 2582
 MIDELET-BOURDIN, G., Superchilled storage of cold-smoked salmon, 2198
 MIE, T., Toxoplasma in ready-to-eat smallgoods, 1442
 MIESCHER SCHWENNINGER, S., Low-molecular-weight antiyeast metabolites of a protective culture, 2481
 MIGHELI, Q., see Schirra, 967
 MILES, B., see Macovei, 435
 MILLER, M. F., Dust exposure and pathogen increases on cattle hides, 2078
 MILLER, M. F., see Brooks, 293
 MILLER, S. A., see Aruscavage, 2384
 MILLETTE, M., see Turgis, 516
 MIMOUN, S., see Kaddouri, 1270
 MINEI, C. C., *L. monocytogenes* biofilms and bacteriocins, 634
 MIRANDA, J. M., Antimicrobial resistance in poultry bacteria, 2537
 MITTAL, G. S., see Xie, 903
 MITZUMACHI, K., see Kimoto-Nira, 2124
 MIYA, S., see Kimura, 468
 MOCHIZUKI, N., see Noba, 1038
 MOE, C. L., see Ailes, 2389
 MOEZELAAR, R., see Van Boeijen, 2007
 MOHAMMED, H. O., see Grace, 257
 MOISAN, H., see Lefebvre, 1774
 MOLINA, M. P., see Zorraquino, 1193
 MOMBA, M. N. B., see Abong'0, 816
 MONEO, I., see Rodríguez-Mahillo, 2578
 MONTVILLE, T. J., see Black, 473; Cruz, 196
 MOORE, M. M., see Yancy, 210
 MOORMAN, M., see Jackson, 445
 MOORMAN, M. A., Altered cell composition in stress-adapted *L. innocua*, 182
 MORALES, P., Multiple-cycle high-pressure treatments, 811
 MORALES, P., see del Olmo, 2468
 MORALES-LOREDO, A., see Gallegos-Robles, 2217
 MORALES-RAYAS, R., Detection of *Campylobacter* spp., 2543
 MORGENSTERN, H., see Tolvanen, 1550
 MORI, M., see Takahashi, 839
 MORITA, Y., see Asai, 1460
 MORLEY, P. S., see Dewell, 1114, 2228
 MORRIS, S., see Phillips, 1232
 MORROW, W. E. M., see Wright, 1791
 MOSQUEDA-MELGAR, J., see Raybaudi-Massilia, 1150
 MOUCHTOURI, V., see Hadjichristodoulou, 1632
 MUKHERJEE, A., Heat inactivation of *E. coli* O157:H7 in nonintact beef, 1349
 MULAS, M., see Schirra, 967
 MULLANE, N., see Healy, 1372
 MUÑOZ, R., see De las Rivas, 657
 MURASE, T., see Matsumoto, 1228
 MURILLO-ARBIZU, M., see Amézqueta, 1422
 MURROS, A. E., see Vihavainen, 2312
 MUSGROVE, M. T., Yeasts from refrigerated commercial shell eggs, 1258
 MWAMBAKANA, J. N., see Abong'0, 816
 NABI, A. Q., see Beneduce, 2082
 NADLE, J., see Nelson, 365
 NAGARAJA, T. G., see Fox, 1761
 NAJARRO, M., see Baker-Austin, 2552
 NAKAGUCHI, Y., see Chai, 2448
 NAKANO, H., see Kikoku, 2030
 NAKAO, M., see Ando, 595
 NAKATANI, M., see Ando, 595
 NANNAPANENI, R., see Wesley, 545
 NASINYAMA, G. W., see Grace, 257
 NATIONAL ADVISORY COMMITTEE ON MICROBIOLOGICAL CRITERIA FOR FOODS, Cooking parameters for safe seafood for consumers, 1287
 NDE, C. W., *Salmonella* detection methods on turkey parts, 386
 NEAL, J. A., E-beam radiation for reducing pathogens in spinach, 2415
 NEILL, S. D., see Hamill, 850
 NELSON, J. M., Food use and practices in long-term care facilities, 365
 NERI, F., see Gregori, 1626
 NERÍN, C., see López, 1889
 NERRIE, B. L., see Pao, 1544
 NGUYEN, N.-L. T., see Yancy, 210
 NIEBUHR, S. E., Surrogates for *Salmonella* in meat, 714
 NIEMIRA, B. A., Cold plasma sanitizing of apples, 1357
 NISBET, D. J., see Crippen, 1981; Edrington, 27
 NISHIBUCHI, M., see Chai, 2448; Koitabashi, 706
 NISHIO, O., see Hansman, 1689
 NIU, Y. D., Fecal versus RAMS detection of *E. coli* O157:H7, 691
 NIXON, C. E., see Zuckschwerdt, 1007
 NOBA, S., Ochratoxin A in wine, 1038
 NÖCKLER, K., see Scherer, 699
 NODA, H., see Asai, 1460
 NODA, M., see Hansman, 1689
 NORTH CUTT, J. K., see Musgrove, 1258
 NOU, X., see Patel, 1922
 NOWAK, B., see Kuhn, 993; Strotmann, 845
 NUNES FILHO, S. A. P., Microbial quality in mineral waters, 1253
 NUÑEZ, M., see del Olmo, 2468; Morales, 811
 NUNEZ, Y. P., Ochratoxin A adsorption by yeast, 1496
 NUTSCH, A. L., see Fox, 1761
 NYCHAS, G.-J. E., see Ammor, 1510; Mataragas, 1835; Panagou, 1393, 1828; Paramithiotis, 1244
 O'BRIEN, S. J., see Jones, 550, 1659
 O'BRIEN, T. W., see Garber, 1868
 OBRITSCH, J. A., Antibacterial effects of polyphosphates, 1401
 ODOMERU, J., see Ilic, 2398; Klotz, 1884
 OH, J.-H., see Brown, 319
 OH, S., see Cho, 2048
 OHKUBO, T., see Sakai, 1855
 OHTSUKA, K., see Asai, 1460
 OKA, T., see Hansman, 1689
 OLAIMAT, A. N., see Osaili, 934
 OLIVEIRA, R., see Silva, 1379
 OLOYA, J., see Khaitsa, 1679; Tabe, 539
 OMORE, A., see Grace, 257
 OMOTE, M., see Noba, 1038

- ONG, C. X., see Yap, 1035
 ONG, K. C., see Yap, 1035
 ONODERA, T., see Shinoda, 2257
 ORAL, N., Extending shelf life of fish, 218
 ORDÓÑEZ, J. A., see Hoz, 2001
 OROZCO, L., Microbiological profile of hydroponic tomatoes, 60
 OROZCO R., L., *Salmonella* on hydroponic tomatoes, 676
 ORTA-RAMIREZ, A., see Stasiewicz, 279; Tuntivanich, 2548; Warsaw, 153
 ORTEGA, Y. R., Parasite inactivation by chlorine dioxide, 2410
 OSAILI, T. M., Inactivation of desiccated *E. sakazakii*, 934
 OSCAR, T. P., Mapping of *Salmonella* contamination on the poultry carcass, 1785; Model for growth of *Salmonella* on chicken frankfurters, 1135
 OYAIZU, H., see Tominaga, 1910
 OZDEMIR, M., see Emer, 914
- PADHI, S., see Black, 473
 PAEZ, P., see Strohehn, 1641
 PAGOTTO, F., Detection and genotyping noroviruses using DNA microarrays, 1434
 PAGOTTO, F., see Blais, 2301
 PAL, A., Estimating *L. monocytogenes* growth kinetics parameters in frankfurters, 1806
 PALANICHAMY, A., Predictive microbiology, 6
 PALCICH, G., see Aragon-Alegro, 2115
 PALMER, S. R., see Jones, 550, 1659
 PAN, Z., see Brandl, 897
 PANAGOU, E. Z., Modeling *L. monocytogenes* in pasteurized vanilla cream, 1828; Neural networks and survival of *L. monocytogenes* in soft cheese, 750; Survival of *B. cereus* during green olive fermentation, 1393
 PAO, S., Microbial quality of fish fillets, 1544
 PAPANIKOLAOU, S., see Belessi, 742
 PAPPELBAUM, K., *L. monocytogenes* in produce processing, 735
 PARAMITHIOTIS, S., Enterococci in spontaneous sausage fermentation, 1244
 PARKER, S., see Forbes, 558
 PAROLARI, G., see Ravyts, 1817
 PARRY, S. M., see Jones, 550, 1659
 PASCUAL, S., *Kudoa* in Patagonian hakes, 2316
 PASTRANA CASTRO, L., see Fajardo Bernárdez, 1797
 PASTUSZCZAK-FRAK, M., see Rasschaert, 146
 PATEL, J. R., Control of airborne microorganisms in meat processing, 1922
 PATIL, B. S., see Soni, 134
 PATRIARCA, A., see Azcarate, 1262
 PATTON, B. S., *E. coli* inhibition by colicin E1, 2519
 PATURI, G., *Lactobacillus* systemic immune functions and bacterial translocation, 796
 PAULSEN, P., Automated MPN determination of *Enterobacteriaceae* in food, 376
 PEARCE, L. E., see Robertson, 1186
 PEJCHALOVÁ, M., Presence and diversity of *Arcobacter* in the Czech Republic, 719
 PEJCHALOVA, M., see Cervenka, 165
 PÉREZ-DÍAZ, I. M., Microbiology of sweet potato puree, 639
 PÉREZ GUERRA, N., see Fajardo Bernárdez, 1797
 PERNI, S., Decontamination of mango and melon by gas plasma, 302; Plasma inactivation of bacteria migrating through fruits, 1619
 PESKOVA, I., see Cervenka, 165
 PESTKA, J. J., see Moorman, 182
 PHAN, L., see Cheng, 2436
 PHEBUS, R. K., see Luchansky, 2190
 PHILLIPS, D., Microbiological quality of Australian retail raw meat, 1232
 PHILLIPS, M., see Paturi, 796
 PHILLIPS, W. A., see Riley, 807
 PHUMKHACHORN, P., see Rattanachaiakunsopon, 2024
 PICART 1 BARROT, L., see Cabedo, 855
 PILLAI, S. D., see Soni, 134; Vega, 522
- PIÑA MILLAN, M. B., see Lai, 578
 PINTO, V. F., see Azcarate, 1262
 PLANTENGA, M., see Nelson, 365
 POAPOLATHEP, A., Deoxynivalenol contamination in wheat products, 1931; Oxytetracycline residue depletion in giant freshwater prawn, 870
 POAPOLATHEP, S., see Poapolathep, 870, 1931
 POINTON, A., Microbiological quality of chicken meat, 1123
 POINTON, A. M., see Mie, 1442
 PORTO-FETT, A. C. S., Controlling pathogens in beef jerky, 918
 POUBOL, J., see Izumi, 52, 530
 PRITT, B., Detection of *Sarcocystis* parasites in retail beef, 2144
 PUEYO, E., see Nunez, 1496
- QUICKERT, S. C., see Seman, 1386
- RADU, S., see Chai, 2448
 RAHMATI, T., *C. perfringens* and *B. cereus* in seafood, 1178
 RAJIC, A., see Fazil, 2323
 RAJKOWSKI, K. T., Radiation D_{10} -values of *L. monocytogenes* on catfish and tilapia, 2278
 RAMASWAMY, H. S., see Gill, 2182
 RAMEZANI, R., see Amiri, 411
 RANDOLPH, S. C., see Yancy, 210
 RANDOLPH, T., see Grace, 257
 RAO, Q., Competitive ELISA to detect bovine blood, 1000
 RASOOLY, R., see He, 2053
 RASSCHAERT, G., Contamination of carcasses with *Salmonella* during poultry slaughter, 146
 RASTALL, R. A., see Rhoades, 2272
 RATTANACHAIKUNSOPON, P., Nisin Z production in *L. lactis*, 2024
 RATTEI, T., see Grimm, 13
 RATTI, R. P., see Minei, 634
 RAVEN, G., see Pointon, 1123
 RAVISHANKAR, S., Inactivation of antibiotic-resistant *Campylobacter*, 1145
 RAVYTS, F., Bacteriocin-producing starters in fermented sausages, 1817
 RAYBAUDI-MASSILIA, R. M., Essential oils in coatings for fresh-cut apples, 1150
 REGLERO, G., see Anadón, 790; Mendiola, 2138
 REICH, F., see Atanassova, 860, 2533
 REID, A. O., see Pao, 1544
 REINEKE, K. F., see Fu, 888
 RENTER, D. G., see Fox, 1761
 REYES-HERRERA, I., Distribution of drug residues in breast muscle tissue, 223
 RHOADES, J., Inhibition of pathogen adhesion, 2272
 RHOADES, J. R., see Little, 19
 RIBEIRO, V. B., see Aragon-Alegro, 2115
 RICE, D. H., see Cobbold, 1023
 RICHT, J. A., see Schönenbrücher, 2059
 RICO-ROMERO, L., see Orozco, 60
 RIDLEY, A. M., see Allen, 264
 RIEDEL, K., see Grimm, 13
 RILEY, D. G., Shedding of foodborne pathogens by Florida cattle, 807
 RILEY, L. W., see Loui, 200
 RITIENI, A., see Tafuri, 2133
 RITZ, M., High-pressure treatment of *L. monocytogenes* in smoked salmon, 1612
 RIVERA, C. S., see Venturini, 1701
 RIVERA, F., see Wesley, 545
 RIVIERE, J. E., see Mason, 584
 RIZZELLO, C. G., see Di Cagno, 1491
 ROBERTSON, L. J., Analysis of shellfish for *Cryptosporidium* and *Giardia*, 959
 ROBERTSON, R. E., *Bacillus* spore inactivation differences, using two pressurizing fluids, 1186
 ROCA, M., see Zorraquino, 1193
 RÖDER, M., Allergen cross-contamination in the manufacture of fine bakery wares, 2263
 RODRIGUEZ, A., Stainless steel surface roughness and finish, 170
 RODRIGUEZ, G., see Gutierrez, 1846

- RODRÍGUEZ, M., see Sánchez, 2497
- RODRÍGUEZ-MAHILLO, A. I., Allergen release by frozen and chilled *Anisakis*, 2578
- RODRIGUEZ-MAHILLO, A. I., see Solas, 1273
- RODRIGUEZ-ROMO, L. A., see Männig, 2249
- RODRÍGUEZ-SAONA, L. E., see Männig, 2249
- ROGERS, P. L., see Yancy, 210
- ROHNER, P., see Templer, 2100
- ROIG-SAGUÉS, A. X., see Velázquez-Estrada, 2283
- ROJAS, M., see Martín, 564
- ROJAS-GRAÜ, M. A., see Raybaudi-Massilia, 1150
- ROMINE, S., see Wright, 1791
- RÖMKENS, P., see Franz, 2504
- ROSENTHAL, B. M., see Pritt, 2144
- RÖSLER, U., see Scherer, 699
- ROSS, T., see Shabala, 1169
- ROTA, C., see Carramiñana, 502
- ROUSE, S., Bioprotective potential of LAB in malting and brewing, 1724
- ROVIRA, P. J., On-farm meat safety, 1919
- ROZEMA, E. A., see Niu, 691
- ROŻYNEK, E., Antibiotic resistance of *Campylobacter* isolates from Poland, 602
- RU, G., see Lai, 578
- RUBY, J., see Carlson, 2223
- RUDELLE, S., see Midelet-Bourdin, 2198
- RUEPP, A., see Grimm, 13
- RUKAYADI, Y., see Lee, 1926
- RUSSELL, S. M., Evaluation of a copper sulfate-based commercial disinfectant, 226
- RYSER, E. T., see Keskinen, 176; Moorman, 182; Tuntivanich, 2548; Warsaw, 153
- RYU, D., see Obritsch, 1401; Voss, 2036
- RYU, J.-H., see Kim, 940
- SACCANI, G., see Ravvys, 1817
- SADO KAMDEM, S., Safety and quality assessment of lagoon grey mullets, 2572
- SAEGERMAN, C., see Delhalle, 1320
- SAIDI, D., see Kaddouri, 1270
- SAJFRTOVA, M., see Kokoska, 2475
- SAKAI, F., Characteristics of enterotoxin H-producing *S. aureus*, 1855
- SAKUMA, K., see Kimura, 468
- SALAFRANCA, J., see López, 1889
- SALMAN, M. D., see Dewell, 573, 1114, 2228
- SALMIERI, S., see Turgis, 516
- SALOMÃO, B. C. M., Sanitizers for inactivation of *P. expansum* on apples, 643
- SALTER, R., see Jackson, 445
- SAMADPOUR, M., see Stopforth, 509, 625, 653
- SÁNCHEZ, B., Efficient fungal DNA extraction method for RAPD-PCR, 2497
- SANCHEZ, J., see Fazil, 2323
- SANDERS, S. Q., Temperature and nutrient effects on *C. jejuni*, 271
- SANDERSON, M. W., see Fox, 1761
- SANT'ANA, A. S., see Nunes Filho, 1253
- SANTOYO, S., see Mendiola, 2138
- SAPUTRA, T., see Pointon, 1123
- SARAVANOS, E. K. A., see Panagou, 1393
- SASANYA, J. J., LC-UV-MS method for DON, DON-3-glucoside, and pigment, 1205
- SATTAR, S. A., see Medeiros, 2087
- SAVARY, W. E., see Yancy, 210
- SAVINO, M., see Angelini, 2488
- SAWYER, J. E., Xylitol to reduce microbial adhesion, 405
- SCALLAN, E., see Nelson, 365
- SCANGA, J. A., see Bowling, 83; Byelashov, 728; Carlson, 1343; Dewell, 573, 1114, 2228; Mukherjee, 1349; Simpson, 494
- SCHAFFNER, D. W., see Dominguez, 2429
- SCHERER, K., Time course of infection with *Salmonella* Typhimurium in fattening pigs, 699
- SCHIRRA, M., Fludioxonil residues and control of blue and gray mold on pears, 967
- SCHNEIDER, M. J., Danofloxacin in beef muscle, 1720
- SCHÖNENBRÜCHER, H., Detection of central nervous system tissues in meat products, 2059
- SCHOPF, E., see Paulsen, 376
- SCHROEDER, M., see Lenhart, 70
- SCHUKKEN, Y. H., see Van Kessel, 1967
- SCHWARZ, P. B., see Zhou, 1266
- SEIDEL, A., see Kuhn, 993
- SEKINE, M., see Tominaga, 1910
- SELMA, M. V., Water disinfection from the fresh-cut vegetable industry, 286
- SELMA, M. V., see Allende, 2514
- SELVAM, A., see Chang, 2094
- SEMAN, D. L., Antimycotics and diacetate in RTE meats, 1386
- SEMBENI, S., see Torriani, 93
- SENEVIRATHNE, R., see Melody, 1475
- SEÑORÁNS, F. J., see Anadón, 790; Mendiola, 2138
- ŞENYUVA, H. Z., LC-TOF-MS analysis of mycotoxins in figs, 1500
- SEOKA, M., see Ando, 595
- SERA, K., see Izumi, 530
- SEXTON, M., see Pointon, 1123
- SHABALA, L., *Listeria* inactivation affected by sequences of stresses, 1169
- SHACKELFORD, S. D., see Arthur, 1685, 1752, 2177
- SHAKER, R. R., see Osaili, 934
- SHALLO THESMAR, H., see Chen, 2073
- SHAMA, G., see Pemi, 302, 1619
- SHARMA, M., see Allen, 264
- SHAZER, A. G., see Keller, 775
- SHEFFIELD, C. L., see Crippen, 1981
- SHELDON, T. A., Effect of HPP on freely suspended and bivalve-associated T7, 345
- SHERWOOD, J. S., see Thorsness, 2295
- SHI, X., see Fox, 1761
- SHIH, D. Y.-C., see Chang, 2094
- SHIM, J.-S., see Lee, 1926
- SHIM, W.-B., Enhanced rapidity for *L. monocytogenes* detection, 781
- SHIN, D.-H., see Kim, 325
- SHIN, K.-J., see Cho, 2048
- SHINODA, N., Developing group-specific primers, 2257
- SILETZKY, R. M., see Wright, 1791
- SILVA, J. L., see Kim, 1465
- SILVA, S., *L. monocytogenes* adhesion and viability, 1379
- SIMMONS, G., see Ameratunga, 2334
- SIMMONS, J., Effect of incubation temperature on APCs, 373
- SIMMONS-ARNOLD, L., see Pritt, 2144
- SIMPSON, C. A., *L. monocytogenes* on salami during storage, 494
- SIMPSON, C. A., see Dewell, 1114, 2228
- SINIGAGLIA, M., see Bevilacqua, 1366; Cibelli, 2129; Conte, 119
- SITES, J., see Niemira, 1357
- SIVAROOBAN, T., Natural antimicrobial combinations against *L. monocytogenes*, 2105
- SKANDAMIS, P. N., see Belessi, 742
- SLÁMOVÁ, M., see Pejchalová, 719
- SLIDER, S. D., see Kamireddy, 973
- SMALL, A., see Goulter, 1338
- SMITH, D. M., see Stasiewicz, 279
- SMITH, G. C., see Bowling, 83; Carlson, 1343, 2223; Dewell, 573, 1114, 2228; Mukherjee, 1349; Sugiura, 802
- SMITH, K. E., Salmonellosis from stuffed chicken products, 2153
- SMITH, M. A., see Jensen, 1028
- SMULDERS, F. J. M., see Paulsen, 376
- SNEED, J., see Strohhahn, 1641
- SOFOS, J. N., see Barmpalia-Davis, 2014; Bowling, 83; Byelashov, 728; Carlson, 1343, 2223; Lenhart, 70; Mukherjee, 1349; Simpson, 494
- SOLAS, M. T., *Anisakis* antigens in fish muscle, 1273
- SOLAS, M. T., see Rodríguez-Mahillo, 2578
- SONI, K. A., Identification of AI-2 inhibitors from ground beef, 134

- SOVOVA, H., see Kokoska, 2475
 SPANO, G., see Beneduce, 2082; El-Sharoud, 2567
 SPARKS, N., see Allen, 264
 SPÖRNDLI, C., see Miescher Schwenninger, 2481
 STAHL, C. H., see Patton, 2519
 STANFORD, K., see Niu, 691; Stephens, 648
 STASIEWICZ, M. J., Modeling the effect of sublethal history, 279
 STEPANAUSKAS, R., see Baker-Austin, 2552
 STEPHAN, R., see Grimm, 13; Hricova, 1934
 STEPHENS, T. P., Fecal pats inoculated with *E. coli* O157:H7 to mimic a super shedder, 648
 STEPHENS, T. P., see Brooks, 293; Niu, 691
 STERGIOU, V., see Mataragas, 1835
 STOJANOVIC, J., see Kim, 1465
 STOPFORTH, J. D., Antimicrobial interventions for leafy greens, 625; Heat resistance of *Salmonella* serotypes, 509; Sterility testing of an aseptic dispensing valve, 653
 STORRS, M., see Healy, 1372
 STROHBEHN, C., Hand washing frequencies and procedures, 1641
 STROTSMANN, C., *Y. enterocolitica* in ground pork packaged under modified atmospheres, 845
 STUTTARD, E., see Eglezos, 402, 613
 SUGITA-KONISHI, Y., see Poapolathep, 1931
 SUGIURA, K., A comparison of BSE risk between Japanese and U.S. beef, 802
 SUGIURA, K., see Shinoda, 2257
 SUGIURA, T., see Shinoda, 2257
 SUHALIM, R., see Stopforth, 509
 SUMNER, J., see Phillips, 1232; Pointon, 1123; Simmons, 373
 SUMNER, S. S., see Gailunas, 629
 SUPAN, J., see Melody, 1475
 SURICO, R. F., see Di Cagno, 1491
 SURMAN-LEE, S., see Little, 19
 SUTTON, M. T., see Hafner, 2307
 SUWALAK, S., see Voravuthikunchai, 1223
 SUWELACK, C., see Röder, 2263
 SUZUKI, C., see Kimoto-Nira, 2124
 SUZZI, G., see Torriani, 93
 SWAISGOOD, H. E., see Clare, 126
 SWANSON, K. M. J., see Jackson, 445
 SWAYNE, D. E., see Thomas, 1214
 SZABÓ, I., see Scherer, 699
 SZYCH, J., see Rożynek, 602
 SZYMANSKI, M., see Cobbold, 1023
- TABE, E. S., Probiotics effect on *E. coli* O157:H7 and *Salmonella* shedding in cattle, 539
 TAFURI, A., Rapid method used to analyze ochratoxin A, 2133
 TAGASHIRA, N., see Kikoku, 2030
 TAKAHASHI, H., Grouping of food isolated bacteria by PCR-SSCP in food, 839
 TAKATORI, K., see Asai, 1460
 TAKEDA, N., see Hansman, 1689
 TAMPLIN, M. L., see Orozco R., 676; Simmons, 373
 TANAKA, Y., see Takahashi, 839
 TANG, J., see Mah, 953
 TANG, J. Y. H., *V. cholerae* survival on different types of cooked rice, 2453
 TANIWAKI, T., see Matsumoto, 1228
 TARR, P. I., see Cobbold, 1023
 TASSOU, C. C., see Panagou, 1393
 TAUXE, R. V., see Vojdani, 356
 TAYLOR, T. M., see Black, 473
 TEIXEIRA, P., see Silva, 1379
 TEIXIDÓ I CANELLES, A., see Cabedo, 855
 TEJADA, M., see Rodríguez-Mahillo, 2578; Solas, 1273
 TEMPLER, S. P., Enterococci from human patients and foods, 2100
 TENGE, B. J., see Yancy, 210
 TERMINIELLO, L., see Azcarate, 1262
 TERSTEEG, M. H. G., see Wijnker, 1199
- TESSARO, S. V., see Forbes, 558
 THELEMANN, C. A., see Moorman, 182
 THIPPAREDDI, H., see Luchansky, 2190; Yarbava, 77
 THOMAS, C., Thermal inactivation of viruses in chicken meat, 1214
 THONART, P., see Yao, 431
 THORSNESS, J. L., *Campylobacter* prevalence in poultry production, 2295
 THOUAND, G., Detection of *Salmonella* using a simple phage assay, 380
 THRELFALL, E. J., see Little, 19
 THRESHOLD WORKING GROUP, Approaches to establish thresholds, 1045
 TIAN, P., Seasonal binding of recombinant norovirus to GI cells in oysters, 1696
 TIPPETT, R., see Jackson, 445
 TITTO, A. W., see Brooks, 293
 TODD, E. C. D., Infective doses and pathogen carriage, 2339; Sources of contamination and pathogen excretion from infected persons, 2582
 TODD, E. C. D., see Hong, 1248; Keskinen, 176
 TOFALO, R., see Torriani, 93
 TOLVANEN, R., Survival of *L. monocytogenes* in dry sausage, 1550
 TOMAZELLI, I. B., BAX system PCR method for *Salmonella* versus Brazil's official method, 2442
 TOMAZELLI JÚNIOR, O., see Tomazelli, 2442
 TOMINAGA, T., Tracing the origin of coliform bacteria, 1910
 TOQUIN, M.-T., see Chemaly, 1996
 TORRES, M. P., see Ortega, 2410
 TORRIANI, S., Evaluation of *tdc* and its transcript in bacteria and foods, 93
 TOWNSEND, J. C., see Bauermeister, 1119
 TRAINER, T., see Pritt, 2144
 TRAN, N. N., see Cheng, 2436
 TRAVAGLIA, F., see Bonetta, Silvia, 205
 TRONCO, V. M., see Tomazelli, 2442
 TRUJILLO, S., see Day, 2208
 TRUONG, V.-D., see Pérez-Díaz, 639
 TRUTTMANN, S., see Miescher Schwenninger, 2481
 TSUJISAWA, T., see Ando, 595
 TSUKADA, Y., see Izumi, 52
 TSUKAMASA, Y., see Ando, 595
 TUCKFIELD, R. C., see Baker-Austin, 2552
 TUNTIVANICH, V., Thermal inactivation of *Salmonella* in turkey breast, 2548
 TURGIS, M., Combined treatments with irradiation on beef microbiology, 1237; Radiosensitivity of pathogens in ground beef, 516
 TYLER, K., see Blais, 2301
- UKUKU, D. O., Membrane damage and inactivation of bacteria in apple juice by RFEF 684; Thermal inactivation of bacteria in liquid egg, 1988
 UYTENDAELE, M., see Baert, 1590
 ÜZGÜN, Y., see Kışla, 617
- VACHON, P., see Thouand, 380
 VALDIVIA, E., see Cobo Molinos, 2460
 VALTEROVA, I., see Kokoska, 2475
 VAMVAKOULA, P., see Panagou, 1393
 VAN, K. T., see Cheng, 2436
 VAN ASSELT, E. D., see van der Fels-Klerx, 1974
 VAN BOEIJEN, I. K. H., *L. monocytogenes* HHP inactivation, 2007
 VAN BRAKEL, R., see van der Fels-Klerx, 1974
 VAN COILLIE, E., see Baert, 1590
 VANCUTSEM, J., see Lai, 578
 VANDAMME, P., see Huys, 339
 VAN DER FELLS-KLERX, H. J., *Salmonella* prevalence in the broiler supply chain, 1974
 VAN DER FELLS-KLERX, I., see Franz, 2504
 VAN DER VOET, H., see van der Fels-Klerx, 1974
 VAN KESSEL, J. S., Predicting *Salmonella* prevalence in a dairy herd, 1967
 VAN LEUSDEN, F. M., see Verhoeff-Bakkenes, 1018
 VANPELT, O. M., see Fu, 888
 VAN RAAMSDONK, L., see Franz, 2504

- VAN SINDEREN, D., see Rouse, 1724
 VAN TASSELL, C. P., see Van Kessel, 1967
 VARZAKAS, T., see Hadjichristodoulou, 1632
 VATANSEVER, L., see Oral, 218
 VÁZQUEZ, B. I., see Miranda, 2537
 VEGA, E., Virus adsorption to lettuce, 522
 VEGA-P., A., see Gallegos-Robles, 2217
 VELARDE, S., see Gallegos-Robles, 2217
 VELAZCO, J. I., see Rovira, 1919
 VELÁZQUEZ-ESTRADA, R. M., Inactivation of *Salmonella* Senftenberg 775W by UHPH, 2283
 VELUGOTI, P. R., see Yarbava, 77
 VENTURINI, M. E., Antibacterial activity of mushroom extracts, 1701
 VERHOEFF-BAKKENES, L., *C. jejuni* cross-contamination, 1018
 VERMEERSCH, M., see Baert, 1590
 VERNOCCHI, P., see Sado Kamdem, 2572
 VERNOOIJ, J. C. M., see Wijnker, 1199
 VIAENE, J., see Gellynck, 479
 VIAZIS, S., High-pressure processing of human milk, 109
 VIETHS, S., see Röder, 2263
 VIHAVAINEN, E. J., LAB spoilage of vegetable sausages, 2312
 VILLAESCUSA, R., see Allende, 2514
 VOJDANI, J. D., U.S. juice-associated outbreaks, 356
 VON MUEFFLING, T., see Strotmann, 845
 VORAVUTHIKUNCHAI, S. P., *Q. infectoria* as anti-*E. coli* agent, 1223
 VOSS, K. A., Reduced toxicity of fumonisin B₁, 2036
 VYTRÁSOVÁ, J., see Cervenka, 165; Pejchalová, 719
- WAGNER, B. A., see Dewell, 1114, 2228
 WAGNER, D., see Wesley, 545
 WAGNER, M., see Pappelbaum, 735
 WAITE, J. G., Antimicrobial synergy between high pressure and Red No. 3, 1861
 WALKER, J. L., see Garber, 1868
 WANG, B., see Brown, 319
 WANG, J.-Y., see Chang, 2094
 WANG, K.-C., see Chuang, 1108
 WANNAPAT, N., see Poapolathep, 870
 WARD, L., see Williams, 835
 WARDAK, S., see Rozynek, 602
 WARREN-SERNA, W., see Carlson, 2223
 WARRINER, K., see Klotz, 1884; Xie, 903
 WARSOW, C. R., *Salmonella* migration into marinated turkey, 153
 WASHINGTON, J. D., see Yancy, 210
 WASSENAAR, T. M., Regulation of safety of probiotics, 1734
 WATANABE, M., Mycotoxins produced in apples, 1714
 WEAVER, H., see Allen, 264
 WEBBER, A., see Pérez-Díaz, 639
 WELLS, A., see Rhoades, 2272
 WELLS, J. E., see Arthur, 1685; Berry, 2233
 WENG, W. L., see Kim, 1465
 WERNITZNIG, F., see Lai, 578
 WESLEY, I. V., *L. monocytogenes* in cull sows and pork, 545; Prevalence of *Y. enterocolitica*, 1162
 WHALEN, L. R., see Dewell, 573
 WHEELER, T. L., see Arthur, 1685, 1752, 2177; Kalchayanand, 621
 WHITING, R. C., see Day, 2208; De Jesús, 1915
 WHITNEY, B. M., HPP in combination with antimicrobials, 820
 WIDMER, K., see Soni, 134
 WIEDMANN, M., see Chan, 420
 WIJNKER, J. J., Lymphoid tissue reduction in bovine small intestines, 1199
 WILDEMAUWE, C., see Rasschaert, 146
 WILKINSON, B. J., see Giotis, 1481
- WILKINSON, M. G., see Cronin, 2168
 WILLFORD, J., Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
 WILLIAMS, D., see Jensen, 1028
 WILLIAMS, J. B., see Dowd, 1673
 WILLIAMS, L. K., Detection methods for *Campylobacter* spp., 835
 WILLIAMS, R. C., see Gailunas, 629; Whitney, 820
 WILSON, A. N. S., see Kwon, 1651
 WILSON, I., see Little, 19
 WOERNER, D. R., see Dewell, 573
 WOLFFS, P. F. G., see Morales-Rayas, 2543
 WOLFGANG, D. R., see Van Kessel, 1967
 WOLF-HALL, C., see Sasanya, 1205
 WONG, H.-C., Low-salinity stress in *V. vulnificus*, 416
 WOROBO, R. W., see Salomão, 643
 WRIGHT, S. L., *Campylobacter* in paired turkey and swine farms, 1791
 WUNDERLICH, J., see Deilmann, 2119
 WÜRZNER, R., see Pappelbaum, 735
- XI, M., PFGE scheme for subtyping *Salmonella*, 2067
 XIE, Y., UV light-hydrogen peroxide decontamination of lettuce, 903
 XIONG, Y. H., see Lai, 865
 XU, S., Inactivation of *B. anthracis* in milk, 333
 XU, Y., see Lai, 865; Niu, 691
- YANAHIRA, S., see Sakai, 1855
 YANCY, H. F., DNA barcoding and the *Regulatory Fish Encyclopedia*, 210
 YANG, C.-H., see Chuang, 1108
 YANG, S.-S., see Chang, 2094
 YANG, Z.-Y., see Shim, 781
 YANO, D. M. Y., see Tomazelli, 2442
 YAO, A. A., Storage stability of freeze-dried lactic acid bacteria, 431
 YAP, K. L., SA11 rotavirus survival in fresh tropical fruit juices, 1035
 YAP, K. L., see Tang, 2453
 YARBAEVA, S. N., Control of *C. perfringens* in Tajik sambusa, 77
 YEH, K.-S., see Chuang, 1108
 YEMM, R. S., see Bowling, 83
 YIN, L.-J., see Chen, 2523
 YOKOI, A., see Takahashi, 839
 YOON, Y., see Carlson, 1343; Mukherjee, 1349; Simpson, 494
 YOSHIDA, T., see Shinoda, 2257
 YOUSEF, A. E., see Männig, 2249; Waite, 1861
 YU, R.-C., see Chiang, 2589
- ZEMPLAK, T. S., see Yancy, 210
 ZHANG, H., see Ukuku, 1988
 ZHANG, H. Q., see Ukuku, 684
 ZHAO, S., see Xi, 2067
 ZHENG, J., see Xi, 2067
 ZHENG, Z., see Clare, 126
 ZHENG, Z.-B., see Kim, 325
 ZHONG, Q., Dense phase CO₂ processing of spinach, 1015
 ZHOU, B., Occurrence of bound deoxynivalenol in *Fusarium*-infected barley, 1266
 ZHOU, S., see Moorman, 182
 ZHU, L., see Ravishankar, 1145
 ZHU, Y., see Brandl, 897
 ZORRAQUINO, M. A., Antibiotics in milk, 1193
 ZUCKSCHWERDT, J. B., Efficacy of drug test kits for adulterated food, 1007
 ZUREK, L., see Macovei, 435
 ZWEIFEL, C., see Hricova, 1934
 ZWIETERING, M. H., see Van Boeiijen, 2007

SUBJECT INDEX

- Abrin
 Detection of abrin, 1868
 Ricin and abrin in beverages, 1875
- Acaricidal activity
T. vulgaris and a stored food mite, 351
- Acetic acid
 Survival of *E. coli* O157:H7 in acid, 2404
- Acid
 Altered cell composition in stress-adapted *L. innocua*, 182
- Acid adaptation
L. monocytogenes on salami during storage, 494
- Acid adapted
Salmonella in tomatoes, 1572
- Acidic
 Evaluation of a copper sulfate-based commercial disinfectant, 226
- Acidified sodium chlorite
 Acidified sodium chlorite treatment of trout fillets, 973
 Antimicrobial interventions for leafy greens, 625
- Acid resistance
 Survival of *L. innocua* in dairy products, 742
- Acid stress
L. monocytogenes sprout colonization and stress response, 1556
Listeria inactivation affected by sequences of stresses, 1169
- Activation energy
 Storage stability of freeze-dried lactic acid bacteria, 431
- Active packaging
 Active packaging, 119
- Acute safety study
 Acute toxicity of rosemary extracts, 790
- Adaptation
 Contribution of sigma B factor in alkali tolerance in *L. monocytogenes*, 1481
 Low-salinity stress in *V. vulnificus*, 416
- Adhesion
L. monocytogenes adhesion and viability, 1379
 Xylitol to reduce microbial adhesion, 405
- Adsorption
 Virus adsorption to lettuce, 522
- Advanced meat recovery
 Detection of DRG in advanced meat recovery product, 2307
- Advanced oxidation technologies
 Water disinfection from the fresh-cut vegetable industry, 286
- Aerobic bacteria colony counts
 Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
- Aerobic colony counts
 Hygiene indicator criteria for meat in Belgium, 35
- Aerobic plate count (APC)
 Aerobic plate counts as a measure of HACCP effectiveness, 1248
 Effect of incubation temperature on APCs, 373
 Grouping of food isolated bacteria by PCR-SSCP in food, 839
- Aeromonas*
 Multiplex PCR detection of enterotoxin genes in *Aeromonas* spp., 2094
- Agricultural water
 On-farm contamination of persimmons, 52
- AI-2
 Identification of AI-2 inhibitors from ground beef, 134
- ail* gene
 Prevalence of *Y. enterocolitica*, 1162
- Air
 Evaluation of a copper sulfate-based commercial disinfectant, 226
- Airborne contamination
 Control of airborne microorganisms in meat processing, 1922
- Air-dried sausage
 Detection of *S. aureus*, 191
- Alfalfa sprouts
 Growth of *Salmonella* during sprouting, 888
 16S rDNA profiling of alfalfa sprouts-associated bacteria, 200
- Alginate-based coating
 Essential oils in coatings for fresh-cut apples, 1150
- Alkaline phosphatase
 Alkaline phosphates in cow's, goat's, and sheep's milk, 1884
- Alkaline stress
 Contribution of sigma B factor in alkali tolerance in *L. monocytogenes*, 1481
- Allergen(s)
 Allergen release by frozen and chilled *Anisakis*, 2578
Anisakis antigens in fish muscle, 1273
 Cleaning and validation to prevent allergen cross-contact, 445
- Allergic reactions
 Latex glove use by food handlers, 2334
- Allergic thresholds
 Approaches to establish thresholds, 1043
- Almonds
 Decontamination of almonds with infrared heat, 897
 Microbiology of raw nuts, 402
 Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
Salmonella migration into almonds, 397
- Alternaria*
Alternaria toxins in wheat, 1262
- Amidation
 Bactericidal effect of lactoferrin and its derivatives, 2468
- Amygdalin
 Enzyme immunoassay for detection for amygdalin, 2048
- Analysis
 Analysis of shellfish for *Cryptosporidium* and *Giardia*, 959
- Animal constituents
 Microscopic method for animal constituents detection in feedstuffs, 578
- Animal feed
 Developing group-specific primers, 2257
- Anisakis simplex*
 Allergen release by frozen and chilled *Anisakis*, 2578
Anisakis antigens in fish muscle, 1273
- Ani s 4
Anisakis antigens in fish muscle, 1273
- Anodic electrolyzed water
 Shelf life of coated and semifried tuna, 770
- Anthrax
 Inactivation of *B. anthracis* in milk, 333
- Antiadhesive
 Inhibition of pathogen adhesion, 2272
- Antibacterial
 Antibacterial activity of xanthorrhizol, 1926
- Antibacterial activity
 Influence of extraction methods on *N. sativa* essential oils, 2475
Q. infectoria as anti-*E. coli* agent, 1223
- Antibacterial enzymes
 Antibacterial milkfat globule membrane fractions, 126
- Antibacterial milk peptides
 Antibacterial milkfat globule membrane fractions, 126
- Antibacterial peptide
 Expression of goat lactoferricin from *E. coli*, 2523
- Antibiotic(s)
 Antibiotic-resistant bacteria in herbal products, 1486
 Antibiotics in milk, 1193
Salmonella prevalence in bovine lymph nodes, 1685
 Sulfamethazine water medication pharmacokinetics and contamination, 584
- Antibiotic residues
 Antibiotics in milk, 1193
 Distribution of drug residues in breast muscle tissues, 223
- Antibiotic resistance
 Antibiotic resistance in *V. parahaemolyticus*, 2552
 Enterococcal contamination of food by houseflies, 435
 Enterococci from human patients and foods, 2100
 Multidrug-resistant *Salmonella*, 27
 Regulation of safety of probiotics, 1734

- S. aureus* and Domiati cheese, 2567
- Antibiotic resistant
Inactivation of antibiotic-resistant *Campylobacter*, 1145
- Antifungal activity
Antifungal activity of *L. paracasei*, 2213
- Antigen-binding fragment (Fab)
Enzyme immunoassay for detection for amygdalin, 2048
- Antigenicity
Immunoreactivity of irradiated milk and whey samples, 1270
- Antiinfective
Inhibition of pathogen adhesion, 2272
- Antimicrobial(s)
Antimicrobial activity of *D. salina* extracts, 2138
Antimicrobial efficacy and sensory acceptability of plant essential oils, 1846
Cold plasma sanitizing of apples, 1357
Control of *L. monocytogenes* on frankfurters, 728
E. coli inhibition by colicin E1, 2519
Effects of interventions on MDR *Salmonella*, 2177
Enterococci in spontaneous sausage fermentation, 1244
Essential oils in coatings for fresh-cut apples, 1150
HPP in combination with antimicrobials, 820
Peracetic acid mixture as an antimicrobial treatment in poultry, 1119
Poultry skin flora after successive washings, 1707
- Antimicrobial activities
Natural antimicrobial combinations against *L. monocytogenes*, 2105
- Antimicrobial activity
Antibacterial activity of mushroom extracts, 1701
Antibacterial milkfat globule membrane fractions, 126
Antimicrobial activity of lactoferrin, 319
Antimicrobial essential oils against *L. monocytogenes* in pork, 502
- Antimicrobial effect
Antibacterial activity of dextran-conjugated lysozyme in cheese curd, 411
Antibacterial effects of polyphosphates, 1401
- Antimicrobial ice
Extending shelf life of fish, 218
- Antimicrobial polymer
Rechargeable antimicrobial polyethylene, 2042
- Antimicrobial resistance
Antimicrobial resistance and class 1 integrons in *E. coli*, 1679
Antimicrobial resistance and virulence in enterococci, 760
Antimicrobial susceptibility of *L. paracasei*, 339
- Antimycotic
Antimycotics and diacetate in RTE meats, 1386
- Antiyeast
Low-molecular-weight antiyeast metabolites of a protective culture, 2481
- Apple juice
Membrane damage and inactivation of bacteria in apple juice by RFEF, 684
- Apples
Mycotoxins produced in apples, 1714
Sanitizers for inactivation of *P. expansum* on apples, 643
- Arcobacter*
Isolation of *Arcobacter* from beef, 850
Prevalence and diversity of *Arcobacter* in the Czech Republic, 719
- Arcobacter* spp.
Arcobacter spp. in poultry processing, 2533
Susceptibility of *Arcobacter* spp. to aromatic compounds, 165
- Aromatic compounds
Susceptibility of *Arcobacter* spp. to aromatic compounds, 165
- Artificial neural networks
Predictive microbiology, 6
- Aseptic
Sterility testing of an aseptic dispensing valve, 653
- Aseptic filling
New sterilization process for aseptic packaging, 2119
- Aspergillus niger*
Sterility testing of an aseptic dispensing valve, 653
- Assay
Detection of *Salmonella* Enteritidis by polymixin-ELISA, 392
- Atomic force microscope
Stainless steel surface roughness and finish, 170
- ATP
Membrane damage and inactivation of bacteria in apple juice by RFEF, 684
Thermal inactivation of bacteria in liquid egg, 1988
- Attachment
Physicochemical properties of *Salmonella*, 2421
Resistance of *E. sakazakii* to desiccation, 940
- Audit
Potentially prohibited tissue removal, 573
- Autoinducer
Identification of AI-2 inhibitors from ground beef, 134
Role of quorum sensing in food spoilage, 1510
- Autoinducing peptide
Role of quorum sensing in food spoilage, 1510
- Autolysis
Ochratoxin A adsorption by yeast, 1496
- Automated microbiology
Automated MPN determination of *Enterobacteriaceae* in food, 376
- Automated rep-PCR
Evaluation of an automated rep-PCR system, 1372
- Avian influenza
Thermal inactivation of viruses in chicken meat, 1214
- Bacillus anthracis*
Efficacy of household products against *Bacillus* spores, 473
- Bacillus cereus*
B. cereus: enterotoxigenic and genetic profile, 2115
C. perfringens and *B. cereus* in seafood, 1178
Efficacy of household products against *Bacillus* spores, 473
Physiological response of *B. cereus* to food processing treatments, 2168
Survival of *B. cereus* during green olive fermentation, 1393
- Bacillus* spores
Bacillus spore inactivation differences, using two pressurizing fluids, 1186
- Bacteria
Antibacterial effects of polyphosphates, 1401
Antibiotic-resistant bacteria in herbal products, 1486
- Bacterial communities
16S rDNA profiling of alfalfa sprouts-associated bacteria, 200
- Bacterial transfer
Listeria transfer from knives to turkey breast, 176
- Bacterial translocation
Lactobacillus systemic immune functions and bacterial translocation, 796
- Bactericidal
Bactericidal effect of lactoferrin and its derivatives, 2468
- Bacteriocin
Bacteriocin-producing starters in fermented sausages, 1817
Bioprotective potential of LAB in malting and brewing, 1724
L. monocytogenes inactivation by AS-48, 2460
- Bacteriological quality
Incidence of *Salmonella* in imported seafood, 1460
- Bacteriology
Time course of infection with *Salmonella* Typhimurium in fattening pigs, 699
- Bacteriophage
Detection of *Salmonella* using a simple phage assay, 380
Fecal versus RAMS detection of *E. coli* O157:H7, 691
Toxin expression in O157 lineages, 1673
- Barley
Occurrence of bound deoxynivalenol in Fusarium-infected barley, 1266
- Barotolerance
Heat shock induces barotolerance in *L. monocytogenes*, 426
- BAX system
BAX system PCR method for *Salmonella* versus Brazil's official method, 2442

- Beef
 Combined treatments with irradiation on beef microbiology, 1237
 Danofloxacin in beef muscle, 1720
 Detection of *Sarcocystis* parasites in retail beef, 2144
E. coli O157:H7 and *Salmonella* in lairages, 1752
 Effect of incubation temperature on APCs, 373
 Effects of interventions on MDR *Salmonella*, 2177
 Heat resistance of *Salmonella* serotypes, 509
 Radiosensitivity of pathogens in ground beef, 516
Salmonella prevalence in bovine lymph nodes, 1685
 Xylitol to reduce microbial adhesion, 405
- Beef casings
 Lymphoid tissue reduction in bovine small intestines, 1199
- Beef cattle
 Shedding of foodborne pathogens by Florida cattle, 807
 Virulence genes in *E. coli* O157 and non-O157 isolates, 1774
- Beef head
 Antimicrobial interventions of bovine heads, 621
- Beef jerky
 Controlling pathogens in beef jerky, 918
- Beer
 Bioprotective potential of LAB in malting and brewing, 1724
- Behavioral theory
 Cancer patients and food safety, 1666
- Benzoic acid
 Microbiology of sweet potato puree, 639
- Beverage
 Ricin and abrin in beverages, 1875
- Bile
 Isolation of *C. jejuni* from bovine bile, 1228
- Bioassay
 Detection of PAHs in smoked meat, 993
 Role of quorum sensing in food spoilage, 1510
- Biocide
 Inactivation of *B. anthracis* in milk, 333
- Biodefense
 Thermal and hydrostatic inactivation, 196
- Biofilm(s)
 Fourteen-day-old chicken cecal microflora, 1981
Listeria transfer from knives to turkey breast, 176
L. monocytogenes biofilms and bacteriocins, 634
 Resistance of *E. sakazakii* to desiccation, 940
 Stainless steel surface roughness and finish, 170
 Temperature and nutrient effects on *C. jejuni*, 271
- Biogenic amines
 Decarboxylating strains in a typical Italian cheese, 205
 Enterococci in spontaneous sausage fermentation, 1244
M. morgani ornithine decarboxylase, 657
 Safety and quality assessment of lagoon grey mullets, 2572
- Bioinformatics
 Detection and genotyping noroviruses using DNA microarrays, 1434
- Biological applications of polymers
 Rechargeable antimicrobial polyethylene, 2042
- Bioluminescence
 Detection of *Salmonella* using a simple phage assay, 380
- Biopreservation
 Low-molecular-weight antiyeast metabolites of a protective culture, 2481
- Bioprotection
 Bioprotective potential of LAB in malting and brewing, 1724
- Bivalve
 Effect of HPP on freely suspended and bivalve-associated T7, 345
- Black pepper
 Decontamination of *E. coli* with ozone, 914
- Blade
 Blade tenderization of beef containing *E. coli*, 2190
- Blanching
L. monocytogenes in produce processing, 735
- Blood
 Competitive ELISA to detect bovine blood, 1000
- Blueberries
 Heat-resistant fungi from blueberries, 2030
- Bluefin tuna
 Mercury in bluefin tuna, 595
- Body weight gain
 Effects of a probiotic culture on performance of weaned piglets, 1797
- Bologna
 Survival of *L. monocytogenes* in a gastrointestinal model, 2014
- Boneless meat
 Microbiological quality of chicken meat, 1123
- Bottle
 New sterilization process for aseptic packaging, 2119
- Bound mycotoxins
 Occurrence of bound deoxynivalenol in Fusarium-infected barley, 1266
- Bovine
 Competitive ELISA to detect bovine blood, 1000
 Non-O157 STEC from humans, cattle, and food in the United States, 1023
- Bovine DNA quantification
 Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
- Bovine spongiform encephalopathy (BSE)
 A comparison of BSE risk between Japanese and U.S. beef, 802
 Cross-contamination of CNS tissue due to beef carcass splitting, 83
 Detection of central nervous system tissues in meat products, 2059
 Developing group-specific primers, 2257
 Lymphoid tissue reduction in bovine small intestines, 1199
 Potentially prohibited tissue removal, 573
- Brazil nuts
 Microbiology of raw nuts, 402
- Breaded
 Salmonellosis from stuffed chicken products, 2153
- Broiler(s)
Campylobacter on retail broiler carcasses in New Zealand, 2526
L. monocytogenes in poultry flocks, 1996
- Broiler flocks
Campylobacter in thinning of broiler flocks, 264
- Buffered sodium citrate
 Control of *C. perfringens* in Tajik sambusa, 77
- Caco-2 cell
 In vitro digestion and bioavailability of LDPs, 979
- Cadmium
 Linking soil quality to human cadmium exposure, 2504
- Calcium chloride
 Effects of chloride salts on microbial growth, 1412
- Campylobacter*
 Antibiotic resistance of *Campylobacter* isolates from Poland, 602
Campylobacter in paired turkey and swine farms, 1791
Campylobacter in thinning of broiler flocks, 264
Campylobacter on retail broiler carcasses in New Zealand, 2526
Campylobacter prevalence in poultry production, 2295
Campylobacter spp. in raw and ready-to-eat foods, 2087
 Detection methods for *Campylobacter* spp., 835
 Detection of *Campylobacter* spp., 2543
 Economics of *Campylobacter* control in poultry, 479
 Peracetic acid mixture as an antimicrobial treatment in poultry, 1119
 Shedding of foodborne pathogens by Florida cattle, 807
- Campylobacter coli*
Campylobacter in paired turkey and swine farms, 1791
- Campylobacter jejuni*
Campylobacter in paired turkey and swine farms, 1791
C. jejuni cross-contamination, 1018
 Contamination of kitchen by *C. jejuni* in raw vegetables, 2448
 Inactivation of antibiotic-resistant *Campylobacter*, 1145
 Isolation of *C. jejuni* from bovine bile, 1228
 Temperature and nutrient effects on *C. jejuni*, 271
- Cancer patients
 Cancer patients and food safety, 1666
- Candida*
 Yeasts from refrigerated commercial shell eggs, 1258

- Cantaloupe(s)
 Identification of *Salmonella* serotypes by PCR-RFLP, 2217
 Microflora on cantaloupes, 46
- Carbon dioxide
 Antimicrobial activity of *D. salina* extracts, 2138
Y. enterocolitica in ground pork packaged under modified atmospheres, 845
- Carbon monoxide
 Spoilage and safety characteristics of ground beef, 293
- Carcass(es)
E. coli inhibition by colicin E1, 2519
E. coli O157:H7 and *Salmonella* in lairages, 1752
 Effect of incubation temperature on APCs, 373
 Xylitol to reduce microbial adhesion, 405
- Carcass mapping
 Mapping of *Salmonella* contamination on the poultry carcass, 1785
- Carcass maturity score
 A comparison of BSE risk between Japanese and U.S. beef, 802
- Carcass prevalence
E. coli O157 in feces and on beef carcasses, 1761
- Carvacrol
 Facilitated thermal inactivation of *E. coli* O157:H7, 1604
 Inactivation of antibiotic-resistant *Campylobacter*, 1145
T. vulgaris and a stored food mite, 351
- Cashews
 Microbiology of raw nuts, 402
- Catalonia, Spain
 Dietary intake of hexachlorobenzene, 2148
- Catering
 Foodborne disease outbreaks in the catering industry, 550
- Catfish
 Radiation D_{10} -values of *L. monocytogenes* on catfish and tilapia, 2278
- Cattle
 Dust exposure and pathogen increases on cattle hides, 2078
E. coli O157:H7 and *Salmonella* in lairages, 1752
 Effect of transportation and lairage on risk of *S. enterica* in cattle, 2228
 Fecal versus RAMS detection of *E. coli* O157:H7, 691
 Impact of transportation and lairage on *E. coli* O157:H7 in cattle, 1114
 Isolation of *C. jejuni* from bovine bile, 1228
 Lymphoid tissue reduction in bovine small intestines, 1199
Salmonella prevalence in bovine lymph nodes, 1685
- Cattle feces
 Detection of Shiga toxin-producing *E. coli*, 927
- Cecal microflora
 Fourteen-day-old chicken cecal microflora, 1981
- Celiac disease
 Approaches to establish thresholds, 1043
Lactobacillus for gluten and phytate hydrolysis in gluten-free bread, 1491
- Cell-free translation assay
 Detection of risin in foods, 2053
- Cell hydrophobicity
 Altered cell composition in stress-adapted *L. innocua*, 182
- Cell signaling
 Identification of AI-2 inhibitors from ground beef, 134
- Cellulose-containing biofilm
 Cellulose production in *E. sakazakii* biofilms, 13
- Central nervous system
 Cross-contamination of CNS tissue due to beef carcass splitting, 83
 Detection of central nervous system tissues in meat products, 2059
- Central nervous system tissue
 Comparison of detection methods for CNST, 1505
- Cetrimide
 Altered cell composition in stress-adapted *L. innocua*, 182
- Chain modeling
 Linking soil quality to human cadmium exposure, 2504
- Cheese
 Antibacterial activity of dextran-conjugated lysozyme in cheese curd, 411
Listeria spp. in farmstead cheese processing environments, 2239
 Pathogen incidence in raw milk, 1580
 Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
- Chemical-activated luciferase expression assay (CALUX)
 Detection of PAHs in smoked meat, 993
- Chemical composition
 Influence of extraction methods on *N. sativa* essential oils, 2475
- Chemical dehairing
 Reduction of hide microbial contamination, 2223
- Chemical oxygen demand
 Impact of water reuse in fresh-cut washing, 2514
- Chicken(s)
 Antibiotic resistance of *Campylobacter* isolates from Poland, 602
 Bacteriology of raw chicken nuggets, 613
C. jejuni cross-contamination, 1018
 Distribution of drug residues in breast muscle tissues, 223
 Evaluation of a copper sulfate-based commercial disinfectant, 226
 Fourteen-day-old chicken cecal microflora, 1981
 Risk assessment of *Listeria* in ready-to-eat chicken production line, 946
 Salmonellosis from stuffed chicken products, 2153
 Virulence genes in *E. coli* O157 and non-O157 isolates, 1774
- Chicken frankfurters
 Model for growth of *Salmonella* on chicken frankfurters, 1135
- Chicken meat
 Economics of *Campylobacter* control in poultry, 479
- Chile pepper
 Identification of *Salmonella* serotypes by PCR-RFLP, 2217
- Chilling
 Allergen release by frozen and chilled *Anisakis*, 2578
 Application of meta-analysis in food safety, 1330
- China
 STEC and *E. coli* O157 from retail meats in China, 706
- Chitosan film
 Antimicrobial activity of lactoferrin, 319
- Chlorinated wash water
 Microflora on cantaloupes, 46
- Chlorine dioxide
 Parasite inactivation by chlorine dioxide, 2410
- Cinnamaldehyde
 Facilitated thermal inactivation of *E. coli* O157:H7, 1604
 Inactivation of antibiotic-resistant *Campylobacter*, 1145
 Susceptibility of *Arcobacter* spp. to aromatic compounds, 165
- Cinnamon
 Combined treatments with irradiation on beef microbiology, 1237
- Citrinin
 Mycotoxins produced in apples, 1714
- Class 1 integrons
 Antimicrobial resistance and class 1 integrons in *E. coli*, 1679
- Cleaning
 Cleaning and validation to prevent allergen cross-contact, 445
 Risk assessment of *Listeria* in ready-to-eat chicken production line, 946
- Cleaning protocols
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
- Clenbuterol residue
 Screening clenbuterol by strip and ELISA, 865
- Clostridium botulinum*
 Proteolytic *C. botulinum* in rice at low pH, 468
- Clostridium perfringens*
 Control of *C. perfringens* in Tajik sambusa, 77
C. perfringens and *B. cereus* in seafood, 1178
- Clostridium sporogenes*
 Heat tolerance of *C. sporogenes* spores, 953
- Coagulase type
 Characteristics of enterotoxin H-producing *S. aureus*, 1855
- Cocoa
 OTA removal procedure for cocoa shells, 1422
- Cold
 Altered cell composition in stress-adapted *L. innocua*, 182
L. monocytogenes growth at low temperature, 420
- Cold atmospheric gas plasma

- Decontamination of mango and melon by gas plasma, 302
 Plasma inactivation of bacteria migrating through fruits, 1619
- Cold shock
 In vitro cold shock gene expression in *Vibrio vulnificus*, 157
- Colicin E1
E. coli inhibition by colicin E1, 2519
- Coliform bacilli
 Isolation of *C. jejuni* from bovine bile, 1228
- Coliform counts
 Effects of a probiotic culture on performance of weaned piglets, 1797
- Coliform flora
 Tracing the origin of coliform bacteria, 1910
- Coliforms
 Coliforms and foodborne pathogens on spinach in packing plants, 2398
 Poultry skin flora after successive washings, 1707
- Colloidal gold
 Enhanced rapidity for *L. monocytogenes* detection, 781
- Colorimetric method
 Alkaline phosphates in cow's, goat's, and sheep's milk, 1884
- Commensal *E. coli*
 Antimicrobial resistance and class 1 integrons in *E. coli*, 1679
- Commercialization conditions
 Microbial quality in mineral waters, 1253
- Comminuted fermented meats
 Toxoplasma in ready-to-eat smallgoods, 1442
- Comparison
 Isolation of *Arcobacter* from beef, 850
- Competitive enzyme-linked immunosorbent assay
 Competitive ELISA to detect bovine blood, 1000
 OTA determination in grapes, 2488
- Compositions
 HPP inactivation of FCV, 1598
- Compost
 Determining *E. coli* O157:H7 levels in manures and soil, 2233
- Consumer education
 Cooking parameters for safe seafood for consumers, 1287
- Consumer food safety
 Consumer food safety knowledge and behaviors, 1651
- Consumers
 Consumer assessment of safety/date labeling on RTE meat products, 70
Listeria and *Enterobacteriaceae* in domestic refrigerators, 608
- Contamination
 Bacteriological safety in São Jorge cheese manufacture, 986
 Contamination of carcasses with *Salmonella* during poultry slaughter, 146
 Factors for produce contamination, 2389
 Incidence of *Salmonella* in imported seafood, 1460
Salmonella contamination of food service eggs, 19
 Sources of contamination and pathogen excretion from infected persons, 2582
 Sulfamethazine water medication pharmacokinetics and contamination, 584
- Continuous-flow culture
 Fourteen-day-old chicken cecal microflora, 1981
- Conventional culture
 Detection methods for *Campylobacter* spp., 835
- Cookies
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
- Cooking
 Heat resistance of *Salmonella* serotypes, 509
 Toxoplasma in ready-to-eat smallgoods, 1442
- Cooking protocols
 Cooking parameters for safe seafood for consumers, 1287
- Copper sulfate
 Evaluation of a copper sulfate-based commercial disinfectant, 226
- Cornish game hen
 Mapping of *Salmonella* contamination on the poultry carcass, 1785
- Corylus avellana*
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
- Cost
 Application of MCDA to a food safety decision-making problem, 2323
- Cost-benefit analysis
 Economics of *Campylobacter* control in poultry, 479
- Cow's milk
 Immunoreactivity of irradiated milk and whey samples, 1270
- Cross-contact
 Cleaning and validation to prevent allergen cross-contact, 445
- Cross-contamination
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
C. jejuni cross-contamination, 1018
 Contamination of kitchen by *C. jejuni* in raw vegetables, 2448
- Cross-protection
 Contribution of sigma B factor in alkali tolerance in *L. monocytogenes*, 1481
- Cross-reactivity
 Latex glove use by food handlers, 2334
- Cryptosporidium*
 Analysis of shellfish for *Cryptosporidium* and *Giardia*, 959
 Parasite inactivation by chlorine dioxide, 2410
- Cull sow
L. monocytogenes in cull sows and pork, 545
- Curcuma xanthorrhiza* Roxb.
 Antibacterial activity of xanthorrhizol, 1926
- Cyanide poisoning
 Enzyme immunoassay for detection for amygdalin, 2048
- Cyclopiazonic acid
 Efficient fungal DNA extraction method for RAPD-PCR, 2497
- Cyclospora*
 Parasite inactivation by chlorine dioxide, 2410
- Cytokine
Lactobacillus systemic immune functions and bacterial translocation, 796
- Dairy
 Inactivation of *B. anthracis* in milk, 333
 Predicting *Salmonella* prevalence in a dairy herd, 1967
- Dairy cattle
 Multidrug-resistant *Salmonella*, 27
- Dairy farming
 Food safety for the Turkish dairy industry, 440
- Dairy industry
 Food safety for the Turkish dairy industry, 440
- Dairy products
 Survival of *L. innocua* in dairy products, 742
 Survival of *L. monocytogenes*, 1835
- Danofloxacin
 Danofloxacin in beef muscle, 1720
- Date labels
 Consumer assessment of safety/date labeling on RTE meat products, 70
- Dates
 Microbial spoilage of rutab, 1406
- Death kinetics
 Bacterial reduction in UV irradiated chill brines, 629
- Decarboxylating strains
 Decarboxylating strains in a typical Italian cheese, 205
- Decimal reduction time
 Storage stability of freeze-dried lactic acid bacteria, 431
- Decontamination
 Contamination of kitchen by *C. jejuni* in raw vegetables, 2448
 Decontamination of almonds with infrared heat, 897
 Effect of different water temperatures on knife decontamination, 1338
 OTA removal procedure for cocoa shells, 1422
 Survival of murine norovirus 1 on onions and spinach, 1590
 UV light-hydrogen peroxide decontamination of lettuce, 903
 Water disinfection from the fresh-cut vegetable industry, 286
- Dense phase carbon dioxide

- Dense phase CO₂ processing of spinach, 1015
- Deoxynivalenol
Deoxynivalenol contamination in wheat products, 1931
LC-UV-MS method for DON, DON-3-glucoside, and pigment, 1205
Occurrence of bound deoxynivalenol in Fusarium-infected barley, 1266
- Deoxynivalenol-3-glucoside
LC-UV-MS method for DON, DON-3-glucoside, and pigment, 1205
- Descriptive analysis
Risk factors for food premises unsatisfactory inspection results, 1632
- Desiccation
Inactivation of desiccated *E. sakazakii*, 934
Resistance of *E. sakazakii* to desiccation, 940
- Detection
Detection and genotyping noroviruses using DNA microarrays, 1434
Detection of abrin, 1868
Detection of *Salmonella* using a simple phage assay, 380
Listeria spp. in farmstead cheese processing environments, 2239
- Detection method(s)
Detection of Shiga toxin-producing *E. coli*, 927
Salmonella detection methods on turkey parts, 386
- Detection of *Salmonella*
BAX system PCR method for *Salmonella* versus Brazil's official method, 2442
- Devriesia* spp.
Heat-resistant fungi from blueberries, 2030
- Dextran
Antibacterial activity of dextran-conjugated lysozyme in cheese curd, 411
- Dietary exposure
Dietary intake of hexachlorobenzene, 2148
- Differentiation
Arcobacter spp. in poultry processing, 2533
Salmonella identification and strain differentiation by HGMP-IR, 2249
- Direct-fed microbial
Probiotics effect on *E. coli* O157:H7 and *Salmonella* shedding in cattle, 539
- Discrimination
Distinguishable control strains for food microbiology testing, 2301
SSCP analysis for rapid discrimination of *Salmonella* isolates, 1960
- Disease control
Fludioxonil residues and control of blue and gray mold on pears, 967
- Disease management
Potassium sorbate for controlling postharvest brown rot, 1626
- Disinfectant(s)
Efficacy of household products against *Bacillus* spores, 473
Electrolyzed water in the food industry, 1934
- Dispensing valve
Sterility testing of an aseptic dispensing valve, 653
- Distribution
Oxytetracycline residue depletion in giant freshwater prawn, 870
- DNA barcoding
DNA barcoding and the *Regulatory Fish Encyclopedia*, 210
- DNA extraction
DNA extraction for PCR detection of food pathogens, 2110
Efficient fungal DNA extraction method for RAPD-PCR, 2497
- Domestic kitchen
Contamination of kitchen by *C. jejuni* in raw vegetables, 2448
- Domestic refrigerator
Domestic refrigeration practices, 1898
- Domiat cheese
S. aureus and Domiat cheese, 2567
- Dorsal root ganglia
Detection of DRG in advanced meat recovery product, 2307
- Double separatory funnel
Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
- Drain
L. monocytogenes in drain pipes, 66
- Drop plating
Mapping of *Salmonella* contamination on the poultry carcass, 1785
- Drug resistant
Heat resistance of *Salmonella* serotypes, 509
- Dry-cured ham
E-beam radiation of dry-cured ham, 2001
- Dry cured salami
E. coli O157:H7 inhibition in dry sausage by mustard powder, 486
- Dry fermented sausage
L. monocytogenes on salami during storage, 494
Predictive microbiology, 6
- Dry sausage
Survival of *L. monocytogenes* in dry sausage, 1550
- D₁₀-value
Inactivation of desiccated *E. sakazakii*, 934
- Dunaliella salina*
Antimicrobial activity of *D. salina* extracts, 2138
- Dust
Dust exposure and pathogen increases on cattle hides, 2078
- D-value
Heat-resistant fungi from blueberries, 2030
Modification of the submerged coil, 775
- Dynamic modeling
Modeling *L. monocytogenes* in pasteurized vanilla cream, 1828
- eaeA*
Prevalence of *E. coli* O157:H7 in vegetables, 816
- E-beam radiation
E-beam radiation of dry-cured ham, 2001
- Effectiveness
Application of MCDA to a food safety decision-making problem, 2323
- Egg(s)
Detection of *Salmonella* Enteritidis by polymixin-ELISA, 392
Salmonella contamination of food service eggs, 19
Salmonella Enteritidis growth on egg yolk membranes, 1905
Salmonella Enteritidis in eggs, 2073
- Egg safety
Salmonella Enteritidis in eggs, 2073
- 18S rRNA
Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
- Electrochemiluminescence
Detection of abrin, 1868
- Electrolyzed oxidizing water
Economics of *Campylobacter* control in poultry, 479
- Electrolyzed water
Electrolyzed water in the food industry, 1934
- Electron beam radiation
E-beam radiation for reducing pathogens in spinach, 2415
- Electrostatic
Virus adsorption to lettuce, 522
- Emetic toxin
C. perfringens and *B. cereus* in seafood, 1178
- Enrichment
E. coli O157:H7 enrichment in produce, 139
- Enrofloxacin
Distribution of drug residues in breast muscle tissues, 223
- Enteric virus(es)
UV light-hydrogen peroxide decontamination of lettuce, 903
Viral contamination in clams, 1689
Virus adsorption to lettuce, 522
- Enterobacteriaceae*
Automated MPN determination of *Enterobacteriaceae* in food, 376
Bacteriological safety in São Jorge cheese manufacture, 986
Hygiene indicator criteria for meat in Belgium, 35
Listeria and *Enterobacteriaceae* in domestic refrigerators, 608
- Enterobacter sakazakii*
Cellulose production in *E. sakazakii* biofilms, 13
Evaluation of an automated rep-PCR system, 1372
Inactivation of desiccated *E. sakazakii*, 934
Resistance of *E. sakazakii* to desiccation, 940
- Enterococci
Enterococcal contamination of food by houseflies, 435
Enterococci in spontaneous sausage fermentation, 1244

- L. monocytogenes* biofilms and bacteriocins, 634
- Enterococcus*
- Enterococci from human patients and foods, 2100
- Enterococcus faecalis*
- Antimicrobial resistance and virulence in enterococci, 760
- Enterohemorrhagic *Escherichia coli* (EHEC)
- Occurrence of STEC in meat products in Morocco, 2082
 - Transcription of *stx*₁, *marA*, and *eaeA* in *E. coli* O157:H7, 1469
- Enterotoxin
- C. perfringens* and *B. cereus* in seafood, 1178
 - Detection of *S. aureus*, 191
- Enterotoxin genes
- Multiplex PCR detection of enterotoxin genes in *Aeromonas* spp., 2094
 - S. aureus* and Domiat cheese, 2567
- Enterotoxin production
- B. cereus*: enterotoxigenic and genetic profile, 2115
- Enumeration
- Automated MPN determination of *Enterobacteriaceae* in food, 376
 - Determining *E. coli* O157:H7 levels in manures and soil, 2233
- Enumeration method
- Mapping of *Salmonella* contamination on the poultry carcass, 1785
- Environmental sampling
- Listeria* spp. in farmstead cheese processing environments, 2239
- Enzyme immunoassay
- Enzyme immunoassay for detection for amygdalin, 2048
- Enzyme-linked immunosorbent assay (ELISA)
- Allergen cross-contamination in the manufacture of fine bakery wares, 2263
 - Characteristics of enterotoxin H–producing *S. aureus*, 1855
 - Comparison of detection methods for CNST, 1505
 - Cross-contamination of CNS tissue due to beef carcass splitting, 83
 - Detection of abrin, 1868
 - Detection of central nervous system tissues in meat products, 2059
 - Detection of melamine, 590
 - Enhanced rapidity for *L. monocytogenes* detection, 781
 - Screening clenbuterol by strip and ELISA, 865
- Epidemic clones
- Epidemiologic relevance of MVLST, 2559
- Epidemiologic concordance
- Epidemiologic relevance of MVLST, 2559
- Epidemiology
- Factors for produce contamination, 2389
- Escherichia coli*
- Antimicrobial resistance in poultry bacteria, 2537
 - Bacteriology of raw chicken nuggets, 613
 - Coliforms and foodborne pathogens on spinach in packing plants, 2398
 - Decontamination of *E. coli* with ozone, 914
 - Dense phase CO₂ processing of spinach, 1015
 - Detection of Shiga toxin–producing *E. coli*, 927
 - E. coli* inhibition by colicin E1, 2519
 - Hygiene indicator criteria for meat in Belgium, 35
 - Inhibition of pathogen adhesion, 2272
 - Membrane damage and inactivation of bacteria in apple juice by RFEF, 684
 - Microbiological profile of hydroponic tomatoes, 60
 - Microbiology of raw nuts, 402
 - On-farm meat safety, 1919
 - Radiosensitivity of pathogens in ground beef, 516
 - Risk assessment for STEC in unpasteurized milk, 257
 - Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
 - Salmonella* on hydroponic tomatoes, 676
 - Surrogates for *Salmonella* in meat, 714
 - Thermal inactivation of bacteria in liquid egg, 1988
 - Toxin expression in O157 lineages, 1673
 - Water disinfection from the fresh-cut vegetable industry, 286
- Escherichia coli* O157
- Dust exposure and pathogen increases on cattle hides, 2078
 - E. coli* O157 in feces and on beef carcasses, 1761
 - Impact of transportation and lairage on *E. coli* O157:H7 in cattle, 1114
 - Killing *E. coli* O157 with high pressure processing, 2182
 - Survival of *E. coli* O157 on lettuce, 2384
 - Virulence genes in *E. coli* O157 and non-O157 isolates, 1774
- Escherichia coli* O157:H7
- Antimicrobial effect of muscadine seed extracts, 1465
 - Antimicrobial interventions for leafy greens, 625
 - Antimicrobial interventions of bovine heads, 621
 - Antimicrobial synergy between high pressure and Red No. 3, 1861
 - Blade tenderization of beef containing *E. coli*, 2190
 - Cold plasma sanitizing of apples, 1357
 - Controlling pathogens in beef jerky, 918
 - Determining *E. coli* O157:H7 levels in manures and soil, 2233
 - Distinguishable control strains for food microbiology testing, 2301
 - E-beam radiation for reducing pathogens in spinach, 2415
 - E. coli* O157:H7 and *Salmonella* in lairages, 1752
 - E. coli* O157:H7 enrichment in produce, 139
 - E. coli* O157:H7 inhibition in dry sausage by mustard powder, 486
 - Effects of interventions on MDR *Salmonella*, 2177
 - Effects of reduced enrichment volumes, 1768
 - Essential oils in coatings for fresh-cut apples, 1150
 - Facilitated thermal inactivation of *E. coli* O157:H7, 1604
 - Fecal pats inoculated with *E. coli* O157:H7 to mimic a super shedder, 648
 - Fecal versus RAMS detection of *E. coli* O157:H7, 691
 - Heat inactivation of *E. coli* O157:H7 in nonintact beef, 1349
 - Hot water treatment decontamination of mung bean seeds, 830
 - HPP in combination with antimicrobials, 820
 - Multiple-cycle high-pressure treatments, 811
 - Pathogen incidence in raw milk, 1580
 - Predictive microbiology, 6
 - Prevalence of *E. coli* O157:H7 in vegetables, 816
 - Probiotics effect on *E. coli* O157:H7 and *Salmonella* shedding in cattle, 539
 - Q. infectoria* as anti-*E. coli* agent, 1223
 - Reduction of hide microbial contamination, 2223
 - Reduction of microbial contamination on cattle hides, 1343
 - Shedding of foodborne pathogens by Florida cattle, 807
 - Spoilage and safety characteristics of ground beef, 293
 - Survival of *E. coli* O157:H7 in acid, 2404
 - Transfer of *E. coli* O157:H7 to lettuce from ice, 252
- Essential oil(s)
- Antimicrobial efficacy and sensory acceptability of plant essential oils, 1846
 - Antimicrobial essential oils against *L. monocytogenes* in pork, 502
 - Combined treatments with irradiation on beef microbiology, 1237
 - Essential oils in coatings for fresh-cut apples, 1150
 - Influence of extraction methods on *N. sativa* essential oils, 2475
 - Radiosensitivity of pathogens in ground beef, 516
 - Shelf life of coated and semifried tuna, 770
- Ethanol shock
- Protein expression in stressed *V. parahaemolyticus*, 2589
- Expansolides A and B
- Mycotoxins produced in apples, 1714
- Experimental infection
- Time course of infection with *Salmonella* Typhimurium in fattening pigs, 699
- Exposure
- Microbiological quality of chicken meat, 1123
- Exposure assessment
- Linking soil quality to human cadmium exposure, 2504
- Extraction
- OTA determination in grapes, 2488
- Extrinsic and intrinsic factors
- Y. enterocolitica* in ground pork packaged under modified atmospheres, 845
- Extrusion
- Reduced toxicity of fumonisin B₁, 2036
- Farmstead
- Pathogen incidence in raw milk, 1580

- Fatty acid composition
 Acidified sodium chlorite treatment of trout fillets, 973
 Altered cell composition in stress-adapted *L. innocua*, 182
- Fatty acids
 Identification of AI-2 inhibitors from ground beef, 134
 Safety and quality assessment of lagoon grey mullets, 2572
- FD&C Red No. 3
 Antimicrobial synergy between high pressure and Red No. 3, 1861
- Fecal-oral route
 Sources of contamination and pathogen excretion from infected persons, 2582
- Fecal pats
 Fecal pats inoculated with *E. coli* O157:H7 to mimic a super shedder, 648
- Fecal prevalence
E. coli O157 in feces and on beef carcasses, 1761
- Fecal shedding
 Probiotics effect on *E. coli* O157:H7 and *Salmonella* shedding in cattle, 539
- Feed conversion efficiency
 Effects of a probiotic culture on performance of weaned piglets, 1797
- Feedlot cattle
 Probiotics effect on *E. coli* O157:H7 and *Salmonella* shedding in cattle, 539
- Feedstuffs
 Microscopic method for animal constituents detection in feedstuffs, 578
 Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
- Feline calicivirus
 HPP inactivation of FCV, 1598
- Fermentation
 Effects of chloride salts on microbial growth, 1412
- Fermented meat safety
E. coli O157:H7 inhibition in dry sausage by mustard powder, 486
- Fermented vegetable product
 Growth inhibitory effects of kimchi against foodborne pathogens, 325
- Feta cheese
 Survival of *L. innocua* in dairy products, 742
- Figs
 LC-TOF-MS analysis of mycotoxins in figs, 1500
- Fimbriae
 PCR detection of *Salmonella* by *stdA* primers, 1108
- Finfish
 Radiation D_{10} -values of *L. monocytogenes* on catfish and tilapia, 2278
- Fish
 Extending shelf life of fish, 218
 Microbial quality of fish fillets, 1544
- Fish culture
 Mercury in bluefin tuna, 595
- Fish muscle
Anisakis antigens in fish muscle, 1273
- fliC_{H7}*
 Prevalence of *E. coli* O157:H7 in vegetables, 816
- Flocks
Campylobacter prevalence in poultry production, 2295
- Flow cytometry
 Physiological response of *B. cereus* to food processing treatments, 2168
- Fluctuating temperature
 Modeling *L. monocytogenes* in pasteurized vanilla cream, 1828
- Fludioxonil
 Fludioxonil residues and control of blue and gray mold on pears, 967
- Fluorescence
 Danofloxacin in beef muscle, 1720
- Fluorophos method
 Alkaline phosphates in cow's, goat's, and sheep's milk, 1884
- Fluoroquinolone
 Distribution of drug residues in breast muscle tissues, 223
- Focus groups
 Cancer patients and food safety, 1666
- Food
 Detection of abrin, 1868
 Detection of *Salmonella* by PCR in foods, 2436
 Dietary intake of hexachlorobenzene, 2148
- Food adulteration
 Efficacy of drug test kits for adulterated food, 1007
- Food allergy
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
 Approaches to establish thresholds, 1043
- Foodborne
 Non-O157 STEC from humans, cattle, and food in the United States, 1023
- Foodborne disease outbreaks
 Foodborne disease outbreaks in the catering industry, 550
 Operational practices associated with foodborne disease outbreaks, 1659
- Foodborne outbreaks
 Infective doses and pathogen carriage, 2339
 Sources of contamination and pathogen excretion from infected persons, 2582
 U.S. juice-associated outbreaks, 356
- Foodborne pathogen(s)
 Antibacterial activity of xanthorrhizol, 1926
 DNA extraction for PCR detection of food pathogens, 2110
 Growth inhibitory effects of kimchi against foodborne pathogens, 325
Listeria and *Enterobacteriaceae* in domestic refrigerators, 608
 Metabolic effects of *Fusarium*, 1366
 Water disinfection from the fresh-cut vegetable industry, 286
- Food contact surfaces
L. monocytogenes adhesion and viability, 1379
- Food defense
 Efficacy of household products against *Bacillus* spores, 473
- Food ecology
 Role of quorum sensing in food spoilage, 1510
- Food handling behaviors
 Consumer food safety knowledge and behaviors, 1651
- Food migration tests
 Role of fat content in trapping volatile compounds, 1889
- FoodNet
 Food use and practices in long-term care facilities, 365
- Food packaging
 Role of fat content in trapping volatile compounds, 1889
- Food pathogens
 Antibacterial milkfat globule membrane fractions, 126
- Food preparation
 Operational practices associated with foodborne disease outbreaks, 1659
- Food products
 Electrolyzed water in the food industry, 1934
- Food quality
 Food safety for the Turkish dairy industry, 440
- Food quality index
M. morgani ornithine decarboxylase, 657
- Food safety
 Application of MCDA to a food safety decision-making problem, 2323
 Blade tenderization of beef containing *E. coli*, 2190
 Controlling pathogens in beef jerky, 918
 Detection of central nervous system tissues in meat products, 2059
 Detection of melamine, 590
 Detection of *Sarcocystis* parasites in retail beef, 2144
 Detection of Shiga toxin-producing *E. coli*, 927
 Domestic refrigeration practices, 1898
 E-beam radiation for reducing pathogens in spinach, 2415
 Enterococcal contamination of food by houseflies, 435
 Foodborne disease outbreaks in the catering industry, 550
 Food safety for the Turkish dairy industry, 440
L. monocytogenes in cull sows and pork, 545
L. monocytogenes on salami during storage, 494
 Mercury in bluefin tuna, 595
 Operational practices associated with foodborne disease outbreaks, 1659
 Potentially prohibited tissue removal, 573
 Rapid method used to analyze Ochratoxin A, 2133
 Risk factors for food premises unsatisfactory inspection results, 1632

- Salmonella contamination of food service eggs, 19
- Sandwiches from health care establishments, 309
- Survival of *E. coli* O157 on lettuce, 2384
- U.S. FDA regulatory management of human listeriosis, 1277
- Food safety education
 - Cancer patients and food safety, 1666
- Food safety knowledge
 - Consumer food safety knowledge and behaviors, 1651
- Food service
 - Campylobacter* spp. in raw and ready-to-eat foods, 2087
 - Salmonella contamination of food service eggs, 19
 - Sterility testing of an aseptic dispensing valve, 653
- Food spoilage
 - Role of quorum sensing in food spoilage, 1510
- Food supplements
 - Regulation of safety of probiotics, 1734
- Food suppliers
 - Operational practices associated with foodborne disease outbreaks, 1659
- Food workers
 - Infective doses and pathogen carriage, 2339
 - Sources of contamination and pathogen excretion from infected persons, 2582
- Forage
 - On-farm meat safety, 1919
- Francisella tularensis*
 - Thermal resistance of *F. tularensis*, 2208
- Frankfurters
 - Control of *L. monocytogenes* on frankfurters, 728
 - Estimating *L. monocytogenes* growth kinetics parameters in frankfurters, 1806
- Freezing
 - Allergen release by frozen and chilled *Anisakis*, 2578
 - Survival of murine norovirus 1 on onions and spinach, 1590
- Fresh bagged salad
 - E-beam radiation for reducing pathogens in spinach, 2415
- Fresh-cut apple
 - Essential oils in coatings for fresh-cut apples, 1150
- Fresh produce
 - Cold plasma sanitizing of apples, 1357
- Fresh vegetables
 - DNA extraction for PCR detection of food pathogens, 2110
- Frozen
 - Salmonellosis from stuffed chicken products, 2153
- Fruit juice
 - L. monocytogenes* inactivation by AS-48, 2460
 - Thermal resistance of *F. tularensis*, 2208
- Fruits
 - Factors for produce contamination, 2389
 - L. monocytogenes* inactivation by AS-48, 2460
- Fumonisin
 - LC-TOF-MS analysis of mycotoxins in figs, 1500
 - Reduced toxicity of fumonisin B₁, 2036
- Fungi
 - Survival of *L. innocua* in dairy products, 742
- Fungicide treatments
 - Fludioxonil residues and control of blue and gray mold on pears, 967
- Furanosyl borate diester
 - Role of quorum sensing in food spoilage, 1510
- Fusarium head blight
 - Occurrence of bound deoxynivalenol in Fusarium-infected barley, 1266
- Fusarium* species
 - Antifungal activity of *L. paracasei*, 2213
 - Metabolic effects of *Fusarium*, 1366
- Gamma radiation
 - Immunoreactivity of irradiated milk and whey samples, 1270
 - Inactivation of desiccated *E. sakazakii*, 934
- Gaping
 - Icing to control *V. vulnificus* and *V. parahaemolyticus*, 1475
- Gas chromatography-mass spectrometry (GC-MS)
 - Antimicrobial activity of *D. salina* extracts, 2138
- Gastroenteritis
 - Antibiotic resistance in *V. parahaemolyticus*, 2552
- Gastrointestinal model
 - Survival of *L. monocytogenes* in a gastrointestinal model, 2014
- Gene expression
 - Transcription of *stx*₁, *marA*, and *eaeA* in *E. coli* O157:H7, 1469
- Genetic profile
 - B. cereus*: enterotoxigenic and genetic profile, 2115
- Germicidal air cleaning unit
 - Control of airborne microorganisms in meat processing, 1922
- Germination
 - Thermal and hydrostatic inactivation, 196
- Giant freshwater prawn
 - Oxytetracycline residue depletion in giant freshwater prawn, 870
- Giardia*
 - Analysis of shellfish for *Cryptosporidium* and *Giardia*, 959
- Glial fibrillary acidic protein
 - Cross-contamination of CNS tissue due to beef carcass splitting, 83
- Gliding arc
 - Cold plasma sanitizing of apples, 1357
- Gluconic acid
 - Survival of *E. coli* O157:H7 in acid, 2404
- Gluten
 - Approaches to establish thresholds, 1043
- Gluten-free bread
 - Lactobacillus* for gluten and phytate hydrolysis in gluten-free bread, 1491
- Glycosylation
 - Antibacterial activity of dextran-conjugated lysozyme in cheese curd, 411
- Goat lactoferricin
 - Expression of goat lactoferricin from *E. coli*, 2523
- Gram-negative bacteria
 - Antibacterial activity of mushroom extracts, 1701
- Gram-positive bacteria
 - Antibacterial activity of mushroom extracts, 1701
 - Evaluation of *tdc* and its transcript in bacteria and foods, 93
- Grape
 - OTA determination in grapes, 2488
- Grape seed extract
 - Natural antimicrobial combinations against *L. monocytogenes*, 2105
- Green fluorescent protein
 - Temperature and nutrient effects on *C. jejuni*, 271
- Greenhouse
 - Microbiological profile of hydroponic tomatoes, 60
 - Salmonella on hydroponic tomatoes, 676
- Green olive fermentation
 - Survival of *B. cereus* during green olive fermentation, 1393
- Green tea extract
 - Natural antimicrobial combinations against *L. monocytogenes*, 2105
- Ground beef
 - Detection of risin in foods, 2053
 - Effects of reduced enrichment volumes, 1768
 - Facilitated thermal inactivation of *E. coli* O157:H7, 1604
 - Multiple-cycle high-pressure treatments, 811
- Ground muscle
 - Thermal inactivation of *Salmonella* in turkey breast, 2548
- Ground mustard
 - E. coli* O157:H7 inhibition in dry sausage by mustard powder, 486
- Ground pork
 - Y. enterocolitica* in ground pork packaged under modified atmospheres, 845
- Group-specific primers
 - Developing group-specific primers, 2257
- Growth
 - L. monocytogenes* growth at low temperature, 420
 - Model for growth of *Salmonella* on chicken frankfurters, 1135
- Growth kinetics
 - Growth of *Salmonella* during sprouting, 888

- Growth limits
Listeria inactivation affected by sequences of stresses, 1169
- Growth media
 Immunomodulatory activity of lactococci, 2124
- Guinea pigs
 Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- Gut enzymes
 In vitro digestion and bioavailability of LDPs, 979
- Half inhibitory concentration (IC₅₀)
 Detection of risin in foods, 2053
- Hamigera striata*
 Heat-resistant fungi from blueberries, 2030
- Hand washing
 Hand washing frequencies and procedures, 1641
- Harvest equipment
 On-farm contamination of persimmons, 52
- Hazard analysis critical control point (HACCP)
 Aerobic plate counts as a measure of HACCP effectiveness, 1248
 Food safety for the Turkish dairy industry, 440
 Risk factors for food premises unsatisfactory inspection results, 1632
- Hazelnut(s)
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
 Microbiology of raw nuts, 402
- Health Belief Model
 Cancer patients and food safety, 1666
- Health care establishments
 Sandwiches from health care establishments, 309
- Heat
 Altered cell composition in stress-adapted *L. innocua*, 182
 Decontamination of almonds with infrared heat, 897
- Heat inactivation
 Antibiotics in milk, 1193
- Heat resistance
 Heat resistance of *Salmonella* serotypes, 509
 Heat-resistant fungi from blueberries, 2030
 Heat tolerance of *C. sporogenes* spores, 953
- Heat shock
 Heat shock induces barotolerance in *L. monocytogenes*, 426
 Protein expression in stressed *V. parahaemolyticus*, 2589
- Heat treatment
 Growth inhibitory effects of kimchi against foodborne pathogens, 325
- Heavy metal
 Linking soil quality to human cadmium exposure, 2504
 Safety and quality assessment of lagoon grey mullets, 2572
- Hemolysis
 In vitro digestion and bioavailability of LDPs, 979
- Herbal
 Antibiotic-resistant bacteria in herbal products, 1486
- Heterogeneity
L. monocytogenes HHP inactivation, 2007
 Physiological response of *B. cereus* to food processing treatments, 2168
- Hexachlorobenzene
 Dietary intake of hexachlorobenzene, 2148
- Hidden allergens
 Allergen cross-contamination in the manufacture of fine bakery wares, 2263
- Hide(s)
E. coli O157:H7 and *Salmonella* in lairages, 1752
 On-farm meat safety, 1919
- Hide decontamination
 Reduction of hide microbial contamination, 2223
 Reduction of microbial contamination on cattle hides, 1343
- High hydrostatic pressure
L. monocytogenes HHP inactivation, 2007
 Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
- High-performance liquid chromatography (HPLC)
 Deoxynivalenol contamination in wheat products, 1931
- Rapid method used to analyze Ochratoxin A, 2133
- High pressure
 HPP inactivation of FCV, 1598
 Hydrolysates of β -lactoglobulin with low IgE binding, 1453
 Killing *E. coli* O157 with high pressure processing, 2182
 Multiple-cycle high-pressure treatments, 811
- High pressure processing (HPP)
Bacillus spore inactivation differences, using two pressurizing fluids, 1186
 Effect of HPP on freely suspended and bivalve-associated T7, 345
 Heat shock induces barotolerance in *L. monocytogenes*, 426
 High-pressure processing of human milk, 109
 High-pressure processing of Turkish white cheese, 102
 High-pressure treatment of *L. monocytogenes* in smoked salmon, 1612
 HPP in combination with antimicrobials, 820
- High shedders
E. coli O157 in feces and on beef carcasses, 1761
- High throughput
 Detection of *Salmonella* Enteritidis by polymixin-ELISA, 392
- Histo-blood group antigen expression
 Seasonal binding of recombinant norovirus to GI cells in oysters, 1696
- Histology
 Detection of *Sarcocystis* parasites in retail beef, 2144
- Honey dew melon
 SA11 rotavirus survival in fresh tropical fruit juices, 1035
- Horse meat
 Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
- Hot water dip
 Fludioxonil residues and control of blue and gray mold on pears, 967
- Hot water treatments
 Hot water treatment decontamination of mung bean seeds, 830
- House fly
 Enterococcal contamination of food by houseflies, 435
- HT-2
 LC-TOF-MS analysis of mycotoxins in figs, 1500
- Human(s)
 Antibiotic resistance of *Campylobacter* isolates from Poland, 602
 Virulence genes in *E. coli* O157 and non-O157 isolates, 1774
- Human milk
 High-pressure processing of human milk, 109
- Hydrogen peroxide
 UV light-hydrogen peroxide decontamination of lettuce, 903
- Hydrolyzable tannin
Q. infectoria as anti-*E. coli* agent, 1223
- Hydrophobic grid membrane filter
Salmonella identification and strain differentiation by HGMP-IR, 2249
- Hydroponics
 Microbiological profile of hydroponic tomatoes, 60
Salmonella on hydroponic tomatoes, 676
- Hydrosol
 Extending shelf life of fish, 218
- Hygiene
 Bacteriological safety in São Jorge cheese manufacture, 986
 Domestic refrigeration practices, 1898
- Hygiene indicator
 Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
- Hygiene indicator criteria
 Hygiene indicator criteria for meat in Belgium, 35
- Hypoosmolarity
 Low-salinity stress in *V. vulnificus*, 416
- Ice
 Transfer of *E. coli* O157:H7 to lettuce from ice, 252
- Icing
 Icing to control *V. vulnificus* and *V. parahaemolyticus*, 1475
- Identification
 Identification of *Salmonella* serotypes by PCR-RFLP, 2217
Salmonella identification and strain differentiation by HGMP-IR, 2249
- IgE-binding

- Hydrolysates of β -lactoglobulin with low IgE binding, 1453
- IgG-binding
- Hydrolysates of β -lactoglobulin with low IgE binding, 1453
- Immunoaffinity column
- Rapid method used to analyze Ochratoxin A, 2133
- Immunoassay drug test kits
- Efficacy of drug test kits for adulterated food, 1007
- Immunocapture
- Detection of *Campylobacter* spp., 2543
- Immunochromatography strip test
- Enhanced rapidity for *L. monocytogenes* detection, 781
- Immunohistochemistry
- Detection of DRG in advanced meat recovery product, 2307
- Immunomagnetic bead
- Enhanced rapidity for *L. monocytogenes* detection, 781
- Immunomagnetic separation
- Fecal versus RAMS detection of *E. coli* O157:H7, 691
- Immunomodulatory activity
- Immunomodulatory activity of lactococci, 2124
- Inactivation
- Decontamination of *E. coli* with ozone, 914
- Hot water treatment decontamination of mung bean seeds, 830
- HPP inactivation of FCV, 1598
- Survival of murine norovirus 1 on onions and spinach, 1590
- Inactivation curve
- Bacterial reduction in UV irradiated chill brines, 629
- Incubation temperature
- Effect of incubation temperature on APCs, 373
- Indicator organisms
- Microbiological quality of sushi, 860
- Indicators
- Surrogates for *Salmonella* in meat, 714
- Inducible enzyme
- M. morgani* ornithine decarboxylase, 657
- Infant formula
- Thermal resistance of *F. tularensis*, 2208
- Infection
- UV-light inactivation of viruses on fresh produce, 908
- Infective dose
- Infective doses and pathogen carriage, 2339
- Infiltration
- Salmonella* migration into almonds, 397
- Infrared
- Decontamination of almonds with infrared heat, 897
- Infrared microspectroscopy
- Salmonella* identification and strain differentiation by HGMF-IR, 2249
- Infrared spectroscopy
- Salmonella* identification and strain differentiation by HGMF-IR, 2249
- Inhibition
- Identification of AI-2 inhibitors from ground beef, 134
- Inoculum preparation
- L. monocytogenes* on salami during storage, 494
- Inspection scores
- Foodborne disease outbreaks in the catering industry, 550
- Internet
- Microbial quality of fish fillets, 1544
- Interventions
- Antimicrobial interventions for leafy greens, 625
- Antimicrobial interventions of bovine heads, 621
- Economics of *Campylobacter* control in poultry, 479
- Intervention selection
- Application of MCDA to a food safety decision-making problem, 2323
- Intestines
- Lymphoid tissue reduction in bovine small intestines, 1199
- In vitro gene expression
- In vitro cold shock gene expression in *Vibrio vulnificus*, 157
- In vivo toxicity
- Reduced toxicity of fumonisin B₁, 2036
- Ionic strength
- Survival of *E. coli* O157:H7 in acid, 2404
- Irradiation
- Food use and practices in long-term care facilities, 365
- Radiosensitivity of pathogens in ground beef, 516
- γ -Irradiation
- Combined treatments with irradiation on beef microbiology, 1237
- Isolation
- Isolation of *Arcobacter* from beef, 850
- Prevalence and diversity of *Arcobacter* in the Czech Republic, 719
- Italian cheese
- Decarboxylating strains in a typical Italian cheese, 205
- Japan
- A comparison of BSE risk between Japanese and U.S. beef, 802
- Japanese clams
- Viral contamination in clams, 1689
- Juice(s)
- SA11 rotavirus survival in fresh tropical fruit juices, 1035
- U.S. juice-associated outbreaks, 356
- Kimchi
- Growth inhibitory effects of kimchi against foodborne pathogens, 325
- Kinetic parameters
- Survival of *B. cereus* during green olive fermentation, 1393
- Kinetics
- Inactivation of *B. anthracis* in milk, 333
- Knife blades
- Listeria* transfer from knives to turkey breast, 176
- Knives
- Effect of different water temperatures on knife decontamination, 1338
- Kudoa*
- Kudoa* in Patagonian hakes, 2316
- Labeling
- Salmonellosis from stuffed chicken products, 2153
- Lactic acid
- Survival of *E. coli* O157:H7 in acid, 2404
- Lactic acid bacteria
- Antimicrobial susceptibility of *L. paracasei*, 339
- Bioprotective potential of LAB in malting and brewing, 1724
- Growth inhibitory effects of kimchi against foodborne pathogens, 325
- High-pressure processing of Turkish white cheese, 102
- L. monocytogenes* biofilms and bacteriocins, 634
- Storage stability of freeze-dried lactic acid bacteria, 431
- Survival of *B. cereus* during green olive fermentation, 1393
- Lactobacillus*
- Bacteriocin-producing starters in fermented sausages, 1817
- Lactobacillus* systemic immune functions and bacterial translocation, 796
- Low-molecular-weight antiyeast metabolites of a protective culture, 2481
- Lactobacillus casei*
- Antimicrobial susceptibility of *L. paracasei*, 339
- Lactobacillus paracasei*
- Antifungal activity of *L. paracasei*, 2213
- Lactobacillus pentosus*
- Effects of chloride salts on microbial growth, 1412
- Lactobacillus plantarum*
- Antimicrobial synergy between high pressure and Red No. 3, 1861
- Survival of *L. monocytogenes* in dry sausage, 1550
- Lactococcus lactis*
- Immunomodulatory activity of lactococci, 2124
- Nisin Z production in *L. lactis*, 2024
- Lactoferrin
- Antimicrobial activity of lactoferrin, 319
- Bactericidal effect of lactoferrin and its derivatives, 2468
- β -Lactoglobulin
- Hydrolysates of β -lactoglobulin with low IgE binding, 1453
- Immunoreactivity of irradiated milk and whey samples, 1270
- Lairage
- E. coli* O157:H7 and *Salmonella* in lairages, 1752
- Effect of transportation and lairage on risk of *S. enterica* in cattle, 2228

- Impact of transportation and lairage on *E. coli* O157:H7 in cattle, 1114
- Lateral flow
- Screening clenbuterol by strip and ELISA, 865
- Latex gloves
- Latex glove use by food handlers, 2334
- Latex sensitization
- Latex glove use by food handlers, 2334
- Lauric acid
- Poultry skin flora after successive washings, 1707
- Laying hens
- L. monocytogenes* in poultry flocks, 1996
- Leafy greens
- Antimicrobial interventions for leafy greens, 625
 - E-beam radiation for reducing pathogens in spinach, 2415
- Leafy vegetable
- Impact of water reuse in fresh-cut washing, 2514
- Legislation
- Regulation of safety of probiotics, 1734
- Lethality
- Controlling pathogens in beef jerky, 918
- Lettuce
- Survival of *E. coli* O157 on lettuce, 2384
 - UV light-hydrogen peroxide decontamination of lettuce, 903
 - Virus adsorption to lettuce, 522
- Leuconostoc* spp.
- LAB spoilage of vegetable sausages, 2312
- Lipid oxidation
- Acidified sodium chlorite treatment of trout fillets, 973
- Lipopeptide
- In vitro digestion and bioavailability of LDPs, 979
- Lipolysis
- Enterococci in spontaneous sausage fermentation, 1244
- Liquid chicken egg
- Detection of risin in foods, 2053
- Liquid chromatography
- Efficacy of drug test kits for adulterated food, 1007
 - LC-TOF-MS analysis of mycotoxins in figs, 1500
- Liquid chromatography-mass spectroscopy (LC-MS)
- LC-UV-MS method for DON, DON-3-glucoside, and pigment, 1205
- Liquid chromatography-tandem mass spectrometry (LC-MS-MS)
- Ochratoxin A in wine, 1038
- Liquid culture
- Antifungal activity of *L. paracasei*, 2213
- Liquid egg
- Thermal inactivation of bacteria in liquid egg, 1988
- Liquid whole egg
- Inactivation of *Salmonella* Senftenberg 775W by UHPH, 2283
- Listeria*
- Antimicrobial efficacy and sensory acceptability of plant essential oils, 1846
 - Bacteriocin-producing starters in fermented sausages, 1817
 - Listeria* and *Enterobacteriaceae* in domestic refrigerators, 608
 - Listeria* spp. in farmstead cheese processing environments, 2239
 - L. monocytogenes* inactivation by AS-48, 2460
 - L. monocytogenes* in cull sows and pork, 545
 - L. monocytogenes* sprout colonization and stress response, 1556
 - Risk assessment of *Listeria* in ready-to-eat chicken production line, 946
 - Survival of *L. innocua* in dairy products, 742
- Listeria innocua*
- Altered cell composition in stress-adapted *L. innocua*, 182
- Listeria monocytogenes*
- Antimicrobial essential oils against *L. monocytogenes* in pork, 502
 - Antimicrobial interventions for leafy greens, 625
 - Antimicrobial resistance in poultry bacteria, 2537
 - Antimicrobial synergy between high pressure and Red No. 3, 1861
 - Antimycotics and diacetate in RTE meats, 1386
 - Consumer assessment of safety/date labeling on RTE meat products, 70
 - Contribution of sigma B factor in alkali tolerance in *L. monocytogenes*, 1481
 - Controlling pathogens in beef jerky, 918
 - Control of *L. monocytogenes* on frankfurters, 728
 - CO₂ influence on *L. monocytogenes* growth, 1915
 - Distinguishable control strains for food microbiology testing, 2301
 - E-beam radiation of dry-cured ham, 2001
 - Enhanced rapidity for *L. monocytogenes* detection, 781
 - Enterococci in spontaneous sausage fermentation, 1244
 - Epidemiologic relevance of MVLST, 2559
 - Estimating *L. monocytogenes* growth kinetics parameters in frankfurters, 1806
 - Heat shock induces barotolerance in *L. monocytogenes*, 426
 - High-pressure processing of Turkish white cheese, 102
 - High-pressure treatment of *L. monocytogenes* in smoked salmon, 1612
 - Listeria* inactivation affected by sequences of stresses, 1169
 - Listeria* transfer from knives to turkey breast, 176
 - L. monocytogenes* adhesion and viability, 1379
 - L. monocytogenes* biofilms and bacteriocins, 634
 - L. monocytogenes* growth at low temperature, 420
 - L. monocytogenes* HHP inactivation, 2007
 - L. monocytogenes* in drain pipes, 66
 - L. monocytogenes* in poultry flocks, 1996
 - L. monocytogenes* in produce processing, 735
 - L. monocytogenes* on salami during storage, 494
 - Microbiology of sweet potato puree, 639
 - Modeling *L. monocytogenes* in pasteurized vanilla cream, 1828
 - Natural antimicrobial combinations against *L. monocytogenes*, 2105
 - Neural networks and survival of *L. monocytogenes* in soft cheese, 750
 - Pathogen incidence in raw milk, 1580
 - Radiation D₁₀-values of *L. monocytogenes* on catfish and tilapia, 2278
 - Salmonella* and *L. monocytogenes* in ready-to-eat food, 855
 - Sandwiches from health care establishments, 309
 - Stainless steel surface roughness and finish, 170
 - Superchilled storage of cold-smoked salmon, 2198
 - Survival of *L. monocytogenes*, 1835
 - Survival of *L. monocytogenes* in a gastrointestinal model, 2014
 - Survival of *L. monocytogenes* in dry sausage, 1550
 - Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
 - U.S. FDA regulatory management of human listeriosis, 1277
 - Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- Listeriosis
- Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- Liver
- Isolation of *C. jejuni* from bovine bile, 1228
- Liza* species
- Safety and quality assessment of lagoon grey mullets, 2572
- Logistic model
- Factors for produce contamination, 2389
- Logistic slaughter
- Contamination of carcasses with *Salmonella* during poultry slaughter, 146
- Long-chain polyphosphates
- Antibacterial effects of polyphosphates, 1401
- Long-term care facilities
- Food use and practices in long-term care facilities, 365
- Low-fat milk
- Detection of risin in foods, 2053
- Low-income consumers
- Consumer food safety knowledge and behaviors, 1651
- Low molecular weight
- Low-molecular-weight antiyeast metabolites of a protective culture, 2481
- Low pressure
- New sterilization process for aseptic packaging, 2119
- Low water activity
- Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
- Lymph node
- Salmonella* prevalence in bovine lymph nodes, 1685
- Lysozyme
- Active packaging, 119

- Antibacterial activity of dextran-conjugated lysozyme in cheese curd, 411
- Macrophage
Lactobacillus systemic immune functions and bacterial translocation, 796
- Macruronus magellanicus*
Kudoa in Patagonian hakes, 2316
- Magnesium chloride
 Effects of chloride salts on microbial growth, 1412
- Magnetic bead
 Enhanced rapidity for *L. monocytogenes* detection, 781
- Malic acid
 Essential oils in coatings for fresh-cut apples, 1150
- Malt
 Bioprotective potential of LAB in malting and brewing, 1724
 Occurrence of bound deoxynivalenol in *Fusarium*-infected barley, 1266
- Mango
 Decontamination of mango and melon by gas plasma, 302
 Plasma inactivation of bacteria migrating through fruits, 1619
- Mannoprotein
 Ochratoxin A adsorption by yeast, 1496
- Manure
 Determining *E. coli* O157:H7 levels in manures and soil, 2233
- Manure-amended soils
 Determining *E. coli* O157:H7 levels in manures and soil, 2233
- Marinade
Salmonella migration into marinated Turkey, 153
- Marination
 Heat inactivation of *E. coli* O157:H7 in nonintact beef, 1349
- Mass spectrometry
 LC-TOF-MS analysis of mycotoxins in figs, 1500
- Mathematical models
 Storage stability of freeze-dried lactic acid bacteria, 431
- Maturation time
 Toxoplasma in ready-to-eat smallgoods, 1442
- Maturity
Salmonella in tomatoes, 1572
- Meat
 Blade tenderization of beef containing *E. coli*, 2190
 Control of airborne microorganisms in meat processing, 1922
 Effect of different water temperatures on knife decontamination, 1338
 Thermal inactivation of viruses in chicken meat, 1214
- Meat-filled pastry
 Control of *C. perfringens* in Tajik sambusa, 77
- Meat microbiology
 Predictive microbiology, 6
- Meat processing
 Surrogates for *Salmonella* in meat, 714
- Meat products manufacturing
 Predictive microbiology, 6
- Meat safety
 On-farm meat safety, 1919
- Melamine
 Detection of melamine, 590
- Melon
 Decontamination of mango and melon by gas plasma, 302
 Plasma inactivation of bacteria migrating through fruits, 1619
- Mercury
 Mercury in bluefin tuna, 595
- Merluccius australis*
Kudoa in Patagonian hakes, 2316
- Merluccius hubbsi*
Kudoa in Patagonian hakes, 2316
- Meta-analysis
 Application of meta-analysis in food safety, 1330
- Metabiosis
 Metabolic effects of *Fusarium*, 1366
- Metabiotic interactions
 Proteolytic activity of molds, 2129
- Methicillin
mecA-positive *Staphylococcus* spp. in raw milk, 186
- Methylation analysis
 Cellulose production in *E. sakazakii* biofilms, 13
- MFO-3 method
 Alkaline phosphates in cow's, goat's, and sheep's milk, 1884
- Microalgae
 Antimicrobial activity of *D. salina* extracts, 2138
- Microarray
 Antimicrobial resistance and virulence in enterococci, 760
 Detection and genotyping noroviruses using DNA microarrays, 1434
 Toxin expression in O157 lineages, 1673
- Microbial contamination
 Microbiological quality of Australian retail raw meat, 1232
 Microbiological quality of chicken meat, 1123
- Microbial destruction
 Modification of the submerged coil, 775
- Microbial flora
 Combined treatments with irradiation on beef microbiology, 1237
- Microbial growth
 Impact of water reuse in fresh-cut washing, 2514
- Microbial inactivation
 Decontamination of mango and melon by gas plasma, 302
 High-pressure processing of human milk, 109
 High-pressure processing of Turkish white cheese, 102
 Plasma inactivation of bacteria migrating through fruits, 1619
- Microbial internalization
 Plasma inactivation of bacteria migrating through fruits, 1619
- Microbial population
 Grouping of food isolated bacteria by PCR-SSCP in food, 839
- Microbial spoilage
 Microbial spoilage of rutab, 1406
- Microbiological indices
 Safety and quality assessment of lagoon grey mullets, 2572
- Microbiological quality
 Microbial quality in mineral waters, 1253
 Microbial quality of stuffed mussels, 616
 Microbiological quality of sushi, 860
 Sandwiches from health care establishments, 309
- Microbiology
 Bacteriological safety in São Jorge cheese manufacture, 986
 Factors for produce contamination, 2389
 On-farm meat safety, 1919
- Micrococcaceae*
 Bacteriological safety in São Jorge cheese manufacture, 986
- Microsporidia
 Parasite inactivation by chlorine dioxide, 2410
- Microwaveable
 Salmonellosis from stuffed chicken products, 2153
- Migration
Salmonella migration into almonds, 397
Salmonella migration into marinated Turkey, 153
- Mild acidification
 Microbiology of sweet potato puree, 639
- Milk
 Alkaline phosphates in cow's, goat's, and sheep's milk, 1884
 Antibiotics in milk, 1193
 Heat shock induces barotolerance in *L. monocytogenes*, 426
- Milkfat globule membranes
 Antibacterial milkfat globule membrane fractions, 126
- Minced pork
 Antimicrobial essential oils against *L. monocytogenes* in pork, 502
- Mineral water
 Microbial quality in mineral waters, 1253
- Modeling
 CO₂ influence on *L. monocytogenes* growth, 1915
 Modeling the effect of sublethal history, 279
 Modeling the growth of *Salmonella* in poultry, 2429
 Neural networks and survival of *L. monocytogenes* in soft cheese, 750
- Modified atmosphere packaging (MAP)

- Combined treatments with irradiation on beef microbiology, 1237
 CO₂ influence on *L. monocytogenes* growth, 1915
 Proteolytic *C. botulinum* in rice at low pH, 468
 Radiosensitivity of pathogens in ground beef, 516
 Spoilage and safety characteristics of ground beef, 293
Y. enterocolitica in ground pork packaged under modified atmospheres, 845
- Molds**
 Efficient fungal DNA extraction method for RAPD-PCR, 2497
 Survival of *L. innocua* in dairy products, 742
- Molecular detection**
 Effects of reduced enrichment volumes, 1768
- Molecular typing**
 Detection and genotyping noroviruses using DNA microarrays, 1434
- Monilinia laxa***
 Potassium sorbate for controlling postharvest brown rot, 1626
- Monitoring**
Salmonella prevalence in the broiler supply chain, 1974
- Monoclonal antibody**
 Competitive ELISA to detect bovine blood, 1000
- Monolithic column**
 Rapid method used to analyze Ochratoxin A, 2133
- Most probable number**
 Automated MPN determination of *Enterobacteriaceae* in food, 376
- MS2 coliphage**
 UV light-hydrogen peroxide decontamination of lettuce, 903
- Multi-criteria decision analysis**
 Application of MCDA to a food safety decision-making problem, 2323
- Multiple antibiotic resistance**
 Transcription of *stx1*, *marA*, and *eaeA* in *E. coli* O157:H7, 1469
- Multiple cycles**
 Multiple-cycle high-pressure treatments, 811
- Multiple genes**
 SSCP analysis for rapid discrimination of *Salmonella* isolates, 1960
- Multiplex PCR**
 Multiplex PCR detection of enterotoxin genes in *Aeromonas* spp., 2094
 Prevalence and diversity of *Arcobacter* in the Czech Republic, 719
- Multiplication**
Salmonella Enteritidis growth on egg yolk membranes, 1905
- Multivariate analytical approach**
 Survival of *B. cereus* during green olive fermentation, 1393
- Multi-virulence-locus sequence typing (MVLST)**
 Epidemiologic relevance of MVLST, 2559
- Mung bean seeds**
 Hot water treatment decontamination of mung bean seeds, 830
- Murine norovirus**
 Survival of murine norovirus 1 on onions and spinach, 1590
- Musca domestica***
 Enterococcal contamination of food by houseflies, 435
- Muscle**
 Danofloxacin in beef muscle, 1720
- Muscle tissue**
 Distribution of drug residues in breast muscle tissues, 223
- Mushrooms**
 Antibacterial activity of mushroom extracts, 1701
- Mussels**
 Analysis of shellfish for *Cryptosporidium* and *Giardia*, 959
- Mustard**
 Combined treatments with irradiation on beef microbiology, 1237
- Mycotoxin(s)**
Alternaria toxins in wheat, 1262
 LC-UV-MS method for DON, DON-3-glucoside, and pigment, 1205
 Mycotoxins produced in apples, 1714
 Ochratoxin A in wine, 1038
 OTA removal procedure for cocoa shells, 1422
- Myxosporean**
Kudoa in Patagonian hakes, 2316
- NaCl stress**
Listeria inactivation affected by sequences of stresses, 1169
- N*-Acyl homoserine lactone
 Role of quorum sensing in food spoilage, 1510
- Naïve penmates**
 Fecal pats inoculated with *E. coli* O157:H7 to mimic a super shedder, 648
- Nalidixic acid resistance**
 Distinguishable control strains for food microbiology testing, 2301
- Natural antimicrobial**
E. coli O157:H7 inhibition in dry sausage by mustard powder, 486
- Neural networks**
 Neural networks and survival of *L. monocytogenes* in soft cheese, 750
- Newcastle disease**
 Thermal inactivation of viruses in chicken meat, 1214
- New Zealand**
Campylobacter on retail broiler carcasses in New Zealand, 2526
- N*-Halamine
 Rechargeable antimicrobial polyethylene, 2042
- Nigella sativa***
 Influence of extraction methods on *N. sativa* essential oils, 2475
- Nisin**
 Natural antimicrobial combinations against *L. monocytogenes*, 2105
 Nisin Z production in *L. lactis*, 2024
- Nisin variant**
 Nisin Z production in *L. lactis*, 2024
- Nonintact beef**
 Heat inactivation of *E. coli* O157:H7 in nonintact beef, 1349
- Nonthermal**
 Dense phase CO₂ processing of spinach, 1015
 UV-light inactivation of viruses on fresh produce, 908
- Nonthermal plasma**
 Cold plasma sanitizing of apples, 1357
- Norovirus**
 Detection and genotyping noroviruses using DNA microarrays, 1434
 Detection of noroviruses in oysters, 1427
 Seasonal binding of recombinant norovirus to GI cells in oysters, 1696
- Norovirus binding**
 Seasonal binding of recombinant norovirus to GI cells in oysters, 1696
- Nuggets**
 Bacteriology of raw chicken nuggets, 613
- Nut**
 Decontamination of almonds with infrared heat, 897
- Nutraceutical products**
 Antibiotic-resistant bacteria in herbal products, 1486
- Nutrient effects**
 Temperature and nutrient effects on *C. jejuni*, 271
- Ochratoxin A**
 Ochratoxin A adsorption by yeast, 1496
 Ochratoxin A in wine, 1038
 OTA determination in grapes, 2488
 OTA removal procedure for cocoa shells, 1422
 Rapid method used to analyze Ochratoxin A, 2133
- Oligosaccharide**
 Inhibition of pathogen adhesion, 2272
- Olympic Games 2004**
 Risk factors for food premises unsatisfactory inspection results, 1632
- On-farm risk factors**
 Prevalence of *Y. enterocolitica*, 1162
- Opportunistic human pathogens**
 16S rDNA profiling of alfalfa sprouts-associated bacteria, 200
- Oral exposure**
 Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- Oregano**
 Combined treatments with irradiation on beef microbiology, 1237
- Organic**
 Antimicrobial resistance in poultry bacteria, 2537
- Organic acid preservative**
 Antimycotics and diacetate in RTE meats, 1386
- Organic acids**
 Survival of *B. cereus* during green olive fermentation, 1393

- Organic matter
Impact of water reuse in fresh-cut washing, 2514
- Organoleptic properties
Decontamination of *E. coli* with ozone, 914
- Outbreak(s)
Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
Salmonellosis from stuffed chicken products, 2153
- Oxidative stress
L. monocytogenes sprout colonization and stress response, 1556
- Oxygen
Spoilage and safety characteristics of ground beef, 293
Survival of *E. coli* O157:H7 in acid, 2404
- Oxytetracycline
Oxytetracycline residue depletion in giant freshwater prawn, 870
- Oyster
Detection of noroviruses in oysters, 1427
Seasonal binding of recombinant norovirus to GI cells in oysters, 1696
- Ozone
Decontamination of *E. coli* with ozone, 914
- Packaging
Campylobacter on retail broiler carcasses in New Zealand, 2526
- Packing shed equipment
On-farm contamination of Satsuma mandarin fruit, 530
- Packing sheds
Microflora on cantaloupes, 46
- Papaya
SA11 rotavirus survival in fresh tropical fruit juices, 1035
- Parasite antigens
Anisakis antigens in fish muscle, 1273
- Pasteurization
Dense phase CO₂ processing of spinach, 1015
Food use and practices in long-term care facilities, 365
- Pasteurized
Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
- Pathogen
Antibiotic resistance in *V. parahaemolyticus*, 2552
Controlling pathogens in beef jerky, 918
Decontamination of almonds with infrared heat, 897
Microbial quality of fish fillets, 1544
- Pathogen carriage
Infective doses and pathogen carriage, 2339
- Pathogenic bacteria
Antimicrobial activity of lactoferrin, 319
Impact of water reuse in fresh-cut washing, 2514
On-farm contamination of persimmons, 52
On-farm contamination of Satsuma mandarin fruit, 530
- Pathogen interventions
E-beam radiation for reducing pathogens in spinach, 2415
- Pathogen reduction
E-beam radiation for reducing pathogens in spinach, 2415
- Pathogens
Microbiological quality of Australian retail raw meat, 1232
Microbiological quality of chicken meat, 1123
- Patulin
LC-TOF-MS analysis of mycotoxins in figs, 1500
Mycotoxins produced in apples, 1714
Sanitizers for inactivation of *P. expansum* on apples, 643
- PCR
Detection of *Sarcocystis* parasites in retail beef, 2144
Developing group-specific primers, 2257
Evaluation of *tdc* and its transcript in bacteria and foods, 93
Listeria spp. in farmstead cheese processing environments, 2239
Occurrence of STEC in meat products in Morocco, 2082
PCR detection of *Salmonella* by *stdA* primers, 1108
Salmonella detection methods on turkey parts, 386
S. aureus and Domiat cheese, 2567
Toxin expression in O157 lineages, 1673
- PCR detection
DNA extraction for PCR detection of food pathogens, 2110
- PCR-restriction fragment length polymorphism
Identification of *Salmonella* serotypes by PCR-RFLP, 2217
- PCR-single-strand conformation polymorphism (PCR-SSCP)
Grouping of food isolated bacteria by PCR-SSCP in food, 839
- Peanuts
Microbiology of raw nuts, 402
- Pectin
Inhibition of pathogen adhesion, 2272
- Penetration
Salmonella Enteritidis growth on egg yolk membranes, 1905
- Penicillium expansum*
Mycotoxins produced in apples, 1714
Sanitizers for inactivation of *P. expansum* on apples, 643
- Pentitol
Xylitol to reduce microbial adhesion, 405
- Pepsin digestion
Bactericidal effect of lactoferrin and its derivatives, 2468
- Peracetic acid
Peracetic acid mixture as an antimicrobial treatment in poultry, 1119
Sanitizers for inactivation of *P. expansum* on apples, 643
Survival of murine norovirus 1 on onions and spinach, 1590
- Pericarp
Decontamination of mango and melon by gas plasma, 302
- Persimmons
On-farm contamination of persimmons, 52
- Persistence
Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- Pesticide solution
On-farm contamination of Satsuma mandarin fruit, 530
- pH
HPP inactivation of FCV, 1598
Proteolytic *C. botulinum* in rice at low pH, 468
- Phage display
Enzyme immunoassay for detection for amygdalin, 2048
- Pharmaceuticals
Regulation of safety of probiotics, 1734
- Phenotypical tracing
Tracing the origin of coliform bacteria, 1910
- Phylogenetic tree
Viral contamination in clams, 1689
- Physicochemical properties
Physicochemical properties of *Salmonella*, 2421
- Physiology
Physiological response of *B. cereus* to food processing treatments, 2168
- Phytopathogens
Survival of *E. coli* O157 on lettuce, 2384
- Pig
Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
Time course of infection with *Salmonella* Typhimurium in fattening pigs, 699
- Pig carcass
Application of meta-analysis in food safety, 1330
- Piglets
Effects of a probiotic culture on performance of weaned piglets, 1797
- Pigment
LC-UV-MS method for DON, DON-3-glucoside, and pigment, 1205
- Pineapple
SA11 rotavirus survival in fresh tropical fruit juices, 1035
- Planktonic
Fourteen-day-old chicken cecal microflora, 1981
- Plant-derived compounds
Inactivation of antibiotic-resistant *Campylobacter*, 1145
- Plants
L. monocytogenes sprout colonization and stress response, 1556
- Plasma sterilization
New sterilization process for aseptic packaging, 2119
- Plasma treatment

- Active packaging, 119
- Polycyclic aromatic hydrocarbon (PAH)
Detection of PAHs in smoked meat, 993
- Polyethylene (PE)
Rechargeable antimicrobial polyethylene, 2042
- Polyethylene terephthalate (PET)
New sterilization process for aseptic packaging, 2119
- Polygalacturonase
Potassium sorbate for controlling postharvest brown rot, 1626
- Polymixin-ELISA
Detection of *Salmonella* Enteritidis by polymixin-ELISA, 392
- Pork
Application of meta-analysis in food safety, 1330
L. monocytogenes in cull sows and pork, 545
- Pork meat
Aerobic plate counts as a measure of HACCP effectiveness, 1248
- Postharvest treatment
Icing to control *V. vulnificus* and *V. parahaemolyticus*, 1475
- Potassium chloride
Effects of chloride salts on microbial growth, 1412
- Potassium lactate
Control of *C. perfringens* in Tajik sambusa, 77
Estimating *L. monocytogenes* growth kinetics parameters in frankfurters, 1806
- Potassium sorbate
Potassium sorbate for controlling postharvest brown rot, 1626
- Potential sources of contamination
On-farm contamination of persimmons, 52
On-farm contamination of Satsuma mandarin fruit, 530
- Poultry
Antimicrobial resistance in poultry bacteria, 2537
Bacteriology of raw chicken nuggets, 613
Contamination of carcasses with *Salmonella* during poultry slaughter, 146
Detection methods for *Campylobacter* spp., 835
Mapping of *Salmonella* contamination on the poultry carcass, 1785
Physicochemical properties of *Salmonella*, 2421
Salmonella prevalence in the broiler supply chain, 1974
- Poultry chilling
Peracetic acid mixture as an antimicrobial treatment in poultry, 1119
- Poultry processing
Arcobacter spp. in poultry processing, 2533
- Poultry skin
Poultry skin flora after successive washings, 1707
- Powdered infant formula
Resistance of *E. sakazakii* to desiccation, 940
- Powdered infant milk formula
Inactivation of desiccated *E. sakazakii*, 934
- Powdered milk
Role of fat content in trapping volatile compounds, 1889
- Practicality
Application of MCDA to a food safety decision-making problem, 2323
- Prawn
Incidence of *Salmonella* in imported seafood, 1460
- Predictive microbiology
Model for growth of *Salmonella* on chicken frankfurters, 1135
Neural networks and survival of *L. monocytogenes* in soft cheese, 750
Survival of *L. monocytogenes*, 1835
- Predictive modeling
Predictive microbiology, 6
- Pregnancy
Virulence of persistent *L. monocytogenes* in pregnant guinea pigs, 1028
- Preharvest food safety
Dust exposure and pathogen increases on cattle hides, 2078
- Preservative
L. monocytogenes inactivation by AS-48, 2460
- Pressure-thermal processing
Bacillus spore inactivation differences, using two pressurizing fluids, 1186
- Prevalence
Application of meta-analysis in food safety, 1330
Arcobacter spp. in poultry processing, 2533
Campylobacter in paired turkey and swine farms, 1791
Campylobacter prevalence in poultry production, 2295
L. monocytogenes in poultry flocks, 1996
Prevalence and diversity of *Arcobacter* in the Czech Republic, 719
Salmonella prevalence in the broiler supply chain, 1974
- Probiotics
Antimicrobial susceptibility of *L. paracasei*, 339
Effects of a probiotic culture on performance of weaned piglets, 1797
Immunomodulatory activity of lactococci, 2124
Regulation of safety of probiotics, 1734
- Processed food
Tracing the origin of coliform bacteria, 1910
- Processed spinach
Coliforms and foodborne pathogens on spinach in packing plants, 2398
- Processing surfaces
Electrolyzed water in the food industry, 1934
- Process validation
Surrogates for *Salmonella* in meat, 714
- Produce
Dense phase CO₂ processing of spinach, 1015
E. coli O157:H7 enrichment in produce, 139
Factors for produce contamination, 2389
L. monocytogenes in produce processing, 735
UV-light inactivation of viruses on fresh produce, 908
- Production equipment
Allergen cross-contamination in the manufacture of fine bakery wares, 2263
- Propionibacterium*
Low-molecular-weight antiyeast metabolites of a protective culture, 2481
- Protein expression
Protein expression in stressed *V. parahaemolyticus*, 2589
- Proteolysis
Enterococci in spontaneous sausage fermentation, 1244
Hydrolysates of β -lactoglobulin with low IgE binding, 1453
Lactobacillus for gluten and phytate hydrolysis in gluten-free bread, 1491
- Proteolytic molds
Proteolytic activity of molds, 2129
- Pseudomonas fluorescens*
Bactericidal effect of lactoferrin and its derivatives, 2468
- Pseudomonas syringae*
In vitro digestion and bioavailability of LDPs, 979
- Psychrotrophic counts
Acidified sodium chlorite treatment of trout fillets, 973
- Pulsed-field gel electrophoresis (PFGE)
Enterococci from human patients and foods, 2100
Enterococci in spontaneous sausage fermentation, 1244
Occurrence of STEC in meat products in Morocco, 2082
PFGE scheme for subtyping *Salmonella*, 2067
- Purge-and-trap optimization
Role of fat content in trapping volatile compounds, 1889
- Putrescine
M. morgani ornithine decarboxylase, 657
- Pyrus communis* L.
Fludioxonil residues and control of blue and gray mold on pears, 967
- Qualitative microbiological method
BAX system PCR method for *Salmonella* versus Brazil's official method, 2442
- Quality assurance
Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
- Quantification
L. monocytogenes HHP inactivation, 2007
- Quantitative
C. jejuni cross-contamination, 1018
- Quercus infectoria*
Q. infectoria as anti-*E. coli* agent, 1223

- Quorum quenching
Role of quorum sensing in food spoilage, 1510
- Quorum sensing
Role of quorum sensing in food spoilage, 1510
- Rabbit IgG
Immunoreactivity of irradiated milk and whey samples, 1270
- Radial basis function
Neural networks and survival of *L. monocytogenes* in soft cheese, 750
- Radial diffusion assay
Potassium sorbate for controlling postharvest brown rot, 1626
- Radiation D_{10} -values
Radiation D_{10} -values of *L. monocytogenes* on catfish and tilapia, 2278
- Radio frequency electric field (RFEF)
Membrane damage and inactivation of bacteria in apple juice by RFEF, 684
- Radiosensitivity
Radiosensitivity of pathogens in ground beef, 516
- Radiosensitization
Combined treatments with irradiation on beef microbiology, 1237
- Randomly amplified polymorphic DNA-PCR
Efficient fungal DNA extraction method for RAPD-PCR, 2497
- Rapid methods
Automated MPN determination of *Enterobacteriaceae* in food, 376
- Rapid molecular method
Detection methods for *Campylobacter* spp., 835
- Rats
Acute toxicity of rosemary extracts, 790
- Raw
Bacteriology of raw chicken nuggets, 613
- Raw fish
Microbiological quality of sushi, 860
- Raw meat
Campylobacter spp. in raw and ready-to-eat foods, 2087
Occurrence of STEC in meat products in Morocco, 2082
- Raw milk
Campylobacter spp. in raw and ready-to-eat foods, 2087
Pathogen incidence in raw milk, 1580
Predicting *Salmonella* prevalence in a dairy herd, 1967
Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
- Raw poultry
Modeling the growth of *Salmonella* in poultry, 2429
- Ready-to-eat food
Campylobacter spp. in raw and ready-to-eat foods, 2087
Enterococcal contamination of food by houseflies, 435
Salmonella and *L. monocytogenes* in ready-to-eat food, 855
- Ready-to-eat meat
Killing *E. coli* O157 with high pressure processing, 2182
- Ready-to-eat meat products
Consumer assessment of safety/date labeling on RTE meat products, 70
- Ready-to-eat poultry products
Consumer assessment of safety/date labeling on RTE meat products, 70
- Real-time PCR (RT-PCR)
Detection of *Campylobacter* spp., 2543
Detection of central nervous system tissues in meat products, 2059
Detection of noroviruses in oysters, 1427
Detection of *Salmonella* by PCR in foods, 2436
Prevalence of *Y. enterocolitica*, 1162
Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
- Recalls
U.S. FDA regulatory management of human listeriosis, 1277
- Recontamination
Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
- Rectoanal mucosal swab (RAMS)
Fecal versus RAMS detection of *E. coli* O157:H7, 691
- Refrigerated storage
Salmonella Enteritidis in eggs, 2073
- Refrigerators
Listeria and *Enterobacteriaceae* in domestic refrigerators, 608
- Regulations
U.S. juice-associated outbreaks, 356
- Regulators
L. monocytogenes growth at low temperature, 420
- Regulatory Fish Encyclopedia
DNA barcoding and the *Regulatory Fish Encyclopedia*, 210
- Regulatory science
DNA barcoding and the *Regulatory Fish Encyclopedia*, 210
- Residuals
Evaluation of a copper sulfate-based commercial disinfectant, 226
- Residue(s)
Fludioxonil residues and control of blue and gray mold on pears, 967
Oxytetracycline residue depletion in giant freshwater prawn, 870
- Resistance
Antibiotic resistance of *Campylobacter* isolates from Poland, 602
Antibiotic-resistant bacteria in herbal products, 1486
Effects of interventions on MDR *Salmonella*, 2177
Salmonella prevalence in bovine lymph nodes, 1685
- Restaurant inspection
Risk factors for food premises unsatisfactory inspection results, 1632
- Retail
Campylobacter on retail broiler carcasses in New Zealand, 2526
Microbial quality of fish fillets, 1544
- Retail beef
Isolation of *Arcobacter* from beef, 850
- Retail food
Antimicrobial resistance and virulence in enterococci, 760
- Retail food services
Hand washing frequencies and procedures, 1641
- Retail meat
Microbiological quality of Australian retail raw meat, 1232
- Reverse transcriptase PCR
Detection and genotyping noroviruses using DNA microarrays, 1434
Viral contamination in clams, 1689
- rfbE*_{O157}
Prevalence of *E. coli* O157:H7 in vegetables, 816
- Ribotype
Distinguishable control strains for food microbiology testing, 2301
Listeria spp. in farmstead cheese processing environments, 2239
- Rice
V. cholerae survival on different types of cooked rice, 2453
- Ricin
Detection of ricin in foods, 2053
Ricin and abrin in beverages, 1875
- Ring trial
Microscopic method for animal constituents detection in feedstuffs, 578
- Ripeness
Salmonella in tomatoes, 1572
- Risk assessment
CO₂ influence on *L. monocytogenes* growth, 1915
Risk assessment for STEC in unpasteurized milk, 257
Risk assessment of *Listeria* in ready-to-eat chicken production line, 946
Survival of *L. monocytogenes*, 1835
- Risk factors
Risk factors for food premises unsatisfactory inspection results, 1632
Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
- Romaine lettuce
Transfer of *E. coli* O157:H7 to lettuce from ice, 252
- Rosemary extracts
Acute toxicity of rosemary extracts, 790
- Rotavirus
SA11 rotavirus survival in fresh tropical fruit juices, 1035
- Rules
U.S. juice-associated outbreaks, 356
- Rutab
Microbial spoilage of rutab, 1406
- Saccharomyces cerevisiae*

- Effects of chloride salts on microbial growth, 1412
- Safety
- Microbial quality of fish fillets, 1544
 - Microbial quality of stuffed mussels, 616
- Safety labeling
- Consumer assessment of safety/date labeling on RTE meat products, 70
- Salami
- Killing *E. coli* O157 with high pressure processing, 2182
 - Survival of *L. monocytogenes* in a gastrointestinal model, 2014
- Salinity
- Low-salinity stress in *V. vulnificus*, 416
- Salmonella*
- Antimicrobial interventions for leafy greens, 625
 - Application of meta-analysis in food safety, 1330
 - Bacteriology of raw chicken nuggets, 613
 - Campylobacter* on retail broiler carcasses in New Zealand, 2526
 - Cold plasma sanitizing of apples, 1357
 - Coliforms and foodborne pathogens on spinach in packing plants, 2398
 - Contamination of carcasses with *Salmonella* during poultry slaughter, 146
 - Controlling pathogens in beef jerky, 918
 - Decontamination of almonds with infrared heat, 897
 - Detection of *Salmonella* by PCR in foods, 2436
 - Detection of *Salmonella* using a simple phage assay, 380
 - Distinguishable control strains for food microbiology testing, 2301
 - Dust exposure and pathogen increases on cattle hides, 2078
 - E-beam radiation for reducing pathogens in spinach, 2415
 - E. coli* O157:H7 and *Salmonella* in lairages, 1752
 - Effects of interventions on MDR *Salmonella*, 2177
 - Growth of *Salmonella* during sprouting, 888
 - Heat resistance of *Salmonella* serotypes, 509
 - Hot water treatment decontamination of mung bean seeds, 830
 - HPP in combination with antimicrobials, 820
 - Identification of *Salmonella* serotypes by PCR-RFLP, 2217
 - Incidence of *Salmonella* in imported seafood, 1460
 - Microbiological profile of hydroponic tomatoes, 60
 - Microbiology of raw nuts, 402
 - Modeling the effect of sublethal history, 279
 - Modeling the growth of *Salmonella* in poultry, 2429
 - Multidrug-resistant *Salmonella*, 27
 - Operational practices associated with foodborne disease outbreaks, 1659
 - Pathogen incidence in raw milk, 1580
 - PCR detection of *Salmonella* by *stdA* primers, 1108
 - Peracetic acid mixture as an antimicrobial treatment in poultry, 1119
 - Physicochemical properties of *Salmonella*, 2421
 - Predicting *Salmonella* prevalence in a dairy herd, 1967
 - Probiotics effect on *E. coli* O157:H7 and *Salmonella* shedding in cattle, 539
 - Proteolytic activity of molds, 2129
 - Reduction of hide microbial contamination, 2223
 - Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
 - Salmonella* and *L. monocytogenes* in ready-to-eat food, 855
 - Salmonella* contamination of food service eggs, 19
 - Salmonella* detection methods on turkey parts, 386
 - Salmonella* identification and strain differentiation by HGMP-IR, 2249
 - Salmonella* in tomatoes, 1572
 - Salmonella* migration into almonds, 397
 - Salmonella* migration into marinated Turkey, 153
 - Salmonella* on hydroponic tomatoes, 676
 - Salmonella* prevalence in bovine lymph nodes, 1685
 - Salmonella* prevalence in the broiler supply chain, 1974
 - Salmonellosis from stuffed chicken products, 2153
 - Shedding of foodborne pathogens by Florida cattle, 807
 - Spoilage and safety characteristics of ground beef, 293
 - SSCP analysis for rapid discrimination of *Salmonella* isolates, 1960
 - Surrogates for *Salmonella* in meat, 714
 - Thermal inactivation of *Salmonella* in turkey breast, 2548
 - Time course of infection with *Salmonella* Typhimurium in fattening pigs, 699
 - Salmonella enterica*
 - Effect of transportation and lairage on risk of *S. enterica* in cattle, 2228
 - Salmonella enterica* serovar Enteritidis
 - Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
 - Salmonella enterica* serovar Senftenberg 775W
 - Inactivation of *Salmonella* Senftenberg 775W by UHPH, 2283
 - Salmonella* Enteritidis
 - Detection of *Salmonella* Enteritidis by polymixin-ELISA, 392
 - Salmonella* Enteritidis growth on egg yolk membranes, 1905
 - Salmonella* Enteritidis in eggs, 2073
 - Thermal inactivation of bacteria in liquid egg, 1988
 - Salmonella* Hadar
 - PFGE scheme for subtyping *Salmonella*, 2067
 - Salmonella* Heidelberg
 - PFGE scheme for subtyping *Salmonella*, 2067
 - Salmonella* Kentucky
 - PFGE scheme for subtyping *Salmonella*, 2067
 - Salmonella* SaintPaul
 - PFGE scheme for subtyping *Salmonella*, 2067
 - Salmonella* Typhi
 - Radiosensitivity of pathogens in ground beef, 516
 - Salmonella* Typhimurium DT104
 - Mapping of *Salmonella* contamination on the poultry carcass, 1785
 - Model for growth of *Salmonella* on chicken frankfurters, 1135
 - Salmonellosis
 - Salmonella* contamination of food service eggs, 19
 - Salt
 - Toxoplasma in ready-to-eat smallgoods, 1442
 - Sambusa
 - Control of *C. perfringens* in Tajik sambusa, 77
 - Sample pretreatment
 - Detection of *Campylobacter* spp., 2543
 - Sandwiches
 - Sandwiches from health care establishments, 309
 - Sanitization
 - Decontamination of almonds with infrared heat, 897
 - Sanitizer(s)
 - Efficacy of household products against *Bacillus* spores, 473
 - Inactivation of *B. anthracis* in milk, 333
 - L. monocytogenes* in drain pipes, 66
 - Sanitizing
 - Risk assessment of *Listeria* in ready-to-eat chicken production line, 946
 - Sanitizers for inactivation of *P. expansum* on apples, 643
 - Sarcocystis cruzi*
 - Detection of *Sarcocystis* parasites in retail beef, 2144
 - Sarcocystis hominis*
 - Detection of *Sarcocystis* parasites in retail beef, 2144
 - Satsuma mandarin
 - On-farm contamination of Satsuma mandarin fruit, 530
 - Sausage fermentation
 - Bacteriocin-producing starters in fermented sausages, 1817
 - Enterococci in spontaneous sausage fermentation, 1244
 - Scalder
 - Evaluation of a copper sulfate-based commercial disinfectant, 226
 - Scanning electron microscopy
 - Allergen release by frozen and chilled *Anisakis*, 2578
 - Seafood
 - Cooking parameters for safe seafood for consumers, 1287
 - C. perfringens* and *B. cereus* in seafood, 1178
 - Incidence of *Salmonella* in imported seafood, 1460
 - Microbiological quality of sushi, 860
 - Seasonal change
 - Seasonal binding of recombinant norovirus to GI cells in oysters, 1696
 - Secretions
 - Sources of contamination and pathogen excretion from infected persons, 2582
 - Semifried tuna
 - Shelf life of coated and semifried tuna, 770
 - Sensory

- Antimicrobial efficacy and sensory acceptability of plant essential oils, 1846
- Sensory analysis
Superchilled storage of cold-smoked salmon, 2198
- Sequence of stresses
Listeria inactivation affected by sequences of stresses, 1169
- Serology
Time course of infection with *Salmonella* Typhimurium in fattening pigs, 699
- Serotype(s)
Heat resistance of *Salmonella* serotypes, 509
Identification of *Salmonella* serotypes by PCR-RFLP, 2217
L. monocytogenes in poultry flocks, 1996
- Serovars
Salmonella identification and strain differentiation by HGMP-IR, 2249
- Sheep
Effect of incubation temperature on APCs, 373
- Shelf life
Acidified sodium chlorite treatment of trout fillets, 973
Combined treatments with irradiation on beef microbiology, 1237
Essential oils in coatings for fresh-cut apples, 1150
Estimating *L. monocytogenes* growth kinetics parameters in frankfurters, 1806
Extending shelf life of fish, 218
Shelf life of coated and semifried tuna, 770
- Shell eggs
Yeasts from refrigerated commercial shell eggs, 1258
- Shellfish
Analysis of shellfish for *Cryptosporidium* and *Giardia*, 959
Detection of noroviruses in oysters, 1427
Effect of HPP on freely suspended and bivalve-associated T7, 345
- Shiga toxinogenic *E. coli*
Non-O157 STEC from humans, cattle, and food in the United States, 1023
- Shiga toxin(s)
Detection of Shiga toxin-producing *E. coli*, 927
STEC and *E. coli* O157 from retail meats in China, 706
- Shiga toxin-containing *E. coli* (STEC)
E. coli O157:H7 enrichment in produce, 139
- Shrimp
Incidence of *Salmonella* in imported seafood, 1460
- Sigma B factor
Contribution of sigma B factor in alkali tolerance in *L. monocytogenes*, 1481
- Simulation
A comparison of BSE risk between Japanese and U.S. beef, 802
- Single-strand conformation polymorphism
SSCP analysis for rapid discrimination of *Salmonella* isolates, 1960
- 16S rDNA
Grouping of food isolated bacteria by PCR-SSCP in food, 839
16S rDNA profiling of alfalfa sprouts-associated bacteria, 200
- 16S rRNA gene
Prevalence and diversity of *Arcobacter* in the Czech Republic, 719
- Slaughter cattle
Reduction of microbial contamination on cattle hides, 1343
- Slaughterhouse
Risk factors for *Salmonella* and hygiene indicators in pig slaughterhouses, 1320
- Smallholder dairy
Risk assessment for STEC in unpasteurized milk, 257
- Small subunit rDNA
Kudoa in Patagonian hakes, 2316
- Smoked pork
Detection of PAHs in smoked meat, 993
- Smoked salmon
High-pressure treatment of *L. monocytogenes* in smoked salmon, 1612
Superchilled storage of cold-smoked salmon, 2198
- Sodium benzoate
Antimicrobials and diacetate in RTE meats, 1386
Transcription of *stx1*, *marA*, and *eaeA* in *E. coli* O157:H7, 1469
- Sodium chloride
Antimicrobials and diacetate in RTE meats, 1386
Effects of chloride salts on microbial growth, 1412
- Sodium diacetate
Antimicrobials and diacetate in RTE meats, 1386
Estimating *L. monocytogenes* growth kinetics parameters in frankfurters, 1806
- Sodium hypochlorite
Sanitizers for inactivation of *P. expansum* on apples, 643
- Soft cheese
Neural networks and survival of *L. monocytogenes* in soft cheese, 750
Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
- Soil
Linking soil quality to human cadmium exposure, 2504
- Sorbic acid
Microbiology of sweet potato puree, 639
- Sourdough *Lactobacillus*
Lactobacillus for gluten and phytate hydrolysis in gluten-free bread, 1491
- Specified risk material
Cross-contamination of CNS tissue due to beef carcass splitting, 83
Lymphoid tissue reduction in bovine small intestines, 1199
Potentially prohibited tissue removal, 573
- Spinach
E-beam radiation for reducing pathogens in spinach, 2415
- Spin-bar digestion
Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
- Spleen
Lactobacillus systemic immune functions and bacterial translocation, 796
- Spoilage
Antimicrobial efficacy and sensory acceptability of plant essential oils, 1846
LAB spoilage of vegetable sausages, 2312
- Spores
Efficacy of household products against *Bacillus* spores, 473
Heat tolerance of *C. sporogenes* spores, 953
Thermal and hydrostatic inactivation, 196
- Sprouting conditions
Growth of *Salmonella* during sprouting, 888
- Sprouts
L. monocytogenes sprout colonization and stress response, 1556
- Stainless steel
Temperature and nutrient effects on *C. jejuni*, 271
- Staphylococcal enterotoxin H
Characteristics of enterotoxin H-producing *S. aureus*, 1855
- Staphylococcal food poisoning
Characteristics of enterotoxin H-producing *S. aureus*, 1855
- Staphylococci
Poultry skin flora after successive washings, 1707
- Staphylococcus*
mecA-positive *Staphylococcus* spp. in raw milk, 186
S. aureus and Domiati cheese, 2567
- Staphylococcus aureus*
Antimicrobial resistance in poultry bacteria, 2537
Detection of *S. aureus*, 191
Pathogen incidence in raw milk, 1580
Sterility testing of an aseptic dispensing valve, 653
- Starter culture
Storage stability of freeze-dried lactic acid bacteria, 431
- Starvation
Altered cell composition in stress-adapted *L. innocua*, 182
- Steamed rice
Proteolytic *C. botulinum* in rice at low pH, 468
- Sterility
Sterility testing of an aseptic dispensing valve, 653
- Stochastic model
A comparison of BSE risk between Japanese and U.S. beef, 802
Model for growth of *Salmonella* on chicken frankfurters, 1135

- Storage
Yeasts from refrigerated commercial shell eggs, 1258
- Storage stability
Storage stability of freeze-dried lactic acid bacteria, 431
- Street food
Microbial quality of stuffed mussels, 616
- Stress
Low-salinity stress in *V. vulnificus*, 416
- Structural component
Heat tolerance of *C. sporogenes* spores, 953
- Stuffed
Salmonellosis from stuffed chicken products, 2153
- Stuffed mussel
Microbial quality of stuffed mussels, 616
- Submerged coil
Modification of the submerged coil, 775
- Subprimals
Blade tenderization of beef containing *E. coli*, 2190
- Subtyping
Evaluation of an automated rep-PCR system, 1372
- Subzero temperatures
High-pressure treatment of *L. monocytogenes* in smoked salmon, 1612
- Superchilling
Superchilled storage of cold-smoked salmon, 2198
- Supercritical fluid extraction
Antimicrobial activity of *D. salina* extracts, 2138
- Supercritical fluid extracts
Acute toxicity of rosemary extracts, 790
- Super shedder
Fecal pats inoculated with *E. coli* O157:H7 to mimic a super shedder, 648
- Surface modification
Rechargeable antimicrobial polyethylene, 2042
- Surface-mold-ripened cheese
Survival of *L. monocytogenes* on surface-mold-ripened soft cheese, 1563
- Surface roughness
Stainless steel surface roughness and finish, 170
- Surrogate
Surrogates for *Salmonella* in meat, 714
- Survey
Domestic refrigeration practices, 1898
- Survival
Listeria inactivation affected by sequences of stresses, 1169
Listeria transfer from knives to turkey breast, 176
L. monocytogenes on salami during storage, 494
Neural networks and survival of *L. monocytogenes* in soft cheese, 750
Survival of *L. innocua* in dairy products, 742
Survival of *L. monocytogenes*, 1835
Survival of murine norovirus 1 on onions and spinach, 1590
V. cholerae survival on different types of cooked rice, 2453
- Survival rate
Storage stability of freeze-dried lactic acid bacteria, 431
- Survival studies
SA11 rotavirus survival in fresh tropical fruit juices, 1035
- Susceptibility
Susceptibility of *Arcobacter* spp. to aromatic compounds, 165
- Sushi
Microbiological quality of sushi, 860
- Sweet potato puree
Microbiology of sweet potato puree, 639
- Swine
Campylobacter in paired turkey and swine farms, 1791
L. monocytogenes in cull sows and pork, 545
- Swine liver
Screening clenbuterol by strip and ELISA, 865
- SYBR Green
Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
- Table olives
Effects of chloride salts on microbial growth, 1412
- Tail muscle
Mercury in bluefin tuna, 595
- Tailing
L. monocytogenes HHP inactivation, 2007
- Tajikistan
Control of *C. perfringens* in Tajik sambusa, 77
- Tandem mass spectrometry
Efficacy of drug test kits for adulterated food, 1007
- TaqMan
Detection of noroviruses in oysters, 1427
- TCID₅₀ (50% tissue culture infective dose)
UV-light inactivation of viruses on fresh produce, 908
- tdc*
Evaluation of *tdc* and its transcript in bacteria and foods, 93
- tdc* expression
Evaluation of *tdc* and its transcript in bacteria and foods, 93
- Temperature
Effect of different water temperatures on knife decontamination, 1338
L. monocytogenes growth at low temperature, 420
Modeling the growth of *Salmonella* in poultry, 2429
Salmonella Enteritidis in eggs, 2073
Temperature and nutrient effects on *C. jejuni*, 271
Toxoplasma in ready-to-eat smallgoods, 1442
- TEMPO system
Automated MPN determination of *Enterobacteriaceae* in food, 376
- Tenax
Role of fat content in trapping volatile compounds, 1889
- Tenderization
Blade tenderization of beef containing *E. coli*, 2190
Heat inactivation of *E. coli* O157:H7 in nonintact beef, 1349
- Terrorism
Efficacy of drug test kits for adulterated food, 1007
- Test-and-hold
Effects of reduced enrichment volumes, 1768
- tet*(M) gene
Antimicrobial susceptibility of *L. paracasei*, 339
- Thai fermented foods
Nisin Z production in *L. lactis*, 2024
- Thailand
Deoxynivalenol contamination in wheat products, 1931
- Thermal death time
Modification of the submerged coil, 775
- Thermal history
Modeling the effect of sublethal history, 279
- Thermal inactivation
Facilitated thermal inactivation of *E. coli* O157:H7, 1604
Heat inactivation of *E. coli* O157:H7 in nonintact beef, 1349
Inactivation of desiccated *E. sakazakii*, 934
Thermal and hydrostatic inactivation, 196
Thermal inactivation of *Salmonella* in turkey breast, 2548
Thermal inactivation of viruses in chicken meat, 1214
Thermal resistance of *F. tularensis*, 2208
- Thermal stability studies
Detection of risin in foods, 2053
- Thermostability
Antibiotics in milk, 1193
- Thermotolerance
Modeling the effect of sublethal history, 279
- Thinning
Campylobacter in thinning of broiler flocks, 264
- Thymol
T. vulgaris and a stored food mite, 351
- Thymoquinone
Influence of extraction methods on *N. sativa* essential oils, 2475
- Thymus vulgaris*
T. vulgaris and a stored food mite, 351
- Tilapia
Radiation D₁₀-values of *L. monocytogenes* on catfish and tilapia, 2278

- Time to growth (TTG)
Antimicrobials and diacetate in RTE meats, 1386
- Tomato
Metabolic effects of *Fusarium*, 1366
Microbiological profile of hydroponic tomatoes, 60
Salmonella in tomatoes, 1572
Salmonella on hydroponic tomatoes, 676
- Tomato juice
Proteolytic activity of molds, 2129
- Toxicity
Ricin and abrin in beverages, 1875
- Toxin
Toxin expression in O157 lineages, 1673
- Transmission electron microscopy
Heat tolerance of *C. sporogenes* spores, 953
- Transportation
Effect of transportation and lairage on risk of *S. enterica* in cattle, 2228
Impact of transportation and lairage on *E. coli* O157:H7 in cattle, 1114
- Trends
Salmonella prevalence in the broiler supply chain, 1974
- Trichinella*
Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
- Trifluoroacetic acid
Occurrence of bound deoxynivalenol in *Fusarium*-infected barley, 1266
- T7 virus
Effect of HPP on freely suspended and bivalve-associated T7, 345
- Turkey
Campylobacter in paired turkey and swine farms, 1791
Campylobacter prevalence in poultry production, 2295
Modeling the effect of sublethal history, 279
Salmonella detection methods on turkey parts, 386
Salmonella migration into marinated Turkey, 153
Thermal inactivation of *Salmonella* in turkey breast, 2548
- Turkey meat
Antimicrobial resistance and class 1 integrons in *E. coli*, 1679
- Turkish white cheese
High-pressure processing of Turkish white cheese, 102
- 12S rRNA
Real-time PCR quantitative detection of bovine DNA in feedstuffs, 564
- Typing
L. monocytogenes in produce processing, 735
Superchilled storage of cold-smoked salmon, 2198
- Tyrophagus putrescentiae*
T. vulgaris and a stored food mite, 351
- Tyrosine decarboxylase
Evaluation of *tdc* and its transcript in bacteria and foods, 93
- Ultrahigh pressure
Antimicrobial synergy between high pressure and Red No. 3, 1861
- Ultrahigh pressure homogenization (UHPH)
Inactivation of *Salmonella* Senftenberg 775W by UHPH, 2283
- Ultrasound
L. monocytogenes in drain pipes, 66
- United States
A comparison of BSE risk between Japanese and U.S. beef, 802
Non-O157 STEC from humans, cattle, and food in the United States, 1023
- Unpasteurized milk
mecA-positive *Staphylococcus* spp. in raw milk, 186
- U.S. Food and Drug Administration
U.S. FDA regulatory management of human listeriosis, 1277
- UV
UV-light inactivation of viruses on fresh produce, 908
- UV-absorbing substances
Membrane damage and inactivation of bacteria in apple juice by RFEF, 684
Thermal inactivation of bacteria in liquid egg, 1988
- UV light
UV light-hydrogen peroxide decontamination of lettuce, 903
- Validation
BAX system PCR method for *Salmonella* versus Brazil's official method, 2442
Cleaning and validation to prevent allergen cross-contact, 445
Controlling pathogens in beef jerky, 918
Model for growth of *Salmonella* on chicken frankfurters, 1135
Validated digestion assay to detect *Trichinella* larvae in horse meat, 558
- Vanilla cream
Modeling *L. monocytogenes* in pasteurized vanilla cream, 1828
- Variety
Salmonella in tomatoes, 1573
- Vegetables
E. coli O157:H7 enrichment in produce, 139
Factors for produce contamination, 2389
Prevalence of *E. coli* O157:H7 in vegetables, 816
Survival of murine norovirus 1 on onions and spinach, 1590
Water disinfection from the fresh-cut vegetable industry, 286
- Vegetable sausage
LAB spoilage of vegetable sausages, 2312
- Vegetative cells
Physiological response of *B. cereus* to food processing treatments, 2168
- Verocytotoxin
Q. infectoria as anti-*E. coli* agent, 1223
- Verotoxin
STEC and *E. coli* O157 from retail meats in China, 706
- Viability
L. monocytogenes adhesion and viability, 1379
- Vibrio*
Antibiotic resistance in *V. parahaemolyticus*, 2552
- Vibrio cholerae*
V. cholerae survival on different types of cooked rice, 2453
- Vibrio parahaemolyticus*
Iceing to control *V. vulnificus* and *V. parahaemolyticus*, 1475
Protein expression in stressed *V. parahaemolyticus*, 2589
- Vibrio vulnificus*
Iceing to control *V. vulnificus* and *V. parahaemolyticus*, 1475
In vitro cold shock gene expression in *Vibrio vulnificus*, 157
Low-salinity stress in *V. vulnificus*, 416
- Virulence
Campylobacter prevalence in poultry production, 2295
- Virulence factor
Antimicrobial resistance and virulence in enterococci, 760
Enterococci from human patients and foods, 2100
Transcription of *stx*₁, *marA*, and *eaeA* in *E. coli* O157:H7, 1469
- Virulence genes
Epidemiologic relevance of MVLST, 2559
Regulation of safety of probiotics, 1734
Virulence genes in *E. coli* O157 and non-O157 isolates, 1774
- Virus
Thermal inactivation of viruses in chicken meat, 1214
- Vitelline membrane
Salmonella Enteritidis growth on egg yolk membranes, 1905
- Washing
Coliforms and foodborne pathogens on spinach in packing plants, 2398
Survival of murine norovirus 1 on onions and spinach, 1590
Water disinfection from the fresh-cut vegetable industry, 286
- Water
Effect of different water temperatures on knife decontamination, 1338
- Water medication
Sulfamethazine water medication pharmacokinetics and contamination, 584
- Water pressurization
Reduction of *Salmonella* Enteritidis on almonds by high-pressure water, 825
- Water-soluble muscadine seed extracts
Antimicrobial effect of muscadine seed extracts, 1465
- Weight of evidence
Application of MCDA to a food safety decision-making problem, 2323

- Western blot
 Comparison of detection methods for CNST, 1505
- Wheat
 Alternaria toxins in wheat, 1262
- Wheat products
 Deoxynivalenol contamination in wheat products, 1931
- Whole muscle
 Thermal inactivation of *Salmonella* in turkey breast, 2548
- WIC participants
 Consumer food safety knowledge and behaviors, 1651
- Wine
 Ochratoxin A adsorption by yeast, 1496
 Ochratoxin A in wine, 1038
 Rapid method used to analyze Ochratoxin A, 2133
- Xanthene derivatives
 Antimicrobial synergy between high pressure and Red No. 3, 1861
- Xanthorrhizol
 Antibacterial activity of xanthorrhizol, 1926
- Xylitol
 Xylitol to reduce microbial adhesion, 405
- Yeast
 Ochratoxin A adsorption by yeast, 1496
 Yeasts from refrigerated commercial shell eggs, 1258
- Yeast cell wall
 Ochratoxin A adsorption by yeast, 1496
- Yersinia enterocolitica*
 Prevalence of *Y. enterocolitica*, 1162
 Y. enterocolitica in ground pork packaged under modified atmospheres, 845
- Yogurt
 Survival of *L. innocua* in dairy products, 742
- Yolk
 Salmonella Enteritidis growth on egg yolk membranes, 1905
- Zearalenone
 LC-TOF-MS analysis of mycotoxins in figs, 1500
- Zoonoses
 Non-O157 STEC from humans, cattle, and food in the United States, 1023
- z-Value
 Heat-resistant fungi from blueberries, 2030
 Storage stability of freeze-dried lactic acid bacteria, 431