

ORIGINAL ARTICLE

Depression, anxiety, and sleep attributes: A cross-sectional study of chiropractic college students

John Ward, DC, MA, MS and Jesse Coats, DC

ABSTRACT

Objective: To measure degree of depression, anxiety, and hours of sleep of chiropractic college students through an anonymous paper survey.

Methods: A convenience sample of 164 chiropractic college students completed a Major Depression Inventory (MDI) survey, a 2-item Generalized Anxiety Disorder (GAD-2) survey, and were asked the number of hours of sleep they had per night and demographic questions. The survey was distributed to trimester 1–6 students 4 weeks into their 15-week semester.

Results: Chiropractic students had an average MDI score of 18.1 ± 10.6 (mean \pm SD), indicating the average chiropractic college student did not display depression. However, subset analysis revealed 18.9% of students had MDI scores over 30, which is associated with possible severe depression. Of students who demonstrated possible severe depression, 80.6% were female. Body mass index demonstrated no correlation with the number of respondents at risk for severe depression. Average GAD-2 score was 3.0 ± 1.9 ; male students scored 2.4 ± 1.8 and female 3.6 ± 1.8 , indicating female students expressed anxiety more. Last, survey respondents reported they averaged 6.3 ± 1.1 hours of sleep per night with females reporting approximately 30 minutes less sleep per night than males.

Conclusion: Chiropractic students, on average, did not display depression. However, a subset of female students near the age of 26 were at an increased risk of severe depression. Chiropractic students displayed a rate of anxiety greater than that of undergraduate college students based on existing data sets and reported approximately 6 hours of sleep per night.

Key Indexing Terms: Chiropractic; Students; Health Occupations; Mental Health; Depression; Anxiety; Sleep

J Chiropr Educ 2024;00(00):000–000 DOI 10.7899/JCE-23-18

INTRODUCTION

Depression rates are increasing across the world, an issue that has been magnified in recent years due to the stress brought on by the COVID-19 pandemic.¹ Experiencing major depression is the strongest risk factor for a person to commit suicide.^{2,3} It is common for individuals suffering from depression to be untreated.^{4,5} One factor that is a barrier to treatment is financial resources, an issue that can prevent college students from seeking appropriate care.⁶ Point prevalence suggests 26.5% of college students suffer from depression at clinically significant levels with 9% of them requiring clinical attention.⁷

Undergraduate college student depression and anxiety levels are shown to negatively impact learning.⁸ Overall, college student anxiety prevalence is 32%.⁹ Furthermore, research on medical school students demonstrates that increased stress

levels negatively impact their academic performance.^{10,11} Medical students who are highly stressed are shown to be more cynical,¹² display decreased empathy,^{13,14} and have a lower desire to provide care to chronically ill patients.¹⁵ Some sources of stress for medical school students are workload,^{16,17} academic pressure,¹¹ sleep deprivation,¹⁶ and financial concerns.¹⁶ All of these factors are likely experienced by chiropractic college students to some degree.

Chiropractic programs are academically and physically stressful.¹⁸ Meckamalil et al¹⁹ found that there was a 19.0% prevalence of depression and 32.6% prevalence of anxiety among undergraduate students at Canadian Memorial Chiropractic College. Similar to other researchers, Kinsinger et al²⁰ found that chiropractic students had elevated levels of depression with female students having higher rates of symptoms.^{19,20} Spegman et al¹⁸ found that the higher reported stress level of a chiropractic student, the more likely the student would feel less confident as an intern. Many lifestyle factors besides academic stress level can impact a chiropractic student's health to include alcohol consumption, exercise

First Published Online September 11 2024

level, caffeine consumption, and the use of tobacco products. Evidence demonstrates that the average chiropractic college student reports drinking alcohol,^{21,22} engaging in regular exercise,²² and not smoking.²² Alcohol consumption sometimes is used as an unhealthy stress-coping mechanism by college students. Furthermore, the proportion of chiropractic students that report eating junk food and consuming 1 or more servings of caffeinated beverages a day is shown to increase during a chiropractic program.²² This is especially true during exam weeks when students are under higher levels of stress.²³

Another variable that can impact mental well-being is body mass index (BMI). Individuals categorized as overweight or obese on BMI charts are shown to have increased rates of depression in some studies.^{24–26} However, this relationship is not seen in all studies^{27–29} and, therefore, warrants additional data collection to help aid in the development of future review articles.

There is little published research on the rate of depression and anxiety among chiropractic college students.^{20,22,30,31} Therefore, the objective of this study was to develop a better understanding of depression, anxiety, and sleep habits of chiropractic college students.

METHODS

This research study was reviewed and approved by the Texas Chiropractic College Institutional Review Board for human subjects in accordance with the Declaration of Helsinki. Multiple existing surveys were reviewed in an effort to develop a 1-page maximum anonymous paper survey that would capture data on student depression, anxiety, hours of sleep per night, and a few respondent questions regarding age, biological sex, height, and weight. The limit of 1 page was intended to reduce survey question fatigue and increase the likelihood that respondents answered all of the questions. The survey was distributed in September 2022, 4 weeks into the 15-week trimester for students in trimesters 1–6 of a chiropractic doctoral program.

Existing surveys on depression and anxiety were screened to be concise, publicly available (not requiring permission for use), and free of charge and such that they had been used in prior research studies. Surveys reviewed and screened against these criteria were the Warwick-Edinburgh Mental Well-being Scale,³² Major Depression Inventory (MDI),³³ World Health Organization-5 Well-Being Index,³⁴ Oxford Happiness Questionnaire,³⁵ Everyday Feelings Questionnaire,³⁶ Mental Health Continuum Short Form,³⁷ Satisfaction with Life Scale,³⁸ Center for Epidemiologic Studies Depression Scale,³⁹ Generalized Anxiety Disorder 2-item (GAD-2),⁴⁰ Generalized Anxiety Disorder 7-item,⁴¹ and Patient Health Questionnaire-9.⁴² From these surveys, researchers chose the MDI; GAD-2; a question about how many hours of sleep the student received per night on average; and the aforementioned questions on age, biologic sex, height, and weight. The MDI response options and scoring for each question are as follows: *all of the time* (5), *most of the time* (4), *more than half of the time* (3), *less than half of the time* (2), *some of the time* (1), and *none of the time* (0). The MDI is composed of 12 questions in total (1–7, 8a, 8b, 9, 10a, 10b) with 10 being graded (using the highest score out of questions 8a and 8b as well as highest out of 10a and 10b). A total score of <20 = *no or doubtful depression*,

20–24 = *mild depression*, 25–29 = *moderate depression*, and 30+ = *severe depression*.⁴³ The MDI is shown to have a sensitivity between 86% and 92% and specificity between 82% and 86% for diagnosis of major depression.⁴⁴ The GAD-2 response options and scoring for each question are as follows: *not at all* (0), *several days* (1), *more than half the days* (2), and *nearly every day* (3). The GAD-2 is composed of 2 questions. A score of 0–3 indicates no to mild anxiety and 4–6 indicates severe anxiety. For scores exceeding 3, the GAD-2 has a sensitivity of 86% and specificity of 83% for diagnosis of generalized anxiety disorder.⁴⁵

The MDI and GAD-2 asked the students to answer based on how they had been feeling over the preceding 2 weeks.

The lead investigator for the research study distributed the survey to 6 different trimester groups of chiropractic students (trimesters 1–6). This was done by coordinating in advance with 1 of their instructors per trimester to attend and distribute the survey during the first 15 minutes of class. Participants were told that completing the survey was entirely voluntary and that, if they completed it, they were giving their implied consent for its use in data analysis. Data were only collected for the purpose of this study and not part of a larger college survey. Students were given the 1-page anonymous survey in class along with a blank cover sheet to shield their answers from their peers. Respondents completed the anonymous survey and then placed the survey and cover sheet in a drop box at the front of the classroom. Once all surveys had been collected across multiple classes, they were transposed to Excel (Microsoft Office).

Statistics

Excel data were analyzed using SPSS version 20.0 (IBM). Results were reported as mean \pm SD unless otherwise specified. Pearson correlation was used to analyze the relationship between BMI and MDI score. Spearman correlation was used to analyze the relationship between trimester number and MDI score.

RESULTS

There were 81 female respondents with the following attributes: 26.4 \pm 5.1 years, 1.66 \pm 0.07 m, 73.8 \pm 15.4 kg, and 6.0 \pm 1.2 hours of reported sleep per night. There were 83 male respondents with the following attributes: 26.3 \pm 4.8 years, 1.78 \pm 0.06 m, 89.2 \pm 17.0 kg, and 6.6 \pm 0.9 hours of reported sleep per night.

Tables 1 and 2 illustrate the survey results per question. Chiropractic students averaged an MDI score of 18.1 \pm 10.6 out of a maximum score of 50, indicating the average chiropractic college student did not display depression. However, 18.9% of students were scored to have possible severe depression according to MDI scores. Of students that demonstrated possible severe depression, 80.6% were female. The average age of the female students that displayed possible severe depression was 25.9 years of age. The 2 MDI questions with the highest response scores were “Have you suffered from increased appetite?” with a score of 2.52 \pm 1.09 and “Have you felt lacking in energy and strength?” with a score of 2.40 \pm 1.35. The MDI question with the lowest response score was “Have you felt that life wasn’t worth living?” with a score of 0.54 \pm 1.11.

Table 1 - Major Depression Inventory Results

How Much of the Time ...	Student Average Score on a Scale of 0-5	Student Average Score on a Scale of 0-5					
		0	1	2	3	4	5
1. Have you felt in low spirits or sad?	1.81 + 1.10	9.8%	33.5%	29.9%	20.7%	4.9%	1.2%
2. Have you lost interest in your daily activities?	1.66 + 1.35	22.6%	30.5%	19.5%	15.2%	10.4%	1.8%
3. Have you felt lacking in energy and strength?	2.40 + 1.35	6.7%	22.6%	25.0%	22.0%	17.7%	6.1%
4. Have you felt less self-confident?	1.91 + 1.47	17.1%	31.7%	16.5%	18.3%	10.4%	6.1%
5. Have you had a bad conscience or feelings of guilt?	1.27 + 1.43	42.7%	21.3%	14.0%	12.8%	6.1%	3.1%
6. Have you felt that life wasn't worth living?	0.54 + 1.11	75.0%	9.8%	6.1%	4.9%	3.7%	0.6%
7. Have you had difficulty in concentrating, eg, when reading the newspaper or watching television?	2.10 + 1.53	16.5%	27.4%	14.0%	21.3%	13.4%	7.3%
8a. Have you felt very restless?	2.34 + 1.57	16.5%	20.1%	17.1%	20.1%	17.7%	8.5%
8b. Have you felt subdued?	1.57 + 1.44	39.0%	23.8%	20.7%	9.8%	3.7%	3.1%
9. Have you had trouble sleeping at night?	2.23 + 1.59	21.3%	26.8%	18.9%	14.0%	8.5%	10.4%
10a. Have you suffered from reduced appetite?	1.85 + 1.65	39.0%	25.0%	11.0%	11.0%	7.9%	6.1%
10b. Have you suffered from increased appetite?	2.52 + 1.09	46.3%	20.7%	14.6%	11.6%	6.1%	0.6%

The included survey questions asked about how the student had been feeling over the preceding 2 weeks. Student average score data listed as mean ± SD for the question, followed by the relative frequency percentage of each response option on the right columns out of 100% (0 = none of the time, 5= all of the time, etc.). The highest scoring question (10b) was students reporting increased appetite.

Based on BMI, the average chiropractic college student in this study was overweight (27.4 ± 5.1). Survey respondents at risk for severe depression based on their MDI score possessed the following BMI categories: healthy weight (18.5–24.9 BMI, n = 15), overweight (25.0–29.9 BMI, n = 10), and obese (30.0+ BMI, n = 6). Pearson correlation between BMI and MDI scores of individuals categorized as likely having severe depression demonstrated no correlation ($r(31) = .142, p = .447$); thus, higher BMI did not mean a higher MDI score.

MDI scores per trimester were tri-1 = 17.9 ± 10.0, tri-2 = 17.9 ± 10.5, tri-3 = 17.1 ± 10.7, tri-4 = 19.0 ± 10.2, tri-5 = 16.4 ± 11.9, and tri-6 = 20.0 ± 11.6. Spearman did not demonstrate a correlation between trimester number and MDI score among participants categorized as likely severely depressed ($r(31) = -.250, p = .175$). Thus, no trend was seen in which students in earlier trimesters had high MDI levels and students in higher trimesters possessed lower MDI levels possibly due to any type of adaptation to academic load. Students were taking the following number of hours on average per trimester: trimester 1: 24 hours, trimester 2: 30 hours, trimester 3: 25 hours, trimester 4: 29 hours, trimester 5: 28 hours, and trimester 6: 26 hours.

Average GAD-2 score was 3.0 ± 1.9 indicative that the average chiropractic college student did display some degree of generalized anxiety. The proportion of respondents that displayed generalized anxiety was 53.7%. Upon further sub-analysis of GAD-2 score by sex, males demonstrated a score of 2.4 ± 1.8 and females a score of 3.6 ± 1.8 with female respondents demonstrating anxiety. Chiropractic students averaged 6.3 ± 1.1 hours of sleep per night with females reporting approximately 30 minutes less sleep than males on average.

DISCUSSION

After the data were analyzed, students in each trimester class were reminded that the college offers free online counseling services to students in case they ever feel anxiety and/or depression. For perspective, depression rates for the general U.S. population are 7%.⁴⁶ Medical school students typically have an MDI between 15 and 16, which is less than what was demonstrated by students in this study.^{47,48} Usually, medical school students have an undergraduate grade point average of nearly 3.75 and are accustomed to elevated levels of academic stress.⁴⁹ The typical grade point average of a student entering into a U.S. chiropractic college is about 3.15.⁵⁰ Perhaps chiropractic students may have a decreased ability to cope with life stress compared with medical school students. This point should be further considered when looking at the MDI score of general undergraduate students. Undergraduate college students demonstrate an MDI score of 17.3⁵¹ with the students in this study still scoring higher than that.

Female chiropractic college students were more likely to have higher scores on the MDI scale than male students. Studies of nursing, medical, and chiropractic college school students similarly demonstrate that female students are more likely to suffer from depression.^{20,48,52-54} Some of the factors suggested in research to be responsible for females demonstrating higher rates of depression are genetics, increased need for social support, and pressures of family life.⁵⁵

The results of this study demonstrate that the average chiropractic college student was not suffering from depression but that a proportion of the overall student population was. In Kinsinger's work on the same topic, he found that 22%–25% of chiropractic

Table 2 - Generalized Anxiety Disorder-2 Results

Statements	Student Average Score on a Scale of 0-3
1. Feeling nervous, anxious, or on edge.	1.58 + 0.98
2. Not being able to stop or control worrying.	1.38 + 1.05

students were suffering from some degree of mild-to-severe depression in any given year in a chiropractic program.²⁰ These numbers are significantly higher than what are seen in general and postsecondary colleges.⁵⁶ Countermeasures to aid chiropractic students who may be suffering from depression should be pursued.

In this study, the students surveyed scored higher on the anxiety scale than average undergraduate college students with a score of 3.0 ± 1.9 . Average undergraduate students demonstrate a GAD-2 score of 2.76.⁵⁷ The GAD-2 question with the highest score was about “feeling nervous, anxious, or on edge?” with a score of 1.58 ± 0.98 . These findings of elevated anxiety among chiropractic college students were similar to what Spelman et al¹⁸ found in their research on this topic. The Spelman study demonstrated that chiropractic students had a comparable anxiety level to medical school students.¹⁸

Based on BMI, the average chiropractic student in this study was overweight. The degree to which college students entering into chiropractic programs are overweight and their progression of possibly gaining weight warrants further study. DuMonthier et al³⁰ demonstrate that, due to the stressful academic nature of chiropractic programs, students report having less time for physical activity, which can be a risk factor for gaining weight. Additional attributes that may impact weight gain throughout a chiropractic program are the consumption of junk food as well as alcohol as unhealthy means of dealing with stress.⁵⁸ In this study, BMI was not shown to be linked to an increased rate of depression.

The health habits and methods of chiropractic students of dealing with their own level of anxiety and depression is paramount to how they will interact with their future patients. Studies suggest that, if a chiropractic student has less effective internal coping mechanisms, they will be less likely to promote healthy behaviors to their patients.²² This is seen in the study by DuMonthier et al³⁰ in which obese chiropractic students were less likely to assign importance to role modeling and patient education for their patients. Chiropractic programs should consider what mechanisms they have in place to aid their students in dealing with stress and anxiety because it will have a subsequent, trickle-down impact on their future patients.

Chiropractic students in this study averaged $6.3 + 1.1$ hours of sleep per night. This value is similar to that reported by medical school students, which ranges between 5.9 ± 1.6 hours and 6.9 ± 1.1 hours of sleep per night.⁵⁹⁻⁶¹

Possible future directions for this study would be to (1) perform a similar study at other chiropractic college campuses to see if the rates of severe depression and anxiety are similar, (2) determine if the rate of depression and anxiety improves as students progress through a chiropractic program to the end, and (3) measure weight gain of students as they progress through a chiropractic program related to stress eating. Family support and personalized interventions are shown to aid college students in decreasing their level of depression.⁶² Although colleges may not be able to impact family support, they could incorporate more organized physical fitness and social activities on campuses that could impact depression and/or anxiety levels.

Strengths and Limitations

A strength of this study is the use of validated survey instruments to measure depression and anxiety degree.^{63,64} This will make it easier for future researchers to replicate portions of this study and compare their results through a standardized method.

A limitation of this study is that it did not track a cohort of students as they progressed through the chiropractic program. Doing this could more effectively show if students are able to adapt to their stress level over time. This point was actually demonstrated by Ward et al²² and Kinsinger et al,²⁰ who show a decrease in the number of students reporting depressive symptoms by the end of the chiropractic program, which is in contrast to what is seen among medical school students.^{53,65,66}

A second limitation of this study is that it was performed at 1 chiropractic college on a small group of students, which limits external validity. As more studies are performed at multiple chiropractic colleges, a more definitive understanding of the mental health of chiropractic students will be gained.

This study did not measure alcohol use and its relationship with depression. Alcohol can be used as a coping mechanism by students in high-stress college programs.⁶⁷ Alcohol use as a student is also predictive of alcohol use as a future health care provider, which could impact care of patients.⁶⁸

Another limitation of this study is that it occurred post-COVID, and that could impact the results and their interpretation in relation to preexisting data sets on depression and anxiety rates of college students. COVID had a significant impact on student mental health.⁶⁹⁻⁷¹ The degree to which students in this study may have been dealing with the lingering effects of the worldwide COVID-19 pandemic in September of 2022 is unclear.

CONCLUSION

Chiropractic students, on average, did not display depression. The students did score higher on the MDI scale than undergraduate college students and medical school students. Future studies should determine if these findings can be corroborated and explore why the differences exist. Chiropractic college students did display a rate of anxiety greater than that of typical undergraduate college students. Chiropractic college programs involve students taking on more hours per week than the typical undergraduate student, and this may partially account for the increase in student anxiety level seen in this study.

FUNDING AND CONFLICTS OF INTEREST

This work was funded internally. The authors have no conflicts of interest to declare relevant to this work.

About the Authors

John Ward (corresponding author) is a professor in the department of foundational sciences at Texas Chiropractic College (5912 Spencer Hwy., Pasadena, TX 77505; jward@txchiro.edu). Jesse Coats is a professor and second year chair at Texas Chiropractic College (5912 Spencer Hwy., Pasadena, TX 77505; jcoats@txchiro.edu). This article was received May 4, 2023; revised January 13, 2024; and accepted March 14, 2024.

Author Contributions

Concept development: JW, JC. Design: JW, JC. Supervision: JW, JC. Data collection/processing: JW, JC. Analysis/interpretation: JW, JC. Literature search: JW. Writing: JW. Critical review: JW, JC.

© 2024 Association of Chiropractic Colleges

REFERENCES

1. Goodwin R, Dierker L, Wu M, Galea S, Hoven C, Weinberger A. Trends in U.S. depression prevalence from 2015 to 2020: The widening treatment gap. *Am J Prev Med.* 2022;63(5):726–733. doi:10.1016/j.amepre.2022.05.014
2. Chang S, Stuckler D, Yip P, Gunnell D. Impact of 2008 global economic crisis on suicide time trend study in 54 countries. *BMJ.* 2013;347:f5239. doi:10.1136/bmj.f5239
3. Hasin D, Goodwin R, Stinson F, Grant B. Epidemiology of major depressive disorder: results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Arch Gen Psychiatry.* 2005;62(10):1097–1106. doi:10.1001/archpsyc.62.10.1097
4. Whiteford H, Harris M, McKeon G, et al. Estimating remission from untreated major depression: a systematic review and meta-analysis. *Psychol Med.* 2013;43(8):1569–1585. doi:10.1017/S0033291712001717
5. Riolo S, Baltrus P, Ye J, Rust G. Prevalence of depression by race/ethnicity: findings from the National Health and Nutrition Examination Survey III. *Am J Public Health.* 2005;95(6):998–1000. doi:10.2105/AJPH.2004.047225
6. Harman J, Edlund M, Fortney J. Disparities in the adequacy of depression treatment in the United States. *Psychiatr Serv.* 2004;55(12):1379–1385. doi:10.1176/appi.ps.55.12.1379
7. Shah T, Pol T. Prevalence of depression and anxiety in college students. *J Mental Health Human Behavior.* 2020;25(1):10–13. doi:10.4103/jmhbb.jmhbb_16_20
8. Edberg R, Elasowich B. Student mental health, diminishing resources: two major challenges for urban community colleges. *Commun Coll Lead.* 2006;1(7):1.
9. Tan G, Soh X, Hartanto A, Goh A, Majeed N. Prevalence of anxiety in college and university students: an umbrella review. *J Affective Disorders Reports.* 2023;14:100658. doi:10.1016/j.jadr.2023.100658
10. Spiegel D, Smolen R, Hopfensperger K. Medical student stress and clerkship performance. *J Med Educ.* 1986;61(11):929–931. doi:10.1097/00001888-198611000-00015
11. Stewart S, Lam T, Betson C, Wong C, Wong A. A prospective analysis of stress and academic performance in the first two years of medical school. *Med Educ.* 1999;33(4):243–250. doi:10.1046/j.1365-2923.1999.00294.x
12. Woloschuk W, Harasym P, Temple W. Attitude change during medical school: a cohort study. *Med Educ.* 2004;38(5):522–534. doi:10.1046/j.1365-2929.2004.01820.x
13. Hojat M, Mangione S, Nasca T, et al. An empirical study of decline in empathy in medical school. *Med Educ.* 2004;38(9):934–941. doi:10.1111/j.1365-2929.2004.01911.x
14. Thomas M, Dyrbye L, Huntington J, et al. How do distress and well-being relate to medical student empathy? A multicenter study. *J Gen Intern Med.* 2007;22(2):177–183. doi:10.1007/s11606-006-0039-6
15. Griffith C, Wilson J. The loss of idealism throughout internship. *Eval Health Prof.* 2003;26(4):415–426. doi:10.1177/0163278703258107
16. Wolf T, Faucett J, Randall H, Balson P. Graduating medical students' ratings of stresses, pleasures, and coping strategies. *Med Educ.* 1988;63(8):636–642. doi:10.1097/00001888-198808000-00008
17. Guthrie E, Black D, Shaw C, et al. Embarking upon a medical career: psychological morbidity in first year medical students. *Med Educ.* 1995;29(5):337–341. doi:10.1111/j.1365-2923.1995.tb00022.x
18. Spegman A, Herrin S. Chiropractic interns' perceptions of stress and confidence. *J Chiropr Educ.* 2007;21(2):129–137. doi:10.7899/1042-5055-21.2.129
19. Meckamalil C, Brodie L, Hogg-Johnson S, Carroll L, Jacobs C, Côté P. The prevalence of anxiety, stress and depressive symptoms in undergraduate students at the Canadian Memorial Chiropractic College. *J Am Coll Health.* 2022;70(2):371–376. doi:10.1080/07448481.2020.1751173
20. Kinsinger S, Puhl A, Reinhart C. Depressive symptoms in chiropractic students: a 3-year study. *J Chiropr Educ.* 2011;25(2):142–150. doi:10.7899/1042-5055-25.2.142
21. Bougie J, Singh B. Screening for harmful and hazardous drinking by chiropractic students. *J Chiropr Educ.* 2005;19(1):85–90.
22. Ward K, Kallop K, Fernando S, Smith M. Mental health and lifestyle behaviors of students in a doctor of chiropractic program. *J Chiropr Educ.* 2023;37(1):1–6. doi:10.7899/JCE-21-30
23. Jaremków A, Markiewicz-Górka I, Pawlas K. Assessment of health condition as related to lifestyle among students in the examination period. *Int J Occup Med Environ Health.* 2020;33(3):339–351. doi:10.13075/ijomh.1896.01563
24. Badillo N, Khatib M, Kahar P, Khanna D. Correlation between body mass index and depression/depression-like symptoms among different genders and races. *Cureus.* 2022;14(2):e21841. doi:10.7759/cureus.21841
25. Becker E, Margraf J, Türke V, Soeder U, Neumer S. Obesity and mental illness in a representative sample of young women. *Int J Obesity.* 2001;25:S5–S9. doi:10.1038/sj.ijo.0801688
26. Baumeister H, Härter M. Mental disorders in patients with obesity in comparison with healthy probands. *Int J Obes.* 2007;31:1155–1164. doi:10.1038/sj.ijo.0803556
27. Han T, Tijlhuis M, Lean M, Seidell J. Quality of life in relation to overweight and body fat distribution. *Am J Public Health.* 1998;88:1814–1820. doi:10.2105/AJPH.88.12.1814
28. John U, Meyer C, Rumpf H, Hapke U. Relationships of psychiatric disorders with overweight and obesity in an adult general population. *Obes Res.* 2005;13:101–109. doi:10.1038/oby.2005.13
29. Kress A, Peterson M, Hartzell M. Association between obesity and depressive symptoms among U.S. military active duty service personnel 2002. *J Psychosom Res.* 2006;60:263–271. doi:10.1016/j.jpsychores.2005.08.011
30. DuMonthier W, Haneline M, Smith M. Survey of health attitudes and behaviors of a chiropractic college population. *J Manipulative Physiol Ther.* 2009;32(6):477–484. doi:10.1016/j.jmpt.2009.06.001
31. Zhang N, Henderson C. Predicting stress and test anxiety among 1st-year chiropractic students. *J Chiropr Educ.* 2019;33(2):133–139. doi:10.7899/JCE-18-11
32. Hoffman S, Rueda H, Lambert M. Confirmatory factor analysis of the Warwick-Edinburgh Mental Wellbeing Scale among youth in Mexico. *IJSSW.* 2019;62(1):309–315. doi:10.1177/0020872817727576
33. Olsen L, Jensen D, Noerholm V, Martiny K, Bech P. The internal and external validity of the Major Depression Inventory in measuring severity of depressive states. *Psychol Med.* 2003;33(2):351–356. doi:10.1017/s0033291702006724
34. Topp C, Østergaard S, Søndergaard S, Bech P. The WHO-5 Well-Being Index: A systematic review of the literature. *Psychother Psychosom.* 2015;84(3):167–169. doi:10.1159/000376585
35. Hills P, Argyle M. The Oxford Happiness Questionnaire: a compact scale for measurement of psychological well-being.

- Pers Individ Differ.* 2002;33(7):1073–1082. doi:10.1016/S0191-8869(01)00213-6
36. Uher R, Goodman R. The Everyday Feeling Questionnaire: the structure and validation of a measure of general psychological well-being and distress. *Soc Psychiatry Psychiatr Epidemiol.* 2010;45(3):413–423. doi:10.1007/s00127-009-0074-9
 37. Lamers S, Westerhof G, Bohlmeijer E, Klooster P, Keyes C. Evaluating the psychometric properties of the Mental Health Continuum-Short Form (MHC-SF). *J Clin Psychol.* 2011; 67(1):99–110. doi:10.1002/jclp.20741
 38. Diener E, Emmons R, Larsen R, Griffin S. The Satisfaction with Life Scale. *J Pers Assess.* 1985;49(1):71–75. doi:10.1207/s15327752jpa4901_13
 39. Radloff L. The use of the Center for Epidemiologic Studies Depression Scale in adolescents and young adults. *J Youth Adolesc.* 1991;20(2):149–166. doi:10.1007/BF01537606
 40. Hughes A, Dunn K, Chaffee T, Bhattari J, Beier M. Diagnostic and clinical utility of the GAD-2 for screening anxiety symptoms in individuals with multiple sclerosis. *Arch Phys Med Rehabil.* 2018;99(10):2045–2049. doi:10.1016/j.apmr.2018.05.029
 41. Sapra A, Bhandari P, Sharma S, Chanpura T, Lopp L. Using Generalized Anxiety Disorder-2 (GAD-2) and GAD-7 in a primary care setting. *Cureus.* 2020;12(5):e8224. doi:10.7759/cureus.8224
 42. Manea L, Gilbody S, McMillan D. A diagnostic meta-analysis of the Patient Health Questionnaire-9 (PHQ-9) algorithm scoring method as a screen for depression. *Gen Hosp Psychiatry.* 2015;37(1):67–75. doi:10.1016/j.genhosppsych.2014.09.009
 43. Peter Cogen. Major Depression Inventory (MDI): scoring key. 2013. Accessed November 15, 2023. <https://www.therapiebreve.be/documents/mdi-scoring-key.pdf>
 44. Bech P, Rasmussen N, Olsen L, Noerholm V, Abildgaard W. The sensitivity and specificity of the Major Depression Inventory, using the Present State Examination as the index of diagnostic validity. *J Affect Disord.* 2001;66(2–3):159–164. doi:10.1016/S0165-0327(00)00309-8
 45. Kroenke K, Spitzer R, Williams J, Monahan P, Lowe B. Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Ann Intern Med.* 2007;146:317–325.
 46. Substance Abuse and Mental Health Services Administration. Behavioral health trends in the United States: Results from the 2014 National Survey On Drug Use And Health (NSDUH). Accessed February 1, 2023. <https://www.samhsa.gov/data/report/behavioral-health-trends-united-states-results-2014-national-survey-drug-use-and-health>
 47. Akhtar M, Herwig B, Faize F. Depression and anxiety among international medical students in Germany: the predictive role of coping styles. *J Pak Med Assoc.* 2019;69(2):230–234.
 48. Dahlin M, Nils J, Bo R. Stress and depression among medical students: a cross-sectional study. *Med Educ.* 2005;39(6):594–604. doi:10.1111/j.1365-2929.2005.02176.x
 49. Kowarski I. What college GPA is needed for med school? U.S. News. Accessed February 15, 2023. <https://www.usnews.com/education/best-graduate-schools/top-medical-schools/articles/2018-10-02/how-high-of-a-college-gpa-is-necessary-to-get-into-medical-school>
 50. Campus Reel. What GPA do you need to get into Texas Chiropractic College? Foundation Inc. Campus Reel. Accessed February 18, 2023. <https://www.campusreel.org/how-to-get-into-college/texas-chiropractic-college-foundation-inc-admissions-requirements-and-gpa#:~:text=What%20GPA%20Do%20You%20Need,Inc's%20average%20GPA%20is%203.15>
 51. Murray A, Marenus M, Cahuas A, et al. The impact of web-based physical activity interventions on depression and anxiety among college students: randomized experimental trial. *JMIR Form Res.* 2022;6(4):e31839. doi:10.2196/31839
 52. Christensson A, Vaez M, Dickman P, Runeson B. Self-reported depression in first-year nursing students in relation to sociodemographic and educational factors: a nationwide cross-sectional study in Sweden. *Soc Psychiatry Psychiatr Epidemiol.* 2011;46(4):299–310. doi:10.1007/s00127-010-0198-y
 53. Goebert D, Thompson D, Takeshita J, et al. Depressive symptoms in medical students and residents: a multischool study. *Acad Med.* 2009;84(2):236–241. doi:10.1097/ACM.0b013e31819391bb
 54. Dyrbye L, Thomas M, Shanafelt T. Systematic review of depression, anxiety, and other indicators of psychological distress among U.S. and Canadian medical students. *Acad Med.* 2006;81(4):354–373. doi:10.1097/00001888-200604000-00009
 55. Piccinelli M, Wilkinson G. Gender differences in depression: critical review. *Br J Psychiatry.* 2000;177:486–492. doi:10.1192/bjp.177.6.486
 56. American College Health Association. National college health assessment spring 2008 reference group data report (abridged). *J Am College Health.* 2009;57(5):477–488. doi:10.3200/JACH.57.5.477-488
 57. Coakley K, Cargas S, Walsh-Dilley M, Mechler H. Basic needs insecurities are associated with anxiety, depression, and poor health among university students in the state of New Mexico. *J Community Health.* 2022;47(3):454–463. doi:10.1007/s10900-022-01073-9
 58. Medisaukaite A, Kamau C. Does occupational distress raise the risk of alcohol use, binge-eating, ill health and sleep problems among medical doctors? A UK cross-sectional study. *BMJ Open.* 2019;9(5):e027362. doi:10.1136/bmjopen-2018-027362
 59. Bahammam A, Al-Khairy O, Al-Taweel A. Sleep habits and patterns among medical students. *Neurosciences.* 2005;10(2):159–162.
 60. Huen L, Chan T, Yu W, Wing Y. Do medical students in Hong Kong have enough sleep? *Sleep Biol Rhythms.* 2007; 5:226–230.
 61. Džaferović A, Ulen K. Sleep habits among medical students and correlation between sleep quality and academic performance. *Eur J Public Health.* 2018;28(4):358. doi:10.1093/eurpub/cky214.141
 62. Liu X, Guo Y, Zhang W, Gao W. Influencing factors, prediction and prevention of depression in college students: a literature review. *World J Psychiatry.* 2022;12(7):860–873. doi:10.5498/wjp.v12.i7.860
 63. Plummer F, Manea L, Trepel D, McMillan D. Screening for anxiety disorders with the GAD-7 and GAD-2: a systematic review and diagnostic metaanalysis. *Gen Hosp Psychiatry.* 2016;39:24–31. doi:10.1016/j.genhosppsych.2015.11.005
 64. Nielsen M, Ørnboel E, Bech P, Vestergaard M, Christensen K. The criterion validity of the web-based Major Depression Inventory when used on clinical suspicion of depression in primary care. *Clin Epidemiol.* 2017;9:355–365. doi:10.2147/CLEP.S132913
 65. Stewart S, Betson C, Lam T, et al. Predicting stress in first year medical students: a longitudinal study. *Med Educ.* 1997; 31(3):163–168. doi:10.1111/j.1365-2923.1997.tb02560.x
 66. Guthrie E, Black D, Bagalkote H, et al. Psychological stress and burnout in medical students: a five-year prospective

- longitudinal study. *J R Soc Med.* 1998;91(5):237–243. doi:10.1177/014107689809100502
67. Frank E, Elon L, Naimi T, Brewer R. Alcohol consumption and alcohol counselling behavior among US medical students: cohort study. *BMJ.* 2008;337:a2155. doi:10.1136/bmj.a2155
68. Moore R, Mead L, Pearson T. Youthful precursors of alcohol abuse in physicians. *Am J Med.* 1990;88(4):332–336. doi:10.1016/0002-9343(90)90485-v
69. Son C, Hegde S, Smith A, Wang X, Sasangohar F. Effects of COVID-19 on college students' mental health in the United States: interview survey study. *J Med Internet Res.* 2020; 22(9):e21279. doi:10.2196/21279
70. Yang C, Chen A, Chen Y. College students' stress and health in the COVID-19 pandemic: The role of academic workload, separation from school, and fears of contagion. *PLoS One.* 2021;16(2):e0246676. doi:10.1371/journal.pone.0246676
71. Barbayannis G, Bandari M, Zheng X, Baquerizo H, Pecor K, Ming X. Academic stress and mental well-being in college students: correlations, affected groups, and COVID-19. *Front Psychol.* 2022;13:886344. doi:10.3389/fpsyg.2022.886344