



# Cystatin C- and Creatinine-Based Estimated Glomerular Filtration Rate, Vascular Disease, and Mortality in Persons With Diabetes in the U.S.

Ching-Wei Tsai,<sup>1</sup> Morgan E. Grams,<sup>2</sup>  
Lesley A. Inker,<sup>3</sup> Josef Coresh,<sup>1,2</sup>  
and Elizabeth Selvin<sup>1,2</sup>

## OBJECTIVE

Serum cystatin C is an alternative to serum creatinine for estimating glomerular filtration rate (GFR), since cystatin C is less influenced by age and muscle mass. Among persons with diabetes, we compared the performance of GFR estimated using cystatin C (eGFR<sub>cys</sub>) with that using creatinine (eGFR<sub>cr</sub>) for the identification of reduced kidney function and its association with diabetes complications.

## RESEARCH DESIGN AND METHODS

We analyzed data from adult participants from the 1999–2002 National Health and Nutrition Examination Survey with available cystatin C ( $N = 4,457$ ). Kidney function was dichotomized as preserved (eGFR  $\geq 60$  mL/min/1.73 m<sup>2</sup>) or reduced (eGFR  $< 60$  mL/min/1.73 m<sup>2</sup>) using the 2012 Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) cystatin C and the 2009 CKD-EPI creatinine equations.

## RESULTS

Among 778 persons with diabetes, the prevalence of reduced kidney function was 16.5% using eGFR<sub>cr</sub> and 22.0% using eGFR<sub>cys</sub>. More persons with diabetes were reclassified from preserved kidney function by eGFR<sub>cr</sub> to reduced kidney function by eGFR<sub>cys</sub> than persons without diabetes (odds ratio 3.1 [95% CI 1.9–4.9],  $P < 0.001$ ). The associations between lower eGFR and higher prevalence of albuminuria, retinopathy, peripheral arterial disease, and coronary artery disease were robust regardless of filtration marker. Similarly, the risk of all-cause mortality increased with lower eGFR<sub>cr</sub> and eGFR<sub>cys</sub>. Only lower eGFR<sub>cys</sub> was significantly associated with cardiovascular mortality.

## CONCLUSIONS

More persons with diabetes had reduced kidney function by eGFR<sub>cys</sub> than by eGFR<sub>cr</sub>, and lower eGFR<sub>cys</sub> was strongly associated with diabetes complications. Whether eGFR<sub>cys</sub> is superior to eGFR<sub>cr</sub> in approximating true kidney function in a diabetic population requires additional study.

*Diabetes Care* 2014;37:1002–1008 | DOI: 10.2337/dc13-1910

<sup>1</sup>Department of Epidemiology, Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD

<sup>2</sup>Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD

<sup>3</sup>Department of Medicine, Tufts Medical Center, Boston, MA

Corresponding author: Morgan E. Grams, mgrams2@jhmi.edu.

Received 12 August 2013 and accepted 15 November 2013.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc13-1910/-/DC1>.

A slide set summarizing this article is available online.

C.-W.T. and M.E.G. contributed equally to this work.

© 2014 by the American Diabetes Association. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Diabetes is the leading cause of chronic kidney disease (CKD) in developed countries, including the U.S. (1,2). Diabetic kidney disease accounts for 40% of prevalent CKD and 50% of incident end-stage renal disease, and it has increased in direct proportion to the increasing prevalence of diabetes (2–4). People with diabetes often suffer from microvascular and macrovascular complications, including retinopathy, nephropathy, coronary artery disease, peripheral arterial disease, and stroke, as well as early mortality (5,6). Compared with persons with diabetes and preserved kidney function, those with diabetes and CKD face even higher risks of morbidity and mortality. Indeed, both reduced kidney function and albuminuria are independent predictors for cardiovascular disease as well as all-cause mortality (7,8). Accurate estimation of glomerular filtration rate (GFR) and identification of CKD are important.

In clinical practice, kidney function is estimated rather than measured. Glomerular filtration rate estimated using serum creatinine (eGFR<sub>cr</sub>) is the most common approach; however, creatinine is influenced by age, muscle mass, sex, and race (9). Given these limitations, serum cystatin C has been proposed as an alternative filtration marker (10). Cystatin C, an endogenous protein believed to be produced by all nucleated cells, is less affected by age, race, and muscle mass and, in the general population, associates more strongly with all-cause and cardiovascular mortality than does serum creatinine (11,12). However, BMI, diabetes, and inflammation may affect cystatin C levels independent of kidney function (13). Given the high prevalence of obesity in the population with diabetes, as well as the suggestion that cystatin may perform differently in patients with diabetes, there is controversy as to whether cystatin C–based or creatinine–based estimated GFR (eGFR) equations should be used to estimate kidney function in this population (14,15). Furthermore, it is unknown whether the associations of diabetes complications with kidney function estimated using cystatin C (eGFR<sub>cys</sub>) are similar to those observed using eGFR<sub>cr</sub>.

Using nationally representative data from the 1999–2002 National Health and Nutrition Examination Survey (NHANES), we estimated the prevalence of reduced kidney function (eGFR <60 mL/min/1.73 m<sup>2</sup>) among persons with diabetes using the 2012 CKD-EPI cystatin C (16) and 2009 CKD-EPI creatinine (17) equations and investigated the discordance in CKD classification by the two filtration markers. We also compared the associations of eGFR<sub>cr</sub> and eGFR<sub>cys</sub> with prevalent complications of diabetes, including albuminuria, peripheral arterial disease, retinopathy, and coronary artery disease, as well as incident all-cause and cardiovascular mortality.

## RESEARCH DESIGN AND METHODS

### Study Population

NHANES is an ongoing cross-sectional, multistage, stratified, clustered probability sample of the U.S. civilian noninstitutionalized population conducted by the National Center for Health Statistics. Cystatin C concentrations were measured in a subsample of the NHANES 1999–2002 participants aged 12 years and older who were not missing serum creatinine (18). For the current study, we included all participants in the cystatin C subsample aged 20 years or older ( $N = 4,457$ ; 778 of whom had diabetes). The ankle-brachial index (ABI), used to define peripheral arterial disease, was measured only in persons aged 40 years or older ( $N = 556$  of the 778 participants with diabetes, 10 of whom with ABI >1.5 were excluded owing to concern for calcified atherosclerosis) (19,20).

### Assessment of Diabetes and Kidney Function

We classified persons as having diabetes if they reported a physician diagnosis of diabetes, took antidiabetes pills or insulin injections, or had a glycated hemoglobin (HbA<sub>1c</sub>) value of  $\geq 6.5\%$ . No distinction was made with regard to type of diabetes. Preserved/reduced kidney function was defined as eGFR  $\geq 60$  mL/min/1.73 m<sup>2</sup> / eGFR <60 mL/min/1.73 m<sup>2</sup>, estimated using standardized creatinine and cystatin C values and the CKD-EPI 2009 and 2012 equations, respectively (16,17). The term

“advanced CKD” was used to indicate CKD stage 4 or 5 (eGFR <30 mL/min/1.73 m<sup>2</sup>). Creatinine values from NHANES 1999–2000 were standardized [standard creatinine (mg/dL) = 0.147 + 1.013 × (NHANES 1999–2000 uncalibrated serum creatinine [mg/dL]), whereas no correction to the creatinine values in the 2001–2002 survey was needed (21,22). Cystatin C values were recalibrated and standardized to the International Federation of Clinical Chemistry and Laboratory Medicine (IFCC) standard: IFCC standard cystatin C (mg/L) = 1.12 × (cystatin C [mg/L] – 0.12) (22).

### Other Variables of Interest

Hypertension was defined as mean systolic pressure  $\geq 140$  mmHg, mean diastolic pressure  $\geq 90$  mmHg, self-reported hypertension, or the use of an antihypertensive medication. Hyperlipidemia was defined as total cholesterol concentration  $\geq 200$  mg/dL. Low HDL was defined as serum HDL <40 mg/dL. Information on age, sex, race, and smoking was self-reported. Smoking status was determined using answers to the questions, “Have you smoked at least 100 cigarettes in your life?” and “Do you now smoke cigarettes?”

### Prevalent Micro- and Macrovascular Outcomes

Coronary artery disease was defined on the basis of a self-reported history of coronary heart disease, angina, or previous heart attack. Albuminuria was defined as a urinary albumin-to-creatinine ratio (ACR)  $\geq 30$  mg/g. Peripheral arterial disease was defined by an ABI <0.90 in either leg (23). Diabetic retinopathy was self-reported (“Has a doctor ever told you that diabetes has affected your eyes or that you had retinopathy?”).

### Mortality Follow-up

Information on all-cause and cardiovascular mortality was obtained using the linkage of NHANES data to death certificate data from the National Death Index (24). The underlying cause of death was coded according to the ICD-10. Outcomes of interest included all-cause and cardiovascular (ICD-10 code I00–I78) mortality. Length of follow-up for each participant was calculated as the date of

the NHANES examination to date of death or 31 December 2006 (whichever occurred first).

### Statistical Analyses

All statistical analyses incorporated modified sampling weights, primary sampling units, and strata specific to the sample with available cystatin C in order to generate nationally representative estimates of the U.S. population (18). SEs were estimated using the Taylor series (linearization) method. Kidney function was analyzed both as a continuous measure using restricted cubic splines and as a categorical measure according to K/DOQI classification (eGFR <15, 15–29, 30–59, 60–89, and 90–200 mL/min/1.73 m<sup>2</sup>) for both eGFR<sub>cr</sub> and eGFR<sub>cys</sub>. Individuals with eGFR values >200 mL/min/1.73 m<sup>2</sup> were reassigned a value of 200 mL/min/1.73 m<sup>2</sup> (two persons with eGFR<sub>cr</sub> >200 mL/min/1.73 m<sup>2</sup>).

Modified Poisson regression models were used to examine the relationship of eGFR with the prevalence of coronary artery disease, peripheral arterial disease, albuminuria, and retinopathy (25). Associations of eGFR and mortality were examined using multivariable Cox proportional hazards models. The proportional hazards assumption was tested using log-log plots by category of eGFR. All multivariable models were adjusted for age (years), female sex (yes, no), and race (non-Hispanic black, non-Hispanic white, Hispanic, other) as well as current smoking status (current, former, never), BMI (measured as weight in kilograms divided by the square of height in meters), hypercholesterolemia (yes, no), low HDL (yes, no), hypertension (yes, no), coronary artery disease (yes, no), and albuminuria (yes, no; only in analyses where albuminuria was not the outcome). Given the possible relationship between cystatin C and obesity, interactions between BMI, eGFR<sub>cys</sub>, and adverse outcomes were tested in multivariable models. All analyses were conducted using Stata, version 12 (Stata, College Station, TX).

## RESULTS

### Study Population

The prevalence of diabetes among U.S. adults was 8.2% (95% CI 7.4–9.0).

Persons with diabetes were older, more likely to be black, and more likely to be male than those without diabetes (Table 1). Persons with diabetes also had poorer lipid profiles. Two-thirds of persons with diabetes had hypertension, which was nearly twice as prevalent compared with those without diabetes (63.8% vs. 33.4%). There was also a substantial difference in the distribution of BMI: 50% of the U.S. population with diabetes had a BMI >30 kg/m<sup>2</sup> compared with 28% without diabetes, and 28% of persons with diabetes had a BMI ≥35 kg/m<sup>2</sup> compared with 12% of those without diabetes. The prevalence of reduced kidney function was almost three times higher in persons with diabetes compared with those without diabetes (eGFR<sub>cr</sub> 16.5% vs. 5.8%, eGFR<sub>cys</sub> 22.0% vs. 7.9%).

### Prevalence of Reduced Kidney Function in Persons With Diabetes by Filtration Marker

The trend of higher prevalence of reduced kidney function by eGFR<sub>cys</sub>

persisted across subgroups of sex, race, age, and BMI (Supplementary Fig. 1). On the absolute scale, the discrepancy in eGFR<sub>cys</sub> versus eGFR<sub>cr</sub> was largest in older individuals. The absolute difference in reduced kidney function prevalence estimated by cystatin C versus creatinine was 6.9% in those aged 60–80 years and 10.3% in those aged 80 years and older. Similarly, the absolute difference was larger among persons with BMI >30 kg/m<sup>2</sup> (reduced kidney function by eGFR<sub>cys</sub> vs. eGFR<sub>cr</sub>: 20.0% vs. 13.1%) than among those with BMI <30 kg/m<sup>2</sup> (reduced kidney function by eGFR<sub>cys</sub> vs. eGFR<sub>cr</sub>: 21.3% vs. 16.9%).

### Reclassification of Reduced Kidney Function by Filtration Marker

Discordance between eGFR<sub>cr</sub> and eGFR<sub>cys</sub> in the classification of reduced kidney function was 11.8% in persons with diabetes and 4.7% in persons without diabetes. In the population with diabetes, 10.4% of those classified as having preserved kidney function by eGFR<sub>cr</sub> (8.7% of the 83.6% with eGFR<sub>cr</sub>

**Table 1—Characteristics of U.S. adults aged ≥20 years by diabetes status: NHANES 1999–2002 cystatin C subsample (N = 4,457)**

	No diabetes	Diabetes	P
<i>n</i>	3,679	778	
Age (years)	45.2 (0.4)	58.2 (1.1)	<0.001
Male, %	47.9 (1.2)	55.8 (3.2)	0.03
Race/ethnicity, %			<0.001
Non-Hispanic white	73.0 (2.1)	62.1 (3.8)	
Non-Hispanic black	10.2 (1.2)	17.2 (3.1)	
Mexican American	7.0 (0.9)	7.1 (1.5)	
Other	9.8 (1.9)	13.6 (3.6)	
Self-reported diabetes or medication use, %	—	79.5 (2.5)	
Hemoglobin A <sub>1c</sub> (mean, %)	5.3 (0.02)	7.7 (0.1)	<0.001
Hypertension, %	33.4 (1.4)	63.8 (3.8)	<0.001
Systolic blood pressure (mmHg), mean (SE)	122.8 (0.6)	132.9 (1.2)	<0.001
Diastolic blood pressure (mmHg), mean (SE)	72.2 (0.4)	71.0 (1.0)	0.3
BMI (kg/m <sup>2</sup> ), mean	27.8 (0.2)	31.6 (0.6)	<0.001
<25, %	37.7 (1.5)	18.6 (3.2)	<0.001
25–<30, %	34.2 (1.5)	31.4 (2.9)	
30–35, %	15.8 (0.7)	22.1 (2.8)	
>35, %	12.3 (1.1)	27.9 (3.2)	
Total cholesterol (mg/dL), mean (SE)	202.2 (1.2)	202.6 (2.8)	0.8
HDL (mg/dL), mean (SE)	51.5 (0.4)	45.7 (0.9)	<0.001
ACR (mg/g), mean (SE)	23.4 (3.0)	185.7 (30.3)	<0.001
eGFR <sub>cr</sub> (mL/min/1.73 m <sup>2</sup> ), mean (SE)	95.1 (0.6)	85.5 (1.7)	<0.001
eGFR <sub>cys</sub> (mL/min/1.73 m <sup>2</sup> ), mean (SE)	100.7 (0.8)	84.3 (2.6)	<0.001
eGFR <sub>cr</sub> <60 mL/min/1.73 m <sup>2</sup> , %	5.8 (0.4)	16.5 (1.4)	<0.001
eGFR <sub>cys</sub> <60 mL/min/1.73 m <sup>2</sup> , %	7.9 (0.6)	22.0 (2.1)	<0.001

$\geq 60$  mL/min/1.73 m<sup>2</sup>) were reclassified as having reduced kidney function by eGFR<sub>cys</sub> (Supplementary Table 1). In the population without diabetes, 3.6% of those classified as having preserved kidney function by eGFR<sub>cr</sub> (3.4% of the 94.2% with eGFR<sub>cr</sub>  $\geq 60$  mL/min/1.73 m<sup>2</sup>) were reclassified as having reduced kidney function by eGFR<sub>cys</sub> (Supplementary Table 1). Within the overall population, diabetes status was significantly associated with reclassification from preserved kidney function by eGFR<sub>cr</sub> to reduced kidney function by eGFR<sub>cys</sub> (odds ratio [OR] 3.1 [95% CI 1.9–4.9],  $P < 0.001$ ) (Supplementary Table 2). This association was attenuated but still significant after sequential adjustment for eGFR<sub>cr</sub> and BMI but not after further adjustment for age. In contrast, eGFR<sub>cr</sub>, BMI, age, and albuminuria were all significantly associated with reclassification by eGFR<sub>cys</sub> in multivariable regression (eGFR<sub>cr</sub>, OR 0.9 per 1 mL/min/1.73 m<sup>2</sup> increase,  $P < 0.001$ ; BMI, OR 1.5 per 5 kg/m<sup>2</sup> increase,  $P < 0.001$ ; age, OR 1.8 per decade increase,  $P < 0.001$ ; ACR  $> 30$  mg/g, OR 2.2,  $P = 0.01$ ).

#### Association of Kidney Function With Micro- and Macrovascular

#### Complications by Filtration Marker

In the population with diabetes, the prevalence of albuminuria, retinopathy, coronary artery disease, and peripheral arterial disease was high, even among those with preserved kidney function (Fig. 1). The most common of these complications was albuminuria, present in 27.0% of those with eGFR<sub>cr</sub>  $\geq 60$  mL/min/1.73 m<sup>2</sup> and 55.9% of those with eGFR<sub>cr</sub>  $< 60$  mL/min/1.73 m<sup>2</sup>. The prevalence of coronary artery disease, retinopathy, and peripheral artery disease was also two to three times higher among those with reduced kidney function than those with preserved kidney function. In general, the probability of microvascular and macrovascular complications increased with lower eGFR; above eGFR 90 mL/min/1.73 m<sup>2</sup>, relationships between eGFR and vascular complications were more variable. The unadjusted relationships between eGFR, coronary artery disease, peripheral arterial disease, albuminuria, and retinopathy

were similar using eGFR<sub>cr</sub> and eGFR<sub>cys</sub>, although there was suggestion of a “U-shape” (with higher risk at both higher and lower levels of GFR) in the association between albuminuria, retinopathy, and eGFR<sub>cr</sub> but not eGFR<sub>cys</sub> (Supplementary Fig. 2A–D).

After adjustment for demographic and traditional cardiovascular risk factors, the prevalence ratios for vascular complications (coronary artery disease, peripheral arterial disease, albuminuria, retinopathy) by eGFR category were similar using creatinine or cystatin C (Table 2). Compared with a reference group of eGFR 60–90 mL/min/1.73 m<sup>2</sup>, persons with advanced CKD (eGFR 15–30 mL/min/1.73 m<sup>2</sup>) had a higher prevalence of coronary artery disease, albuminuria, and retinopathy, regardless of filtration marker used. The adjusted relationships between eGFR category and peripheral arterial disease were similar in magnitude but not statistically significant. There were no significant interactions between BMI, eGFR<sub>cys</sub>, and vascular complications.

#### All-Cause and Cardiovascular Mortality in Those With and Without Reduced Kidney Function

There were 153 deaths (63 from a cardiovascular cause) during a median follow-up of 5.3 years among the 778 participants with diabetes. The risk of both all-cause and cardiovascular mortality increased with lower eGFR, regardless of filtration marker used (Table 3). Compared with the reference group of eGFR 60–90 mL/min/1.73 m<sup>2</sup>, persons with eGFR 15–30 mL/min/1.73 m<sup>2</sup> had a significantly higher risk of all-cause mortality (eGFR<sub>cys</sub>, hazard ratio [HR] 3.8,  $P = 0.007$ ; eGFR<sub>cr</sub>, HR 2.6,  $P = 0.04$ ). By contrast, eGFR 15–30 mL/min/1.73 m<sup>2</sup> was significantly associated with increased cardiovascular mortality when estimated by cystatin C (HR 5.3,  $P = 0.007$ ) but not by creatinine (HR 2.0,  $P = 0.3$ ). There were no significant interactions between BMI, eGFR<sub>cys</sub>, and either all-cause or cardiovascular mortality.


#### CONCLUSIONS

This nationally representative study of persons with diabetes suggests that the use of cystatin C to estimate kidney function would result in a higher

prevalence of reduced kidney function than would estimates using serum creatinine. Reclassification from preserved kidney function using creatinine to reduced kidney function using cystatin C occurred more commonly among persons with diabetes than those without, but this observation was explained by differences in the distributions of eGFR, BMI, age, and albuminuria between the two populations: reclassification was significantly associated with lower eGFR<sub>cr</sub>, higher BMI, older age, and ACR  $> 30$  mg/g. Lower eGFR as determined by either creatinine or cystatin was associated with higher odds of prevalent vascular complications; however, the shape of the relationship with albuminuria and retinopathy at higher levels of eGFR differed slightly by filtration marker in unadjusted analysis. Similarly, while low eGFR was robustly associated with all-cause mortality, only eGFR<sub>cys</sub> showed significant association with cardiovascular mortality.

Differences in eGFR<sub>cr</sub> and eGFR<sub>cys</sub> have been noted previously in the general population (26,27). In the U.S. noninstitutionalized civilian population, kidney function estimated using cystatin C resulted in a reduced kidney function prevalence of 8.7% compared with an estimated 6.5% using creatinine (26). In a cross-sectional study of 1,360 inhabitants of the Alpine region in Europe, Pattaro et al. (28) noted that the Lin concordance correlation coefficient of eGFR<sub>cr</sub> and eGFR<sub>cys</sub> was 0.56, with significant differences by age (0.57 in those  $\geq 65$  years old vs. 0.38 in those  $< 65$  years old) but not by diabetes status. Our results differ somewhat from those of this prior study; the presence of diabetes differentially affected kidney disease classification by eGFR<sub>cr</sub> and eGFR<sub>cys</sub>, at least in univariable analysis—an observation that may be attributable to our larger sample size and distinct American population.

Neither creatinine nor cystatin C is a perfect marker of glomerular filtration; each has non-GFR determinants. Some have argued that neither marker can adequately estimate true GFR in persons with diabetes; however, this concern is primarily relevant for those



**Figure 1**—Prevalence of microvascular and macrovascular conditions among U.S. adults with diabetes according to the presence or absence of reduced kidney function (estimated using creatinine and cystatin C).

with high GFR—not reduced GFR as in the current study (29,30). The strong relationship between reclassification to reduced kidney function by eGFR<sub>cys</sub> and age may be due to inherent properties of the filtration markers. In the case of creatinine, muscle mass and diet are significantly associated with creatinine levels (31). Older persons may be sicker than their younger peers; thus, kidney

function estimated using serum creatinine may be confounded by cachexia and muscle wasting. A similar explanation could apply to the differences seen with albuminuria (i.e., those with albuminuria are sicker than their peers without albuminuria). To our knowledge, the observation that reclassification by cystatin C occurs more frequently among those with

albuminuria is novel; however, it is fully consistent with a recent study demonstrating that the decrement in eGFR<sub>cys</sub> associated with 24-h albuminuria >30 mg was greater than that of eGFR<sub>cr</sub> or measured GFR (27). In the population with diabetes, both serum creatinine and cystatin C may have drawbacks. Serum creatinine may poorly estimate kidney function given

**Table 2**—Adjusted prevalence ratios (95% CI) for categories of kidney function with complications of diabetes, by filtration marker, among U.S. adults with diabetes (N = 778)\*

	Coronary artery disease	Peripheral arterial disease**	Albuminuria†	Retinopathy
<b>eGFR<sub>cr</sub> (mL/min/1.73 m<sup>2</sup>)</b>				
>90	1.1 (0.7–1.6)	0.9 (0.4–1.9)	0.97 (0.6–1.7)	1.8 (0.9–3.4)
60–90	1.0 (ref.)	1.0 (ref.)	1.0 (ref.)	1.0 (ref.)
30–60	1.4 (1.0–1.9)	1.7 (0.9–3.5)	1.7 (1.2–2.3)	1.9 (1.2–3.0)
15–30	2.8 (1.4–5.6)	2.7 (0.8–8.8)	2.3 (1.4–3.6)	3.9 (2.0–7.8)
<b>eGFR<sub>cys</sub> (mL/min/1.73 m<sup>2</sup>)</b>				
>90	0.8 (0.4–1.5)	0.8 (0.3–2.1)	1.0 (0.6–1.7)	0.6 (0.3–1.4)
60–90	1.0 (ref.)	1.0 (ref.)	1.0 (ref.)	1.0 (ref.)
30–60	1.2 (0.8–1.6)	1.4 (0.7–2.8)	2.2 (1.5–3.2)	1.1 (0.6–1.9)
15–30	2.4 (1.4–4.1)	1.9 (0.5–7.3)	3.1 (2.0–4.8)	2.1 (1.2–3.9)

\*All analyses adjusted for age, sex, race, hypercholesterolemia, low HDL, smoking, hypertension, BMI, and albuminuria, †except model of albuminuria. Insufficient numbers in eGFR <15 mL/min/1.73 m<sup>2</sup> category to accurately estimate. \*\*Limited to adults aged ≥40 years with ABI measurements.

**Table 3—Adjusted hazard ratios (95% CI) for categories of kidney function with all-cause and cardiovascular mortality, by filtration marker, among U.S. adults with diabetes (N = 778)\***

	Number at risk	All-cause mortality (n = 153 events)	Cardiovascular mortality (n = 63 events)
<b>eGFR<sub>cr</sub> (mL/min/1.73 m<sup>2</sup>)</b>			
>90	243	0.7 (0.2–3.5)	0.1 (0.0–0.8)
60–90	336	1.0 (ref.)	1.0 (ref.)
30–60	162	1.8 (1.0–3.2)	1.8 (0.6–5.4)
15–30	23	2.6 (1.1–6.4)	2.0 (0.5–7.9)
<b>eGFR<sub>cys</sub> (mL/min/1.73 m<sup>2</sup>)</b>			
>90	241	0.5 (0.2–1.1)	0.7 (0.1–3.8)
60–90	289	1.0 (ref.)	1.0 (ref.)
30–60	191	1.5 (0.6–3.3)	1.3 (0.5–3.2)
15–30	39	3.8 (1.5–9.9)	5.3 (1.6–17.3)

\*All analyses adjusted for age, sex, race, smoking, hypertension, coronary artery disease, hypercholesterolemia, low HDL, albuminuria, and BMI. Insufficient numbers in eGFR <15 mL/min/1.73 m<sup>2</sup> category to accurately estimate.

the tendency of persons with diabetes to have a lower than average muscle mass. Cystatin C may be directly affected by both BMI and diabetes (13,32,33). In obese individuals, cystatin C levels are higher, and eGFR<sub>cys</sub> significantly underestimates true kidney function (27,31). Indeed, our study suggests that much of the association between diabetes and cystatin C is driven by differences in the distribution of age and BMI. Additional work is needed to determine whether an approach using cystatin C or both filtration markers (the latter of which better approximates measured GFR in the overall population) would improve kidney function estimates in the population with diabetes (16).

Conventional wisdom is that, among persons with diabetes, kidney function decline and vascular complications go hand in hand. Certainly, our results support the association of prevalent complications with very low eGFR whether estimated by creatinine or cystatin C. Interestingly, in the upper ranges of preserved kidney function, the association between eGFR<sub>cr</sub> and retinopathy reversed, with higher levels of eGFR<sub>cr</sub> conferring increased odds of retinopathy, although this was not statistically significant in adjusted analysis. A similar pattern was seen in the univariate association of eGFR<sub>cr</sub> and albuminuria. These observations are consistent with previous studies demonstrating weaker-than-expected correlation between eGFR<sub>cr</sub>,

albuminuria, and retinopathy (34). Because of the more monotonic relationship seen between eGFR<sub>cys</sub>, albuminuria, and retinopathy, it is possible that kidney function based on cystatin C may prove a better predictor of diabetes complications than that based on creatinine. However, this may be more useful in defining a low-risk group than a high-risk group, given the larger differences in the upper ranges of eGFR. We also observed that the relationship between eGFR<sub>cys</sub> and all-cause and cardiovascular mortality was stronger than the corresponding relationship with eGFR<sub>cr</sub>, similar to findings in the general population (11,12,35) and previous studies of persons with diabetes (36). Additional prospective studies are needed to determine whether eGFR<sub>cys</sub> provides better risk stratification for subsequent diabetes complications.

Our study has certain limitations. We relied on a single measurement of creatinine and cystatin; GFR is estimated from these filtration markers and not measured directly. As such, we cannot assess which filtration marker most closely approximates true kidney function. Additional research is needed to determine whether the confounding by age and BMI (or other unmeasured confounders such as thyroid disease) in diabetes favors eGFR<sub>cr</sub>, eGFR<sub>cys</sub>, or perhaps a combination of the two. Additionally, a single random sample of urine was used to quantify albuminuria. Next, despite being

nationally representative, the subset of NHANES with diabetes was relatively small, with a short duration of follow-up for mortality outcomes. Some of the vascular complications were self-reported, and insofar as reporting may vary by level of eGFR, this may lead to bias. Finally, persons with more severe kidney disease are likely underrepresented in NHANES, thus limiting accuracy in the very low ranges of GFR.

In summary, this study demonstrates that using cystatin C to classify kidney function among persons with diabetes results in a higher prevalence of reduced kidney function yet the same or stronger associations with vascular complications and mortality. Future studies are needed to determine whether incorporating cystatin C measurement into clinical care and kidney function estimation would improve outcomes in persons with diabetes.

**Funding.** This study was supported by the National Kidney Foundation (grant to E.S. at Johns Hopkins) and Merck Sharp and Dohme Corp., a subsidiary of Merck & Co, Inc., Whitehouse Station, NJ (professional service agreement to the National Kidney Foundation). M.E.G. receives support from the National Institute of Diabetes and Digestive and Kidney Diseases (K08-DK-092287).

**Duality of Interest.** No potential conflicts of interest relevant to this article were reported.

**Author Contributions.** C.-W.T. and M.E.G. researched data and wrote the manuscript. L.A.I. and J.C. reviewed and edited the manuscript. E.S. contributed to the discussion and reviewed and edited the manuscript. M.E.G. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## References

1. National Kidney Foundation. KDOQI clinical practice guideline for diabetes and CKD: 2012 update. *Am J Kidney Dis* 2012;60:850–886
2. U.S. Renal Data System. *USRDS 2012 Annual Data Report: Atlas of End-Stage Renal Disease in the United States*. Bethesda, MD, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, 2012
3. de Boer IH, Rue TC, Hall YN, Heagerty PJ, Weiss NS, Himmelfarb J. Temporal trends in the prevalence of diabetic kidney disease in the United States. *JAMA* 2011;305:2532–2539

4. Cowie CC, Rust KF, Byrd-Holt DD, et al. Prevalence of diabetes and high risk for diabetes using A1C criteria in the U.S. population in 1988-2006. *Diabetes Care* 2010;33:562-568
5. Goff DC Jr, Gerstein HC, Ginsberg HN, et al.; ACCORD Study Group. Prevention of cardiovascular disease in persons with type 2 diabetes mellitus: current knowledge and rationale for the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Am J Cardiol* 2007;99:4i-20i
6. Tandon N, Ali MK, Narayan KM. Pharmacologic prevention of microvascular and macrovascular complications in diabetes mellitus: implications of the results of recent clinical trials in type 2 diabetes. *Am J Cardiovasc Drugs* 2012;12:7-22
7. Matsushita K, van der Velde M, Astor BC, et al.; Chronic Kidney Disease Prognosis Consortium. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet* 2010;375:2073-2081
8. Fox CS, Matsushita K, Woodward M, et al.; Chronic Kidney Disease Prognosis Consortium. Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without diabetes: a meta-analysis. *Lancet* 2012;380:1662-1673
9. Ferguson MA, Waikar SS. Established and emerging markers of kidney function. *Clin Chem* 2012;58:680-689
10. Stevens LA, Coresh J, Schmid CH, et al. Estimating GFR using serum cystatin C alone and in combination with serum creatinine: a pooled analysis of 3,418 individuals with CKD. *Am J Kidney Dis* 2008;51:395-406
11. Astor BC, Levey AS, Stevens LA, Van Lente F, Selvin E, Coresh J. Method of glomerular filtration rate estimation affects prediction of mortality risk. *J Am Soc Nephrol* 2009;20:2214-2222
12. Shlipak MG, Wassel Fyr CL, Chertow GM, et al. Cystatin C and mortality risk in the elderly: the health, aging, and body composition study. *J Am Soc Nephrol* 2006;17:254-261
13. Stevens LA, Schmid CH, Greene T, et al. Factors other than glomerular filtration rate affect serum cystatin C levels. *Kidney Int* 2009;75:652-660
14. Iliadis F, Didangelos T, Ntemka A, et al. Glomerular filtration rate estimation in patients with type 2 diabetes: creatinine- or cystatin C-based equations? *Diabetologia* 2011;54:2987-2994
15. Silveiro SP, Araújo GN, Ferreira MN, Souza FD, Yamaguchi HM, Camargo EG. Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation pronouncedly underestimates glomerular filtration rate in type 2 diabetes. *Diabetes Care* 2011;34:2353-2355
16. Inker LA, Schmid CH, Tighiouart H, et al.; CKD-EPI Investigators. Estimating glomerular filtration rate from serum creatinine and cystatin C. *N Engl J Med* 2012;367:20-29
17. Levey AS, Stevens LA, Schmid CH, et al.; CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604-612
18. Köttgen A, Selvin E, Stevens LA, Levey AS, Van Lente F, Coresh J. Serum cystatin C in the united states: The third national health and nutrition examination survey (NHANES III). *Am J Kidney Dis* 2008;51:385-394
19. Centers for Disease Control and Prevention (CDC). Survey operations manuals, brochures, and consent documents: 1999-2000 NHANES. Lower extremity disease procedures manual [Internet], 2001. Available from <http://www.cdc.gov/nchs/data/nhanes/le.pdf>. Accessed 28 April 2013
20. Feringa HH, Bax JJ, van Waning VH, et al. The long-term prognostic value of the resting and postexercise ankle-brachial index. *Arch Intern Med* 2006;166:529-535
21. Selvin E, Manzi J, Stevens LA, et al. Calibration of serum creatinine in the national health and nutrition examination surveys (NHANES) 1988-1994, 1999-2004. *Am J Kidney Dis* 2007;50:918-926
22. Selvin E, Juraschek SP, Eckfeldt J, Levey AS, Inker LA, Coresh J. Calibration of cystatin C in the National Health and Nutrition Examination Surveys (NHANES). *Am J Kidney Dis* 2013;62:253-260
23. Newman AB, Siscovick DS, Manolio TA, et al.; Cardiovascular Heart Study (CHS) Collaborative Research Group. Ankle-arm index as a marker of atherosclerosis in the Cardiovascular Health Study. *Circulation* 1993;88:837-845
24. NHANES (1999-2004) public use linked mortality files [article online]. Available from [http://www.cdc.gov/nchs/data\\_access/data\\_linkage/mortality/nhanes\\_99\\_04\\_linkage.htm#description](http://www.cdc.gov/nchs/data_access/data_linkage/mortality/nhanes_99_04_linkage.htm#description). Accessed 31 July 2013
25. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702-706
26. Grams ME, Juraschek SP, Selvin E, et al. Trends in the prevalence of reduced GFR in the United States: a comparison of creatinine- and cystatin C-based estimates. *Am J Kidney Dis* 2013;62:253-260
27. Rule AD, Bailey KR, Lieske JC, Peysers PA, Turner ST. Estimating the glomerular filtration rate from serum creatinine is better than from cystatin C for evaluating risk factors associated with chronic kidney disease. *Kidney Int* 2013;83:1169-1176
28. Pattaro C, Riegler P, Stifter G, Modenese M, Minelli C, Pramstaller PP. Estimating the glomerular filtration rate in the general population using different equations: effects on classification and association. *Nephron Clin Pract* 2013;123:102-111
29. Gaspari F, Ruggenti P, Porrini E, et al.; GFR Study Investigators. The GFR and GFR decline cannot be accurately estimated in type 2 diabetics. *Kidney Int* 2013;84:164-173
30. Rossing P, Rossing K, Gaede P, Pedersen O, Parving HH. Monitoring kidney function in type 2 diabetic patients with incipient and overt diabetic nephropathy. *Diabetes Care* 2006;29:1024-1030
31. Chew-Harris JS, Florkowski CM, George PM, Elmslie JL, Endre ZH. The relative effects of fat versus muscle mass on cystatin C and estimates of renal function in healthy young men. *Ann Clin Biochem* 2013;50:39-46
32. Knight EL, Verhave JC, Spiegelman D, et al. Factors influencing serum cystatin C levels other than renal function and the impact on renal function measurement. *Kidney Int* 2004;65:1416-1421
33. Macdonald J, Marcora S, Jibani M, et al. GFR estimation using cystatin C is not independent of body composition. *Am J Kidney Dis* 2006;48:712-719
34. Wolf G, Müller N, Mandecka A, Müller UA. Association of diabetic retinopathy and renal function in patients with types 1 and 2 diabetes mellitus. *Clin Nephrol* 2007;68:81-86
35. Menon V, Shlipak MG, Wang X, et al. Cystatin C as a risk factor for outcomes in chronic kidney disease. *Ann Intern Med* 2007;147:19-27
36. de Boer IH, Katz R, Cao JJ, et al. Cystatin C, albuminuria, and mortality among older adults with diabetes. *Diabetes Care* 2009;32:1833-1838