

Carotenoid Intake from Natural Sources and Head and Neck Cancer: A Systematic Review and Meta-analysis of Epidemiological Studies

Emanuele Leoncini¹, Darko Nedovic¹, Nikola Panic¹, Roberta Pastorino¹, Valeria Edefonti², and Stefania Boccia¹

Abstract

Because of their role as antioxidants, the intake of carotenoids has been hypothesized to reduce the risk of head and neck cancer (HNC). We conducted a systematic review and meta-analysis of the epidemiological studies to investigate whether the intake of specific carotenoids from dietary sources, as well as combined carotenoids, is associated with the risk of HNC according to cancer subsites. A comprehensive literature search of the Medline and Scopus databases was conducted. Sixteen articles were identified from the literature search, of which 15 were case-control studies and one prospective cohort study. The risk reduction associated with β -carotene equivalents intake was 46% (95% CI, 20%–63%)

for cancer of oral cavity and 57% (95% CI, 23%–76%) for laryngeal cancer. Lycopene and β -cryptoxanthin also reduced the risk for laryngeal cancer; the ORs for the highest category compared with the lowest one of carotenoid intake were 50% (95% CI, 11%–72%) and 59% (95% CI, 49%–67%), respectively. Lycopene, α -carotene, and β -cryptoxanthin were associated with at least 26% reduction in the rate of oral and pharyngeal cancer (95% CI, 2%–44%). Our systematic review and meta-analysis on dietary carotenoids intake and HNC showed carotenoids to act protectively against HNC, in relation to most of single nutrients and subsites. *Cancer Epidemiol Biomarkers Prev*; 24(7); 1003–11. ©2015 AACR.

Introduction

Head and neck cancer (HNC) encompasses tumors that occur in several sites of the head and neck region, including the oral cavity, pharynx, and larynx, with 90% being squamous cell carcinomas (1). Worldwide, cancers of the head and neck are the sixth most common cancer, with an estimated half million new cases and over 450,000 deaths in 2012 (2). Carotenoids represent a diverse group of natural pigments of the polyene type (3) and are present in non-starchy vegetables and fruit. Carotenoids can be classified into oxygenated carotenoids (β -cryptoxanthin, lutein, and zeaxanthin), called xanthophylls, and hydrocarbon carotenoids (e.g., α -carotene, β -carotene, and lycopene), called carotenes (4). They are known to be very efficient physical and chemical quenchers of singlet oxygen, and potent scavengers of other reactive oxygen species (5, 6). Because of their role, the intake of carotenoids has been hypothesized to reduce the risk of HNC (7). However, carotenoids from natural sources might have a different effect on HNC, depending on the degree of their biologic activity and antioxidant capacities (8).

Several epidemiological studies have examined the association between carotenoid intake from dietary sources and HNC risk (9–18); however, only a few assessed the association

between single-nutrient intake and HNC, with conflicting results (17, 19–24).

Although some reviews report that carotenoids intake have a protective effect on overall HNC (25, 26), the effect on specific cancer subsites and that of single carotenoids so far were not addressed. The aim of this study is to conduct a systematic review and meta-analysis of the epidemiological studies to investigate whether the intake of specific carotenoids from dietary sources, as well as combined carotenoids, reduces the risk of HNC according to specific cancer subsites.

Materials and Methods

Search strategy

We conducted a comprehensive literature search of the MEDLINE and Scopus databases. We selected the following keywords for the literature search: "diet", "nutrients", "carotenoids", "cryptoxanthin", "lycopene", "lutein", "zeaxanthin", "oral carcinoma", "oral cancer", "pharyngeal cancer", "laryngeal cancer", "hypopharyngeal cancer", "oropharyngeal cancer", and "head and neck cancer". The search was limited to human subjects with language restriction to English studies until May 1, 2014. The snowball strategy, including manual search of the references listed by studies retrieved from the online databases, was also adopted to identify additional studies. Abstracts, systematic reviews, editorials, and case reports were not included.

Inclusion and exclusion criteria

The eligibility criteria for inclusion in the review required that the study: (i) refers to one or more conditions corresponding with HNC clinical entity; (ii) examines the association of HNC risk with carotenoid intakes from natural sources either when considered separately or when combined; (iii) provides effect

¹Section of Hygiene, Institute of Public Health, Università Cattolica del Sacro Cuore, Rome, Italy. ²Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy.

Corresponding Author: Stefania Boccia, Università Cattolica del Sacro Cuore, Largo F. Vito 1, 00168 Rome, Italy. Phone: 39-06-30154396/35001527; Fax: 39-06-35001522; E-mail: sboccia@rm.unicatt.it

doi: 10.1158/1055-9965.EPI-15-0053

©2015 American Association for Cancer Research.

measures, such as a risk ratio (RR) or odds ratio (OR), and associated 95% confidence interval (CI). We excluded comments, letters, ecological studies, and animal studies. Studies were also included if they were case-control or cohort studies, studied the effects of levels of carotenoid intakes, and reported HNC cancer as the outcomes of interest. For articles including the same population resources or overlapping datasets, the study with the largest number of cases was included.

Data extraction

Two investigators (N. Panic and E. Leoncini) independently extracted information from the included studies and entered into an Excel 2010 (Microsoft Corp.) spreadsheet. Any discrepancies regarding individual study inclusion, data extraction, and interpretation were resolved by consulting a third investigator (D. Nedovic). We extracted data related to study characteristics, including first author name, year of publication, location of the study, study period, study design, number of cases and controls, site of cancer, and carotenoid examined.

Exposure to different carotenoids was investigated by extracting information on both combined carotenoids and on the intakes of individual carotenoids: α -carotene, β -carotene, β -cryptoxanthin, lycopene, lutein, and zeaxanthin.

When the general term "carotene" is used in the article, we contacted the corresponding author to ask what is meant by carotene. The term carotene has been used to indicate β -carotene equivalents in four articles (17, 18, 23, 27). We were not able to track down an updated email address for two authors (12, 15). We hypothesized that the term carotene has also been used to indicate β -carotene equivalents.

As the level of carotenoids was often categorized into quantiles, the highest intake group for HNC compared with the lowest intake group was reported in each study.

Quality assessment

We performed the quality assessment of the included studies by using Newcastle-Ottawa quality assessment scale. This scale varies from 0 (lowest quality score) to a maximum possible score of 9 (highest quality score) and incorporates information on participant selection, outcome and exposure ascertainment, and the potential for confounding (28). The systematic review and meta-analysis was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses Statement (29).

Statistical analysis

The association between carotenoid intake and HNC risk overall and by cancer subsites was estimated by calculating pooled OR and 95% CI, comparing the highest with the lowest category of carotenoid intake. We used random-effects models to account for variation between studies as this can provide more conservative results than a fixed-effects model (30). We quantified the proportion of the total variation due to that heterogeneity by using the I^2 statistic (31). Analyses were conducted using Stata software (StataCorp. 2013. Stata Statistical Software: Release 13; StataCorp LP).

Results

Figure 1 reports the flowchart of the bibliographic search strategy and the results. We identified a total of 432 articles in




Figure 1. The search strategy and flow diagram for database search.

our initial search. Of these, 395 articles were excluded as duplicates, reviews, or unrelated to the research topic. The remaining 37 articles were assessed for eligibility, and 13 articles were excluded because they did not meet the inclusion criteria. An additional 8 articles were excluded as they did not report a measure of association between carotenoid intakes and HNC risk (9–11, 13, 14, 16, 19, 32). Finally, 16 articles were included, of which 15 were case-control studies (12, 15, 17, 18, 20–24, 27, 33–37) and one prospective cohort study (38).

Table 1 shows the main characteristics of the 16 studies included. Case-control studies were published between 1991 and 2013 and involved a total of 5,482 cases and 14,130 controls. The prospective cohort study included 34,691 postmenopausal Iowa women over the period 1986 to 1992. Eight studies were conducted in Europe (17, 22–24, 27, 33, 34, 36), four in North America (12, 35, 37, 39), two in South America (20, 21), and two in Asia (15, 18). Most of the studies included provided effect measures estimates adjusted at least for age, gender, energy intake, and smoking (12, 15, 17, 18, 20, 21, 23, 24).

Regarding the methodological quality of the included studies, six (37.5%) were scored 8 (12, 15, 20, 21, 35, 37) according to the Newcastle-Ottawa scale, five (31.3%) were scored 7 (22, 23, 27, 33, 36), three (18.7%) were scored 6 (17, 18, 34), and two (12.5%) were scored 5 (25, 38).

Table 1. Characteristics of the studies of carotenoid intake and head and neck cancer (HNC) risk included in the final analysis ($n = 16$)

| First author, year (ref.) | Study location | Year | Population (cases/controls) | OR (95% CI) for the highest vs. the lowest category of consumption by site of cancer | Adjusted for | Newcastle-Ottawa score |
|---------------------------|--------------------|-----------|---------------------------------|--|--|------------------------|
| Bravi, 2013 (24) | Italy, Switzerland | 1997–2009 | 768 OCP/2,078 | α -Carotene 0.51 (0.34–0.76), β -carotene 0.28 (0.18–0.43), β -cryptoxanthin 0.37 (0.24–0.56), lycopene 0.92 (0.54–1.55), lutein plus zeaxanthin 0.34 (0.23–0.51) | Age, gender, center, education, year of interview, BMI, tobacco smoking, alcohol drinking, nonalcohol energy intake | 5 |
| Suzuki, 2006 (18) | Japan | 2001–2004 | 385 (193 OC; 132 P, 60 L)/1,925 | HNC: 0.59 (0.42–0.84); OC: carotene 0.56 (0.35–0.91); P: carotene 0.61 (0.31–1.20); L: carotene 0.51 (0.21–1.25) | Smoking status, drinking status, multivitamin use, nonalcohol energy intake, dental brushing, year of first visit (matched for age and gender) | 6 |
| Gallus, 2003 (27) | Italy, Switzerland | 1986–2000 | 68 L/340 | Carotene 0.74 (0.32–1.74), lycopene 0.5 (0.10–2.10) | Education, BMI, tobacco and alcohol consumption, nonalcohol energy intake (conditioned on age, year of interview, study center) | 7 |
| Bidoli, 2003 (23) | Italy, Switzerland | 1992–2000 | 527 L/1,297 | Carotene 0.30 (0.20–0.40), α -carotene 0.30 (0.20–0.50), β -carotene 0.20 (0.20–0.40), β -cryptoxanthin 0.40 (0.20–0.50), lycopene 0.70 (0.50–1.10), lutein plus zeaxanthin 0.40 (0.30–0.60) | Education, BMI, alcohol drinking, smoking habits, nonalcohol energy (conditioned on age, gender, center) | 7 |
| Petridou, 2002 (22) | Greece | nr | 106 OC/106 | β -Carotene 1.01 (0.67–1.51) | BMI, height, education, oral hygiene, tobacco and alcohol consumption, coffee consumption, total energy intake | 7 |
| Negri, 2000 (17) | Italy | 1992–1997 | 754 OCP/1,775 | Carotene 0.43 (0.28–0.66), lycopene 0.67 (0.44–1.02) | Age, gender, study center, education, occupation, BMI, tobacco smoking, alcohol drinking, nonalcohol energy | 6 |
| De Stefani, 2000 (27) | Uruguay | 1996–1998 | 142 (66 OCP, 76 L)/491 | OCP: lycopene 0.42 (0.19–0.91); L: lycopene 0.21 (0.10–0.43) | Age, gender, residence, urban/rural status, education, BMI, tobacco smoking, alcohol drinking, total energy intake | 8 |
| De Stefani, 1999 (20) | Uruguay | 1996–1997 | 67 (33 OCP, 34 L)/393 | OCP: α -carotene 0.70 (0.40–1.20), β -carotene 1.20 (0.70–2.10), lycopene 0.90 (0.50–1.50), lutein 0.80 (0.40–1.40), cryptoxanthin 0.60 (0.30–1.00); L: α -carotene 0.70 (0.40–1.10), β -carotene 1.50 (0.90–2.40), lycopene 0.70 (0.50–1.20), lutein 0.90 (0.50–1.40), cryptoxanthin 0.50 (0.30–0.90) | Age, gender, residence, urban/rural status, education, BMI, tobacco smoking (pack-years), total alcohol intake, total energy intake | 8 |
| Schantz, 1997 (37) | USA | 1992–1994 | 167 HNC/177 | α -Carotene 1.30 (0.67–2.55), β -carotene 1.39 (0.73–2.67), β -cryptoxanthin 0.30 (0.15–0.60), lycopene 0.60 (0.32–1.11), lutein 0.95 (0.52–1.73) | Age, gender, ethnicity, caloric intake | 8 |

(Continued on the following page)

Table 1. Characteristics of the studies of carotenoid intake and head and neck cancer (HNC) risk included in the final analysis ($n = 16$). (Cont'd)

| First author, year (ref.) | Study location | Year | Population (cases/controls) | OR (95% CI) for the highest vs. the lowest category of consumption by site of cancer | Adjusted for | Newcastle-Ottawa score |
|-------------------------------|-----------------------------------|-----------|-------------------------------|--|---|------------------------|
| Esteve, 1996 (36) | Spain, Italy, Switzerland, France | 1977-1983 | 1147 (727 L, 399 HE)/2,736 | L: β -carotene 1.66 (1.13-2.43) ^a ; HE: β -carotene 1.31 (0.81-2.10) ^a | Age, study center, energy intake, tobacco, alcohol | 7 |
| Zheng, 1995 ^b (38) | USA | 1986-1992 | 34,691 subjects, 33 MPE cases | MPE: carotene 0.70 (0.30-1.80) ^c | Age, smoking status, pack-years of cigarettes, total energy intake | 5 |
| Zheng, 1993 (15) | China | 1989 | 404 OC/404 | Carotene 0.51 (0.27-0.96) | Age, gender, education, inadequate dentition, tobacco smoking, alcohol drinking, Quetelet index, total energy intake | 8 |
| Marshall, 1992 (12) | USA | 1975-1983 | 290 OCP/290 | Carotene 1.00 (0.80-1.20) | Age, gender, total calories, Quetelet index, smoking, alcohol risk imposed by teeth lost but not replaced (matched for age, gender, social class) | 8 |
| Freudenheim, 1992 (35) | USA | 1975-1985 | 250 L/250 | Carotenoids 0.40 (0.20-0.83) | Kilocalories, cigarettes, alcohol, education | 8 |
| La Vecchia, 1991 (34) | Italy | 1987-1989 | 105 OCP/1,169 | Carotenoids 0.30 (0.20-0.40) | Age, gender, area of residence, education, social class, smoking | 6 |
| Franceschi, 1991 (33) | Italy | 1985-1991 | 302 OCP/699 | Carotenoids 0.80 (0.60-1.20) | Age, gender, occupation, smoking and drinking habits | 7 |

Abbreviations: BMI, body mass index; E, endolarynx; HE, hypopharynx and epilarynx; L, larynx; MPE, mouth, pharynx, and esophagus; na, not applicable; nr, not reported; OC, oral cavity; OCP, oral cavity and pharynx; P, pharynx.


^aThe reference group is the highest category.

^bProspective cohort study.

^cRelative risk was reported.

Figure 2.

Meta-analysis of studies reporting HNC risk in relation to natural intake of combined carotenoids and β -carotene equivalents. Quantile cutoff points (or average intake): Franceschi, 1991: not reported; La Vecchia, 1991: not reported; Freudenheim, 1992: $\geq 8,126$ vs. $\leq 3,119$ IU/day; Negri, 2000: mean (SD) = 4,067.5 (2,398.1) $\mu\text{g}/\text{day}$ among controls; Marshall, 1992: not reported; Suzuki, 2006: $>4,029$ vs. $<1,906$ $\mu\text{g}/\text{day}$; Zheng, 1993: $\geq 3,486$ vs. $<2,054$ IU/day; Bidoli, 2003: mean (SD) = 3,700 (2,400) $\mu\text{g}/\text{day}$ among controls; Gallus, 2003: not reported.



Association between carotenoid intake and HNC

Combined carotenoids. Three studies addressed the association between the intake of combined carotenoids in general and HNC (33–35). We reported a nonsignificant inverse association for the highest level of intake of carotenoids for cancer of oral cavity and pharynx (OR, 0.48; 95% CI, 0.19–1.27; $I^2 = 93.3\%$; $P < 0.001$) (Fig. 2). Only a case–control study from the United States addressed the association between combined carotenoids intake and larynx, and observed a consistent protective effect (OR, 0.40; 95% CI, 0.19–0.83; ref. 35).

β -Carotene equivalents. Six case–control studies investigated the association between β -carotene equivalents intake and HNC (12, 15, 17, 18, 23, 27). The meta-analysis identified β -carotene equivalents to act protectively against oral cavity cancer (OR, 0.54; 95% CI, 0.37–0.80; $I^2 = 0.0\%$; $P = 0.82$) and laryngeal cancer (OR, 0.43; 95% CI, 0.24–0.77; $I^2 = 57.4\%$; $P = 0.10$; Fig. 2). In a case–control study conducted in Japan, a nonsignificant association was observed between β -carotene equivalents intakes and pharynx (OR, 0.61; 95% CI, 0.31–1.20). When oral cancer and pharynx cancer were considered together, an inverse nonsignificant association was observed (OR, 0.67; 95% CI, 0.29–1.53; $I^2 = 92.1\%$; $P < 0.001$).


Moreover, one prospective cohort study of 34,691 postmenopausal women in Iowa examined the association between β -carotene equivalents intake and the risk of cancers of the upper digestive tract (mouth, pharynx, and esophagus), with no significant association found (RR, 0.70; 95% CI, 0.30–1.80; ref. 38).

α -Carotene. Three studies examined the association between α -carotene intake and subsites of HNC (20, 23, 24). A strong inverse association was found for a high intake of α -carotene for cancer of the oral cavity and pharynx (OR, 0.57; 95% CI, 0.41–0.79; $I^2 = 0.0\%$; $P = 0.35$), whereas a nonsignificant inverse association was reported for cancer of the larynx (OR, 0.46; 95% CI, 0.20–1.06; $I^2 = 83.1\%$; $P < 0.05$; Fig. 3). Only one case–control study from the United States examined the association between α -carotene intake and the risk of HNC in general, with no significant association found (OR, 1.30; 95% CI, 0.66–0.255; ref. 37).

β -Carotene. Six studies reported data on β -carotene intake and HNC (20, 22–24, 36, 37). The meta-analysis showed a nonsignificant inverse association for a high intake of β -carotene, both for cancer of the oral cavity and pharynx (OR, 0.57; 95% CI, 0.14–2.38, $I^2 = 93.9\%$, $P < 0.001$) and for that of the larynx (OR, 0.58; 95% CI, 0.22–1.55; $I^2 = 91.4\%$; $P < 0.001$; Fig. 3).

Results from studies on β -carotene intake in relation to risk of HNC in general (OR, 1.39; 95% CI, 0.72–2.67; ref. 37), epilararyngeal and hypopharyngeal cancer (OR, 0.76; 95% CI, 0.47–1.23; ref. 36), and oral cancer (OR, 1.01; 95% CI, 0.68–1.51; ref. 22) were inconsistent, with both inverse and positive associations reported.

β -Cryptoxanthin. Four studies reported data on β -cryptoxanthin in relation to HNC (20, 23, 24, 37). We observed a protective effect of the highest versus the lowest category

**Figure 3.**

Meta-analysis of studies reporting HNC risk in relation to natural intake of α -carotene and β -carotene. Quantile cutoff points (or average intake): Schantz, 1997: >609.5 vs. <181.9 $\mu\text{g}/\text{day}$; Bravi, 2013: mean (SD) = 696.6 (607.3) $\mu\text{g}/\text{day}$ among controls; De Stefani, 1999: not reported; Bidoli, 2003: mean (SD) = 800 (900) $\mu\text{g}/\text{day}$ among controls; Schantz, 1997: $>4,421$ vs. $<2,064$ $\mu\text{g}/\text{day}$; Bravi, 2013: mean (SD) = 4,457.1 (2,303.3) $\mu\text{g}/\text{day}$ among controls; De Stefani, 1999: not reported; Esteve, 1996: $>1,300$ vs. ≤ 400 $\mu\text{g}/\text{day}$; Petridou, 2002: mean (SD) = 3,642 (1,920) $\mu\text{g}/\text{day}$ among cases, mean (SD) = 3,410 (1,870) $\mu\text{g}/\text{day}$ among controls; Bidoli, 2003: mean (SD) = 4,100 (2,600) $\mu\text{g}/\text{day}$ among controls.

of β -cryptoxanthin intake for oral and pharyngeal cancer (OR, 0.46; 95% CI, 0.29–0.74; $I^2 = 51.8\%$; $P = 0.15$) and for laryngeal cancer (OR, 0.41; 95% CI, 0.33–0.51; $I^2 = 0.0\%$; $P = 0.49$; Fig. 4). Only a case-control study from the United States addressed the association between β -cryptoxanthin intake and HNC in general, and observed a consistent protective effect (OR, 0.30; 95% CI, 0.15–0.60; ref. 37).

Lycopene. Seven studies reported data on lycopene intake and HNC (17, 20, 21, 23, 24, 27, 37). The meta-analysis identified lycopene to act protectively against oral and pharyngeal cancer (OR, 0.74; 95% CI, 0.56–0.98; $I^2 = 14.5\%$; $P = 0.32$), as well as against laryngeal cancer (OR, 0.50; 95% CI, 0.28–0.89; $I^2 = 65.9\%$; $P < 0.05$; ref. Fig. 4). Oppositely, a case-control study from the United States did not report a consistent association between lycopene intake and HNC in general (OR, 0.60; 95% CI, 0.32–1.11; ref. 37).

Lutein and zeaxanthin. Four studies reported data of lutein and zeaxanthin in relation to HNC (20, 23, 24, 37). A nonsignificant inverse association was observed for the highest level of intake of lutein and zeaxanthin, both for cancer of the oral cavity and pharynx (OR, 0.51; 95% CI, 0.22–1.18; $I^2 = 83.0\%$; $P < 0.05$) and for that of the larynx (OR, 0.60; 95% CI, 0.27–1.32; $I^2 = 85.8\%$; $P < 0.01$; Fig. 5). Similarly, a case-control study from the United States did not find any significant association between lutein intake and HNC in general (OR, 0.95; 95% CI, 0.52–1.73).

Discussion

Our review reports that the intake of carotenoids from natural sources protects against the development of HNC in relation to most individual carotenoids, with some differences between cancer subsites. We observed that a high intake of β -carotene equivalents significantly decreases the risk for cancer of oral cavity as well as for laryngeal cancer. Lycopene and β -cryptoxanthin also reduced the risk of laryngeal cancer. Lycopene, α -carotene, and β -cryptoxanthin appear to significantly reduce the risk of oral and pharyngeal cancer.


The effect of carotenoid intake was investigated in relation to several cancer sites (39–43). A consistent protective effect was reported so far for cancers of the lung (41), prostate (42), colon (43), cervix (44), and esophagus (40).

Carotenoids represent a diverse group of natural pigments of the polyene type (3). There are several possible cancer prevention mechanisms that can explain the favorable effect of carotenoids toward cancer development (7, 45). Carotenoids can function as a provitamin A having effect on cellular differentiation and proliferation (46, 47). Moreover their anticarcinogenic effect lies in the ability to act as antioxidants, quenching free radicals, reducing damage from reactive oxidant species, and inhibiting lipid peroxidation (45).

Although the bioavailability and absorption of the synthetic form of carotenoids are different with respect to carotenoids from natural sources (48), several studies have reported an increased risk of cancer development in individuals taking

Figure 4.

Meta-analysis of studies reporting HNC risk in relation to natural intake of β -cryptoxanthin and lycopene. Quantile cutoff points (or average intake): Schantz, 1997: >148 vs. <61.4 $\mu\text{g}/\text{day}$; Bravi, 2013: mean (SD) = 463.0 (559.1) $\mu\text{g}/\text{day}$ among controls; De Stefani, 1999: not reported; Bidoli, 2003: mean (SD) = 200 (300) $\mu\text{g}/\text{day}$ among controls; Schantz, 1997: >1,898.4 vs. <910.7 $\mu\text{g}/\text{day}$; Bravi, 2013: mean (SD) = 4,873.2 (3,418.4) $\mu\text{g}/\text{day}$ among controls; De Stefani, 1999: not reported; Negri, 2000: mean (SD) = 7,084.0 (3,763.1) $\mu\text{g}/\text{day}$ among controls; De Stefani, 2000: $\geq 3,388$ vs. $\leq 2,240$ $\mu\text{g}/\text{day}$; Bidoli, 2003: mean (SD) = 6,400 (3,700) $\mu\text{g}/\text{day}$ among controls; Gallus, 2003: not reported.




β -carotene preventively. The Alpha-Tocopherol, Beta-Carotene Cancer Prevention Trial (ATBC) indicated a significantly higher occurrence of lung cancer among heavy cigarette-smoking men who were taking α -tocopherol and β -carotene in compar-

ison to individuals taking the placebo (49). The Beta-Carotene and Retinol Efficacy Trial (CARET) study, as well as others, also reported that the combination of β -carotene and vitamin A supplementation was associated with increased lung cancer

Figure 5.

Meta-analysis of studies reporting HNC risk in relation to natural intake of lutein and zeaxanthin. Quantile cutoff points (or average intake): Schantz, 1997: >3,732 vs. <1,678 $\mu\text{g}/\text{day}$; Bravi, 2013: mean (SD) = 4,381.5 (2,286.3) $\mu\text{g}/\text{day}$ among controls; De Stefani, 1999: not reported; Bidoli, 2003: mean (SD) = 4,300 (2,500) $\mu\text{g}/\text{day}$ among controls; De Stefani, 1999: not reported.



Downloaded from http://aacrjournals.org/cebp/article-pdf/24/7/1003/2280948/1003.pdf by guest on 15 September 2024

risk among men and women at a high risk, such as asbestos workers and smokers, and in subjects who consumed larger amounts of alcohol (50–52). Furthermore, it has been reported that β -carotene could also be susceptible to an alternative mechanism, not having an anti-oxygenic, but a pro-oxygenic, effect (5). These studies suggest that the pro-oxidant effects of carotenoids may be explained in terms of their strong interference with unhealthy lifestyle factors (53).

Few systematic reviews and meta-analyses have investigated the relationship between dietary habits and the risk of HNC. Two of them have reviewed the findings of studies that have examined the association between HNC and carotenoid intake (25, 26). The first review, examining the association of cancers of the upper digestive tract with diet, found that carotene, vitamin C, and vitamin E are protective, most likely in combination with each other and with other micronutrients (25). In the second review, β -carotene has been inversely related to the risk of oral and pharyngeal cancer, although it remains difficult to disentangle its potential effect from that of fruit and vegetables (26). However, in both studies, the effect on specific cancer subsites and the single carotenoids so far was not addressed.

The strength of our review is that we have investigated the effect of individual and combined carotenoids from diet on overall and specific subsites of HNC. In interpreting our results, we acknowledge a few limitations. First, although most of the included studies provided estimates adjusted for major recognized risk factors for HNC, residual confounding from other dietary factors (or other factors, e.g., HPV) cannot be excluded.

In fact, carotenoids intake tends to be associated with healthy behaviors that may be protective against HNC, such as a better educational background. Second, although cigarette smoking is known to be a major risk factor for HNC, we were unable to conduct stratified meta-analyses according to smoking status (54). Lastly, heterogeneity may have inadvertently been introduced because of the methodological differences among studies, including different ranges of exposure used to define categories.

In conclusion, our systematic review and meta-analysis on dietary carotenoids intake and HNC suggests that high levels of carotenoid intake may protect against the risk of HNC, in relation to most of single nutrients and HNC subsites.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Grant Support

This work has been supported by Fondazione Veronesi and Italian Association for Research on Cancer (AIRC, IG 2013, No. 14220). The work of Dr. Nikola Panic and Dr. Darko Nedovic was supported by the ERAWEB project, funded with support of the European Commission.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received January 21, 2015; revised March 16, 2015; accepted April 1, 2015; published OnlineFirst April 14, 2015.

References

- Curado MP, Boyle P. Epidemiology of head and neck squamous cell carcinoma not related to tobacco or alcohol. *Curr Opin Oncol* 2013; 25:229–34.
- GLOBOCAN 2012 v1.0 Cancer incidence and mortality worldwide: IARC CancerBase No.11. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>.
- Landrum JT. Carotenoids: physical, chemical, and biological functions and properties. Boca Raton, FL: CRC Press; 2010.
- Paiva SA, Russell RM. Beta-carotene and other carotenoids as antioxidants. *J Am Coll Nutr* 1999;18:426–33.
- Fiedor J, Fiedor L, Haessner R, Scheer H. Cyclic endoperoxides of beta-carotene, potential pro-oxidants, as products of chemical quenching of singlet oxygen. *Biochim Biophys Acta* 2005;1709:1–4.
- Cvetkovic D, Fiedor L, Fiedor J, Wisniewska-Becker A, Markovic D. Molecular base for carotenoids antioxidant activity in model and biological systems: the health-related effects. In: Yamaguchi M, editor. Carotenoids: food sources, production and health benefits. New York, NY: Nova Science Publishers; 2013. p. 93–126.
- Fiedor J, Burda K. Potential role of carotenoids as antioxidants in human health and disease. *Nutrients* 2014;6:466–88.
- Salter A, Wiseman H, Tucker G. Phytonutrients. West Sussex, UK: Blackwell Publishing Ltd.; 2012. 312 p.
- McLaughlin JK, Gridley G, Block G, Winn DM, Preston-Martin S, Schoenberg JB, et al. Dietary factors in oral and pharyngeal cancer. *J Natl Cancer Inst* 1988;80:1237–43.
- Rossing MA, Vaughan TL, McKnight B. Diet and pharyngeal cancer. *Int J Cancer* 1989;44:593–7.
- Gridley G, McLaughlin JK, Block G, Blot WJ, Winn DM, Greenberg RS, et al. Diet and oral and pharyngeal cancer among blacks. *Nutr Cancer* 1990; 14:219–25.
- Marshall JR, Graham S, Haughey BP, Shedd D, O'Shea R, Brasure J, et al. Smoking, alcohol, dentition and diet in the epidemiology of oral cancer. *Eur J Cancer B Oral Oncol* 1992;28B:9–15.
- Zheng W, Blot WJ, Shu XO, Diamond EL, Gao YT, Ji BT, et al. Risk factors for oral and pharyngeal cancer in Shanghai, with emphasis on diet. *Cancer Epidemiol Biomarkers Prev* 1992;1:441–8.
- Day GL, Blot WJ, Austin DF, Bernstein L, Greenberg RS, Preston-Martin S, et al. Racial differences in risk of oral and pharyngeal cancer: alcohol, tobacco, and other determinants. *J Natl Cancer Inst* 1993; 85:465–73.
- Zheng T, Boyle P, Willett WC, Hu H, Dan J, Evstifeeva TV, et al. A case-control study of oral cancer in Beijing, People's Republic of China. Associations with nutrient intakes, foods and food groups. *Eur J Cancer B Oral Oncol* 1993;29B:45–55.
- Prasad MP, Krishna TP, Pasricha S, Quereshi MA, Krishnaswamy K. Diet and oral cancer—a case control study. *Asia Pac J Clin Nutr* 1995;4:259–64.
- Negri E, Franceschi S, Bosetti C, Levi F, Conti E, Parpinel M, et al. Selected micronutrients and oral and pharyngeal cancer. *Int J Cancer* 2000;86:122–7.
- Suzuki T, Wakai K, Matsuo K, Hirose K, Ito H, Kuriki K, et al. Effect of dietary antioxidants and risk of oral, pharyngeal and laryngeal squamous cell carcinoma according to smoking and drinking habits. *Cancer Sci* 2006; 97:760–7.
- Kune GA, Kune S, Field B, Watson LF, Cleland H, Merenstein D, et al. Oral and pharyngeal cancer, diet, smoking, alcohol, and serum vitamin A and beta-carotene levels: a case-control study in men. *Nutr Cancer* 1993; 20:61–70.
- De Stefani E, Ronco A, Mendilaharsu M, Deneo-Pellegrini H. Diet and risk of cancer of the upper aerodigestive tract—II. *Nutrients. Oral Oncol* 1999; 35:22–6.
- De Stefani E, Oreggia F, Boffetta P, Deneo-Pellegrini H, Ronco A, Mendilaharsu M. Tomatoes, tomato-rich foods, lycopene and cancer of the upper aerodigestive tract: a case-control in Uruguay. *Oral Oncol* 2000;36:47–53.
- Petridou E, Zavras AI, Lefatzis D, Dessypris N, Laskaris G, Dokianakis G, et al. The role of diet and specific micronutrients in the etiology of oral carcinoma. *Cancer* 2002;94:2981–8.

23. Bidoli E, Bosetti C, La Vecchia C, Levi F, Parpinel M, Talamini R, et al. Micronutrients and laryngeal cancer risk in Italy and Switzerland: a case-control study. *Cancer Causes Control* 2003;14:477-84.
24. Bravi F, Bosetti C, Filomeno M, Levi F, Garavello W, Galimberti S, et al. Foods, nutrients and the risk of oral and pharyngeal cancer. *Br J Cancer* 2013;109:2904-10.
25. Chainani-Wu N. Diet and oral, pharyngeal, and esophageal cancer. *Nutr Cancer* 2002;44:104-26.
26. Lucenteforte E, Garavello W, Bosetti C, La Vecchia C. Dietary factors and oral and pharyngeal cancer risk. *Oral Oncol* 2009;45:461-7.
27. Gallus S, Bosetti C, Franceschi S, Levi F, Negri E, La Vecchia C. Laryngeal cancer in women: tobacco, alcohol, nutritional, and hormonal factors. *Cancer Epidemiol Biomarkers Prev* 2003;12:514-7.
28. Wells G, Shea B, O'Connell J, Robertson J, et al. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 2011. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
29. Moher D, Liberati A, Tetzlaff J, Altman DG. The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *BMJ* 2009;339:b2535, doi: 10.1136/bmj.b2535.
30. Jackson D, White IR, Thompson SG. Extending DerSimonian and Laird's methodology to perform multivariate random effects meta-analyses. *Stat Med* 2010;29:1282-97.
31. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
32. Mackerras D, Buffler PA, Randall DE, Nichaman MZ, Pickle LW, Mason TJ. Carotene intake and the risk of laryngeal cancer in coastal Texas. *Am J Epidemiol* 1988;128:980-8.
33. Franceschi S, Bidoli E, Baron AE, Barra S, Talamini R, Serraino D, et al. Nutrition and cancer of the oral cavity and pharynx in north-east Italy. *Int J Cancer* 1991;47:20-5.
34. La Vecchia C, Negri E, D'Avanzo B, Boyle P, Franceschi S. Dietary indicators of oral and pharyngeal cancer. *Int J Epidemiol* 1991;20:39-44.
35. Freudenheim JL, Graham S, Byers TE, Marshall JR, Haughey BP, Swanson MK, et al. Diet, smoking, and alcohol in cancer of the larynx: a case-control study. *Nutr Cancer* 1992;17:33-45.
36. Esteve J, Riboli E, Pequinot G, Terracini B, Merletti F, Crosignani P, et al. Diet and cancers of the larynx and hypopharynx: the IARC multi-center study in southwestern Europe. *Cancer Causes Control* 1996;7:240-52.
37. Schantz SP, Zhang ZF, Spitz MS, Sun M, Hsu TC. Genetic susceptibility to head and neck cancer: interaction between nutrition and mutagen sensitivity. *Laryngoscope* 1997;107:765-81.
38. Zheng W, Sellers TA, Doyle TJ, Kushi LH, Potter JD, Folsom AR. Retinol, antioxidant vitamins, and cancers of the upper digestive tract in a prospective cohort study of postmenopausal women. *Am J Epidemiol* 1995;142:955-60.
39. Freedman ND, Park Y, Subar AF, Hollenbeck AR, Leitzmann MF, Schatzkin A, et al. Fruit and vegetable intake and head and neck cancer risk in a large United States prospective cohort study. *Int J Cancer* 2008;122:2330-6.
40. Cheng KK, Day NE. Nutrition and esophageal cancer. *Cancer Causes Control* 1996;7:33-40.
41. Michaud DS, Feskanich D, Rimm EB, Colditz GA, Speizer FE, Willett WC, et al. Intake of specific carotenoids and risk of lung cancer in 2 prospective US cohorts. *Am J Clin Nutr* 2000;72:990-7.
42. Chan JM, Gann PH, Giovannucci EL. Role of diet in prostate cancer development and progression. *J Clin Oncol* 2005;23:8152-60.
43. McGarr SE, Ridlon JM, Hylemon PB. Diet, anaerobic bacterial metabolism, and colon cancer: a review of the literature. *J Clin Gastroenterol* 2005;39:98-109.
44. Zhang X, Dai B, Zhang B, Wang Z. Vitamin A and risk of cervical cancer: a meta-analysis. *Gynecol Oncol* 2012;124:366-73.
45. Krinsky NI, Johnson EJ. Carotenoid actions and their relation to health and disease. *Mol Aspects Med* 2005;26:459-516.
46. De Luca L, Maestri N, Bonanni F, Nelson D. Maintenance of epithelial cell differentiation: the mode of action of vitamin A. *Cancer* 1972;30:1326-31.
47. Sporn MB, Roberts AB. Role of retinoids in differentiation and carcinogenesis. *Cancer Res* 1983;43:3034-40.
48. Brewer MS. Natural Antioxidants: Sources, compounds, mechanisms of action and potential applications. *Compr Rev Food Sci Food Safety* 2011;10:221-47.
49. The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers: The alpha-tocopherol, beta carotene cancer prevention study group. *N Engl J Med* 1994;330:1029-35.
50. Albanes D, Heinonen OP, Taylor PR, Virtamo J, Edwards BK, Rautalahti M, et al. Alpha-Tocopherol and beta-carotene supplements and lung cancer incidence in the alpha-tocopherol, beta-carotene cancer prevention study: effects of base-line characteristics and study compliance. *J Natl Cancer Inst* 1996;88:1560-70.
51. Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, et al. Risk factors for lung cancer and for intervention effects in CARET, the beta-carotene and retinol efficacy trial. *J Natl Cancer Inst* 1996;88:1550-9.
52. Goodman GE, Thornquist MD, Balmes J, Cullen MR, Meyskens FL Jr, Omenn GS, et al. The beta-carotene and retinol efficacy trial: incidence of lung cancer and cardiovascular disease mortality during 6-year follow-up after stopping beta-carotene and retinol supplements. *J Natl Cancer Inst* 2004;96:1743-50.
53. Goralczyk R. Beta-carotene and lung cancer in smokers: review of hypotheses and status of research. *Nutr Cancer* 2009;61:767-74.
54. Hashibe M, Brennan P, Benhamou S, Castellsague X, Chen C, Curado MP, et al. Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled analysis in the international head and neck cancer epidemiology consortium. *J Natl Cancer Inst* 2007;99:777-89.