

## Null Results in Brief

# No Difference Between Red Wine or White Wine Consumption and Breast Cancer Risk

Polly A. Newcomb,<sup>1,3</sup> Hazel B. Nichols,<sup>3</sup> Jeannette M. Beasley,<sup>1,2</sup> Kathleen Egan,<sup>5</sup> Linda Titus-Ernstoff,<sup>6</sup> John M. Hampton,<sup>3</sup> and Amy Trentham-Dietz<sup>3,4</sup>

<sup>1</sup>Fred Hutchinson Cancer Research Center and <sup>2</sup>Group Health Cooperative, Seattle, Washington; <sup>3</sup>University of Wisconsin Paul P. Carbone Comprehensive Cancer Center and <sup>4</sup>Department of Population Health Sciences, University of Wisconsin, Madison, Wisconsin; <sup>5</sup>H. Lee Moffitt Cancer Center & Research Institute, Tampa, Florida; and <sup>6</sup>Dartmouth Medical School, Lebanon, New Hampshire

## Introduction

Epidemiologic studies have reported an increased risk of breast cancer among women who drink alcohol, including wine (1, 2). Two meta-analyses estimated a ~10% [95% confidence interval (CI), 5-15%] increased risk of breast cancer with each additional 10 grams (~1 drink) of alcohol/day regardless of beverage type (3, 4). Few studies have evaluated breast cancer risk separately for red and white wine (5-8). There is some evidence of beneficial health effects of red wine from laboratory (9) and epidemiologic studies of heart disease (10) and prostate cancer (11, 12). We evaluated overall alcohol as well as red and white wine consumption to examine beverage-specific effects on breast cancer.

## Materials and Methods

The analysis was conducted in a population-based case-control study described fully elsewhere (13). Briefly, the study enrolled breast cancer cases identified from population-based registries in Wisconsin, Massachusetts (excluding metropolitan Boston), and New Hampshire, age 20 to 69 y, with incident invasive breast cancer diagnosed between 1995 and 2000. Controls were frequency matched to cases by 5-y age groups, selected from lists of licensed drivers (<65 y) or a roster of Medicare beneficiaries (≥65 y). Approximately 80% of cases ( $n = 6,327$ ) and 76% of controls ( $n = 7,558$ ) participated in the study; participation was similar in the three states.

Telephone interviews were administered concurrently for case (typically within 1 y of diagnosis) and control participants. Participants were queried on suspected risk factors for breast cancer, including age at menarche and menopause, age at first pregnancy, parity, postmeno-

pausal hormone use, family history of breast cancer, smoking, education, and body mass index evaluated before the reference date. Alcohol use was evaluated as frequency of beer, red wine, white wine, or liquor (distilled spirits) consumption (number of drinks per day, week, or month) during the year previous to the reference date, defined as the date of diagnosis for cases or approximately the same period for controls. Logistic regression was used to estimate odds ratios (OR) and 95% CIs. Variables were included in the multivariable model if statistically significantly associated with breast cancer ( $P \leq 0.05$ ) in age-adjusted models. Models estimating the effect of wine type on risk were adjusted for all other sources of alcohol. To obtain  $P_{\text{trend}}$ , we included alcohol both as a continuous and categorical exposure. We tested for interaction by type of alcohol. Analyses were conducted using SAS (Version 9.1; SAS Institute).

## Results

Cases tended to have a younger age at menarche, later age at first birth, lower parity, and later age at menopause (data not shown). Current use of postmenopausal hormones, first-degree family history of breast cancer, greater body mass index, and higher education were also associated with increased breast cancer.

Any alcohol consumption in the year before the reference date was reported by similar proportions of cases (82%) and controls (82%). After multivariable adjustment, consumption of 14 or more drinks per week was associated with a 24% increase in breast cancer (95% CI, 3-49%;  $P_{\text{trend}} = 0.003$ ) compared with non-drinkers (Table 1). Menopausal status modified this association, with elevated risk observed only among postmenopausal women ( $P_{\text{interaction}} = 0.05$ ). This effect was driven by a statistical interaction only among liquor drinkers. Higher intake of all beverage types was suggestive of increased risk; only elevated consumption of liquor was significantly elevated per drink (OR, 1.02; 95% CI, 1.00-1.03). Wine consumption was not associated with risk of breast cancer (OR, 1.01; 95% CI, 0.99-1.02) and no differential was observed between red and white wine ( $P_{\text{interaction}} = 0.6$ ; Table 2).

Cancer Epidemiol Biomarkers Prev 2009;18(3):1007-10

Received 8/27/08; revised 12/9/08; accepted 12/30/08.

**Grant support:** National Cancer Institute grants R01 CA47147, R01 CA47305, and R01 CA69664.

**Requests for reprints:** Polly A. Newcomb, Fred Hutchinson Cancer Center, M4-B402, 1100 Fairview Avenue North, Seattle, WA 98109-1024. Phone: 206-667-3476; Fax: 206-667-7850. E-mail: pnewcomb@fhcrc.org

Copyright © 2009 American Association for Cancer Research.

doi:10.1158/1055-9965.EPI-08-0801

## Discussion

In this large, multicenter, population-based case-control study, alcohol was associated with an elevated risk of breast cancer, primarily because of increased risk in postmenopausal liquor drinkers. Neither red nor white wine was related to breast cancer.

The Women's Health Study, a randomized, chemopreventive trial of 39,876 health professionals followed for ~10 years, reported similar associations of white (multivariable relative risk, 1.1; 95% CI, 0.9-1.2) and red (multivariable relative risk, 1.0; 95% CI, 0.8-1.2) wine per 10 grams/day increment on risk of breast cancer (6). Furthermore, two other studies conducted in France reported findings that did not support inverse effects of red wine versus white wine. Hirvonen et al. (8) reported no significantly increased risk of breast cancer associated with either red or white wine after a median 6.6 years follow-up in a placebo-controlled antioxidant primary-prevention trial with 95 breast cancer cases. A small

case-control study of premenopausal women ( $n = 154$  cases and 154 controls) reported an increase in breast cancer for women who reported consuming >4 liters of red wine per month, or ~1 drink/day (OR, 4.0; 95% CI, 1.6-9.8;  $P_{\text{trend}} = 0.003$ ; ref. 5). Whereas over half of the study population consumed red wine, <10% reported consuming white wine (5). This differs from our study where equal proportions of women reported consuming red and white wine. Another case-control study conducted in France reported a lower risk of breast cancer among women consuming 10 to 12 grams/day of wine (OR, 0.51; 95% CI, 0.30-0.97) when compared with nonwine drinkers, but they did not evaluate red and white wine separately (14).

Separate effects of red versus white wine on risk may have biological plausibility. Resveratrol, a polyphenol in grapes, peanuts, and other plant-based foods, has displayed anticarcinogenic properties in mouse mammary cultures (15, 16). Concentrations are greater in red compared with white wine, due to the variable amounts

**Table 1. Association of recent alcoholic beverage consumption and breast cancer risk**

Drinks per week	All women			Premenopausal women			Postmenopausal women <sup>*,†</sup>		
	Cases ( $n = 6,327$ )	Controls ( $n = 7,558$ )	OR <sup>‡</sup> (95% CI)	Cases ( $n = 2,462$ )	Controls ( $n = 2,925$ )	OR <sup>‡</sup> (95% CI)	Cases ( $n = 3,449$ )	Controls ( $n = 4,162$ )	OR <sup>‡</sup> (95% CI)
All alcohol	No. (%)	No. (%)		No. (%)	No. (%)		No. (%)	No. (%)	
Nondrinkers	1,122 (18)	1,379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	2,129 (34)	2,712 (36)	0.94 (0.85-1.04)	894 (36)	1,045 (36)	1.02 (0.86-1.22)	1,106 (32)	1,480 (36)	0.90 (0.79-1.02)
1-3.4	1,450 (23)	1,731 (23)	0.99 (0.88-1.10)	592 (24)	767 (26)	0.90 (0.75-1.08)	745 (22)	863 (21)	1.03 (0.89-1.19)
3.5-6.9	702 (11)	793 (11)	1.02 (0.90-1.17)	306 (12)	363 (12)	0.96 (0.78-1.19)	345 (10)	384 (9)	1.05 (0.88-1.26)
7-13.9	619 (10)	658 (9)	1.11 (0.97-1.28)	221 (9)	223 (8)	1.17 (0.92-1.48)	355 (10)	395 (10)	1.07 (0.89-1.28)
≥14	305 (5)	285 (4)	1.24 (1.03-1.49)	98 (4)	103 (4)	1.10 (0.80-1.52)	189 (6)	164 (4)	1.37 (1.08-1.73)
$P_{\text{trend}}$			0.003			0.55			0.004
1-drink increase			1.01 (1.00-1.02)			1.00 (0.99-1.01)			1.02 (1.01-1.03)
Wine									
Nondrinkers	1,122 (18)	1,379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	2,455 (39)	2,920 (39)	1.03 (0.94-1.12)	1,021 (42)	1,183 (40)	1.09 (0.95-1.26)	1,283 (37)	1,550 (37)	0.97 (0.86-1.09)
1-3.4	1,048 (17)	1,248 (17)	1.00 (0.89-1.11)	424 (17)	546 (19)	0.98 (0.82-1.17)	538 (16)	634 (15)	0.95 (0.81-1.11)
3.5-6.9	343 (5)	363 (5)	1.13 (0.95-1.34)	134 (5)	149 (5)	1.15 (0.87-1.50)	179 (5)	192 (5)	1.05 (0.83-1.33)
7-13.9	256 (4)	299 (4)	1.07 (0.89-1.29)	90 (4)	86 (3)	1.42 (1.02-1.96)	147 (4)	192 (5)	0.93 (0.73-1.19)
≥14	82 (1)	70 (1)	1.37 (0.98-1.91)	29 (1)	24 (1)	1.58 (0.90-2.79)	47 (1)	43 (1)	1.23 (0.80-1.90)
$P_{\text{trend}}$			0.20			0.12			0.84
1-drink increase			1.01 (0.99-1.02)			1.00 (0.99-1.02)			1.01 (0.99-1.03)
Beer									
Nondrinkers	1,122 (18)	1,379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	1,756 (28)	2,040 (27)	1.01 (0.93-1.10)	714 (29)	870 (30)	0.92 (0.80-1.05)	916 (27)	1,044 (25)	1.05 (0.94-1.18)
1-3.4	754 (12)	836 (11)	1.06 (0.95-1.20)	359 (15)	444 (15)	0.91 (0.77-1.08)	345 (10)	346 (8)	1.21 (1.02-1.44)
3.5-6.9	163 (3)	210 (3)	0.90 (0.72-1.12)	105 (4)	120 (4)	0.91 (0.69-1.21)	48 (1)	76 (2)	0.81 (0.55-1.17)
7-13.9	129 (2)	139 (2)	1.08 (0.84-1.39)	52 (2)	55 (2)	1.05 (0.71-1.56)	66 (2)	74 (2)	1.08 (0.76-1.52)
≥14	82 (1)	84 (1)	1.12 (0.81-1.53)	32 (1)	39 (1)	0.89 (0.55-1.45)	45 (1)	39 (1)	1.44 (0.92-2.25)
$P_{\text{trend}}$			0.55			0.28			0.11
1-drink increase			1.01 (1.00-1.02)			1.01 (0.99-1.02)			1.01 (1.00-1.03)
Liquor									
Nondrinkers	1,122 (18)	1,379 (18)	1	351 (14)	424 (15)	1	709 (21)	876 (21)	1
<1	2,463 (39)	3,132 (41)	0.90 (0.83-0.98)	1,098 (45)	1,371 (47)	0.90 (0.79-1.02)	1,186 (34)	1,529 (37)	0.93 (0.83-1.04)
1-3.4	940 (15)	1,044 (14)	1.00 (0.90-1.12)	334 (14)	402 (14)	0.92 (0.77-1.11)	541 (16)	582 (14)	1.07 (0.92-1.24)
3.5-6.9	154 (2)	146 (2)	1.14 (0.90-1.44)	55 (2)	61 (2)	0.93 (0.63-1.37)	93 (3)	75 (2)	1.40 (1.02-1.93)
7-13.9	162 (3)	175 (2)	1.00 (0.80-1.26)	37 (2)	38 (1)	1.00 (0.62-1.60)	115 (3)	131 (3)	0.98 (0.75-1.29)
≥14	95 (2)	73 (1)	1.41 (1.03-1.94)	18 (1)	19 (1)	1.10 (0.56-2.13)	74 (2)	50 (1)	1.63 (1.12-2.38)
$P_{\text{trend}}$			0.28			0.37			0.04
1-drink increase			1.02 (1.00-1.03)			0.99 (0.97-1.02)			1.03 (1.01-1.05)

\*For the test of the interaction between total alcohol consumption (continuous) and menopausal status  $P = 0.05$ .

†For the test of the interaction between alcohol consumption (continuous) and menopausal status: all alcohol,  $P = 0.05$ ; wine,  $P = 0.89$ ; beer,  $P = 0.89$ ; liquor,  $P = 0.03$ .

‡Adjusted for age, state, family history of breast cancer, age at menarche, age at first birth, parity, menopausal status, age at menopause, postmenopausal hormone use, body mass index, education, wine, beer, and liquor consumption.

**Table 2. Association of recent red and white wine consumption and breast cancer risk**

Wine type	Cases ( <i>n</i> = 6,327)	Controls ( <i>n</i> = 7,558)	OR (95% CI)* <sup>†</sup>
Drinks per week	No. (%)	No. (%)	
<b>Red wine</b>			
Nondrinkers	1,122 (18)	1,379 (18)	1
<1	2,145 (34)	2,502 (33)	1.03 (0.94-1.12)
1-3.4	738 (12)	885 (12)	0.94 (0.83-1.06)
3.5-6.9	98 (2)	119 (2)	0.91 (0.69-1.21)
7-13.9	124 (2)	133 (2)	1.17 (0.90-1.51)
≥14	28 (<1)	24 (<1)	1.27 (0.73-2.22)
<i>P</i> <sub>trend</sub>			0.99
1-drink increase			1.00 (0.98-1.01)
<b>White wine</b>			
Nondrinkers	1,122 (18)	1,379 (18)	1
<1	2,145 (34)	2,587 (34)	0.98 (0.90-1.07)
1-3.4	880 (14)	997 (13)	1.06 (0.94-1.19)
3.5-6.9	161 (3)	143 (2)	1.40 (1.10-1.78)
7-13.9	103 (2)	137 (2)	0.90 (0.69-1.18)
≥14	40 (1)	34 (<1)	1.37 (0.86-2.19)
<i>P</i> <sub>trend</sub>			0.34
1-drink increase			1.02 (1.00-1.04)

\*Adjusted for age, state, family history of breast cancer, age at menarche, age at first birth, parity, menopausal status, age at menopause, postmenopausal hormone use, body mass index, education, and total alcohol consumption (minus red or white wine consumption).

<sup>†</sup>For the test of the interaction wine type (red versus white, continuous) *P* = 0.60.

of time that red grape skins are present during processing (17). However, resveratrol is metabolized quickly: consistent intake of two glasses of red wine daily would be equivalent to a dose of ~27 μg/kg body weight for a 70-kg individual, leading to detectable concentrations of derivatives but not free resveratrol (9). Thus, these low levels may not result in bioeffective concentrations.

Limitations should be considered in interpreting our findings. Exposure measurement was limited to recent use, and patterns may have changed over time. Although we were able to control for many confounders, we did not have data on drinking patterns (i.e., moderate versus binge drinking). Self-reported alcohol intake measures are prone to error due to both retrospective reporting and recall bias. However, alcohol intake measured via food frequency is correlated with food records (Spearman *r* = 0.9; ref. 18) and the reliability of reporting is acceptable (*r* = 0.8; ref. 19).

Wine consumption has been increasing over the past decade in the United States (20), so future studies may more precisely evaluate the effects red and white wine on incidence of breast cancer.

### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

### Acknowledgments

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank Drs. Henry Anderson, Patrick L. Remington, Meir J. Stampfer, Walter C. Willett, John A. Baron, E. Robert Greenberg, Laura Stephenson and the staff of the Wisconsin Cancer Reporting System, Susan T. Gershman, and the staff of the Massachusetts Tumor Registry; Marguerite Stevens and the staff

of the New Hampshire Cancer Registry; Linda Haskins, Heidi Judge, Laura Mignone, and Shafika Abrahams-Gessel along with the study interviewers and programmers in all three states for assistance with data collection; and the study participants, whose generosity made this research possible.

### References

- Smith-Warner SA, Spiegelman D, Yaun SS, et al. Alcohol and breast cancer in women: a pooled analysis of cohort studies. *JAMA* 1998; 279:535-40.
- WCFR/AIRC. Alcoholic Drinks. In: Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective. Washington (DC): AICR; 2007. p. 157-71.
- Ellison RC, Zhang Y, McLennan CE, Rothman KJ. Exploring the relation of alcohol consumption to risk of breast cancer. *Am J Epidemiol* 2001;154:740-7.
- Key J, Hodgson S, Omar RZ, et al. Meta-analysis of studies of alcohol and breast cancer with consideration of the methodological issues. *Cancer Causes Control* 2006;17:759-70.
- Viel JF, Perarnau JM, Challier B, Faivre-Nappey I. Alcoholic calories, red wine consumption and breast cancer among premenopausal women. *Eur J Epidemiol* 1997;13:639-43.
- Zhang SM, Lee IM, Manson JE, et al. Alcohol consumption and breast cancer risk in the Women's Health Study. *Am J Epidemiol* 2007;165: 667-76.
- Vachon CM, Kushi LH, Cerhan JR, Kuni CC, Sellers TA. Association of diet and mammographic breast density in the Minnesota breast cancer family cohort. *Cancer Epidemiol Biomarkers Prev* 2000;9: 151-60.
- Hirvonen T, Mennen LI, de Bree A, et al. Consumption of antioxidant-rich beverages and risk for breast cancer in French women. *Ann Epidemiol* 2006;16:503-8.
- Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the *in vivo* evidence. *Nat Rev Drug Discov* 2006;5:493-506.
- Klatsky AL, Friedman GD, Armstrong MA, Kipp H. Wine, liquor, beer, and mortality. *Am J Epidemiol* 2003;158:585-95.
- Schoonen WM, Salinas CA, Kiemeny LA, Stanford JL. Alcohol consumption and risk of prostate cancer in middle-aged men. *Int J Cancer* 2005;113:133-40.
- Velicer CM, Kristal A, White E. Alcohol use and the risk of prostate cancer: results from the VITAL cohort study. *Nutr Cancer* 2006;56: 50-6.
- Sprague BL, Trentham-Dietz A, Newcomb PA, et al. Lifetime recreational and occupational physical activity and risk of *in situ*

- and invasive breast cancer. *Cancer Epidemiol Biomarkers Prev* 2007; 16:236–43.
14. Bessaoud F, Daures JP. Patterns of alcohol (especially wine) consumption and breast cancer risk: a case-control study among a population in Southern France. *Ann Epidemiol* 2008; 18:467–75.
  15. Whitsett T, Carpenter M, Lamartiniere CA. Resveratrol, but not EGCG, in the diet suppresses DMBA-induced mammary cancer in rats. *J Carcinog* 2006;5:15.
  16. Jang M, Cai L, Udeani GO, et al. Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 1997;275: 218–20.
  17. Kopp P. Resveratrol, a phytoestrogen found in red wine. A possible explanation for the conundrum of the 'French paradox'? *Eur J Endocrinol* 1998;138:619–20.
  18. Giovannucci E, Colditz G, Stampfer MJ, et al. The assessment of alcohol consumption by a simple self-administered questionnaire. *Am J Epidemiol* 1991;133:810–7.
  19. Longnecker MP, Newcomb PA, Mittendorf R, et al. The reliability of self-reported alcohol consumption in the remote past. *Epidemiology* 1992;3:535–9.
  20. Wine Institute DoC. San Francisco: c 2005–2008 [cited 2007 Sept 11]. Available from: [http://www.wineinstitute.org/industry/statistics/2007/wine\\_sales.php](http://www.wineinstitute.org/industry/statistics/2007/wine_sales.php).