

A Calcium-Rich Multimineral Intervention to Modulate Colonic Microbial Communities and Metabolomic Profiles in Humans: Results from a 90-Day Trial

Muhammad N. Aslam¹, Christine M. Bassis², Ingrid L. Bergin³, Karsten Knuver¹, Suzanna M. Zick^{4,5}, Ananda Sen^{4,6}, D. Kim Turgeon⁷, and James Varani¹



ABSTRACT

Aquamín is a calcium-, magnesium-, and multiple trace element-rich natural product with colon polyp prevention efficacy based on preclinical studies. The goal of this study was to determine the effects of Aquamin on colonic microbial community and attendant metabolomic profile. Thirty healthy human participants were enrolled in a 90-day trial in which Aquamin (delivering 800 mg of calcium per day) was compared with calcium alone or placebo. Before and after the intervention, colonic biopsies and stool specimens were obtained. All 30 participants completed the study without serious adverse event or change in liver and renal function markers. Compared with pretreatment values, intervention with Aquamin led to a reduction in total bacterial DNA ($P = 0.0001$) and a shift in the microbial community measured by

thetaYC (θ_{YC} ; $P = 0.0087$). Treatment with calcium also produced a decline in total bacteria, but smaller than seen with Aquamin, whereas no reduction was observed with placebo in the colon. In parallel with microbial changes, a reduction in total bile acid levels ($P = 0.0375$) and a slight increase in the level of the short-chain fatty acid (SCFA) acetate in stool specimens ($P < 0.0001$) from Aquamin-treated participants were noted. No change in bile acids or SCFAs was observed with calcium or placebo. We conclude that Aquamin is safe and tolerable in healthy human participants and may produce beneficial alterations in the colonic microbial community and the attendant metabolomic profile. Because the number of participants was small, the findings should be considered preliminary.

Introduction

Epidemiologic studies have shown an inverse relationship between calcium intake and colon cancer incidence (1). Experimental studies in animals have substantiated antitumor efficacy in the colon (2), and *in vitro* studies have provided mechanistic insight into how calcium influences epithelial cell proliferation and differentiation (3). In spite of these data, interventional trials with calcium have had only modest success in reducing colon polyp formation, with some trials demonstrating a reduction in incidence (4, 5), while others showing

essentially no protection (6, 7) or an increase in incidence of colon polyps with a more-aggressive sessile-serrated phenotype (8). This complicated picture supports the idea that adequate or optimal calcium intake throughout life may be beneficial, but in a realistic dietary setting, the efficacy of supplementation by calcium alone may be limited by dietary complexity. Potentially adverse effects at higher dose levels further complicate the picture (9).

Recent evidence suggests that the antineoplastic activity of calcium supplementation may be enhanced by concomitant inclusion of additional trace minerals along with calcium. This has been demonstrated in our own studies using Aquamin, a calcium- and magnesium-rich multimineral product obtained from mineralized red marine algae. In two long-term (15–18 months) studies in mice, Aquamin more effectively suppressed colon polyp formation than calcium alone (10, 11). Further, Aquamin was more effective than calcium alone at suppressing proliferation and inducing differentiation in human colon carcinoma cells in monolayer culture (12, 13) and human adenoma-derived colonoids (14). Thus, Aquamin (as a source of calcium along with magnesium and additional trace minerals) may, ultimately, prove to be a more effective dietary colon polyp chemopreventive agent than calcium alone.

Exact mechanisms by which calcium alone or multimineral supplementation exerts antineoplastic activity are currently unknown. In addition to direct antiproliferative and prodifferentiating effects on colonic epithelium, beneficial activity may be mediated indirectly through effects on the gut microbial

¹Department of Pathology, The University of Michigan Medical School, Ann Arbor, Michigan. ²Division of Infectious Diseases, Department of Internal Medicine, The University of Michigan Medical School, Ann Arbor, Michigan. ³The Unit for Laboratory Animal Medicine, The University of Michigan Medical School, Ann Arbor, Michigan. ⁴Department of Family Medicine, The University of Michigan Medical School, Ann Arbor, Michigan. ⁵Department of Nutritional Science, The University of Michigan School of Public Health, Ann Arbor, Michigan. ⁶Department of Biostatistics, The University of Michigan Medical School, Ann Arbor, Michigan. ⁷Division of Gastroenterology, Department of Internal Medicine, The University of Michigan Medical School, Ann Arbor, Michigan.

Note: Supplementary data for this article are available at Cancer Prevention Research Online (<http://cancerprevres.aacrjournals.org/>).

Corresponding Author: Muhammad N. Aslam, University of Michigan Medical School, 1150 W Medical Center Dr., Ann Arbor, MI 48109. Phone: 734-936-1897; E-mail: mnaslam@med.umich.edu

Cancer Prev Res 2020;13:101-16

doi: 10.1158/1940-6207.CAPR-19-0325

©2019 American Association for Cancer Research.

population and/or changes in gut microbial metabolic activity. The bacterial community in the gastrointestinal tract plays several important roles that contribute to health. A “healthy” microbiome is important for food digestion and production of important metabolites. In addition, a healthy microbial community helps maintain the tissue barrier, regulates the host immune response, and provides protection against pathogen overgrowth (15). Dysbiosis, especially in the colon, can lead to barrier breakdown and initiate a chronic inflammatory response. Dysbiosis has been directly linked to inflammatory bowel diseases and indirectly to the formation of premalignant colon polyps (15–17). In previous studies in mice, dietary calcium supplementation has been shown to cause a shift in the gut microbial community in comparison with control (18, 19). In another study, Aquamin itself induced gut microbial changes (20).

Metabolic changes may also be important. At high concentrations, bile acids are membrane-active agents and can be cytotoxic (21). In addition, certain bacterially derived secondary bile acids have carcinogenic properties (22). Alterations in the gut microbial population can affect bile acid profiles (23). In our own murine study (19), calcium supplementation altered the gut microbiome and reduced the level of total bile acids and certain microbially derived secondary bile acids. In addition to bile acids, gut microbes also produce short-chain fatty acids (SCFA; e.g., acetate, propionate, and butyrate) in the colon. SCFAs have demonstrable protective effect against colonic inflammation and carcinogenesis (24). Thus, a shift in the gut microbial community may affect colon health through alterations in bacterially derived metabolites.

Whether similar microbial/metabolic changes might also be seen in human participants is not known. To test the feasibility of using dietary Aquamin as an interventional strategy in humans, we conducted a 90-day, FDA-approved pilot study, comparing daily supplementation with Aquamin with calcium alone and placebo in healthy human participants. The main purpose of the study was to assess Aquamin's effect on gut microbial community structure and bile acid, SFCA, and eicosanoid profiles at endpoint in comparison with baseline values in participants from all three groups. Elucidating the microbial and metabolic changes resulting from intervention is a first step toward identifying potential mechanisms of action with Aquamin. Identifying measurable alterations in microbial and metabolomic parameters and determining effect-size with intervention will also be helpful for determining the necessary sample size going forward in a larger-scale clinical trial. As part of the study, we also obtained information on tolerability and safety of Aquamin. Initial results from the current study are reported herein.

Materials and Methods

Aquamin and control interventions

Aquamin is a calcium- and magnesium-rich natural product obtained from the skeletal remains of red marine algae of the *Lithothamnion* genus. In addition to calcium and magnesium,

Aquamin contains detectable levels of 72 additional trace minerals including trace elements from the lanthanide family (essentially all of the minerals accumulated by the algae from seawater). Aquamin is sold as a food supplement (GRAS 000028; Marigot Ltd.) and is used in various products for human consumption in Europe, Asia, Australia, and North America. A single batch of Aquamin-Food Grade was used for this study. Mineral composition was established via an independent laboratory (Advanced Laboratories). Supplementary Table S1 provides a complete list of elements detected in Aquamin and their relative amounts. Aquamin has been used in past clinical studies in human subjects with no serious safety or tolerability issues reported (25–27). Calcium carbonate was used for active comparison, and maltodextrin was used as a placebo.

Study participants

This study was a pilot, double-blind, parallel assignment, randomized clinical interventional trial in which 30 participants were included. Participants were male or nonpregnant female, in general good health, but having “an increased risk for colon cancer” based on (i) a personal history of colorectal polyp, early stage (stage I or II) colon cancer treated by surgical removal without recommendation for adjuvant therapy, or stage III colon cancer treated with surgery > 5 years prior or (ii) a first-degree relative diagnosed under the age of 60 with colorectal cancer. Exclusion criteria included history of kidney disease or kidney stones, Crohn's disease or ulcerative colitis, gastrointestinal hemorrhagic disorders, or coagulopathy, hereditary nonpolyposis coli, or familial adenomatous polyposis. The participants were recruited through the Michigan Medicine web portal and by posting flyers in the hospital.


This clinical interventional trial was conducted with FDA approval of Aquamin as an Investigational New Drug (IND#118194) and with oversight by the Institutional Review Board at the University of Michigan Medical School (IRBMED)—IRB#HUM00076276. The study was registered as an interventional clinical trial with details at Clinicaltrials.gov (study identifier NCT02647671). All participants provided written informed consent prior to inclusion. This phase I trial involving human participants was carried out in accordance with recognized ethical guidelines, for example, Declaration of Helsinki, International Ethical Guidelines for Biomedical Research Involving Human Subjects (CIOMS), the Belmont Report, and the U.S. Common Rule.

Study design

Figure 1 summarizes the study design of this trial in a flowchart. Briefly, at screening participants were given the NIH Diet History Questionnaire II (DHQ II), a food frequency questionnaire which includes portion size and dietary supplement questions as a way to evaluate baseline calcium levels in the past year (28). Participants were also asked about use of dietary supplements, antibiotics, and nonsteroidal anti-inflammatory drugs. No participants were on antibiotics at or

Figure 1.

Study design. Study flow diagram highlighting enrollment, group randomization, intervention allocation, study duration, and study sample collection plan.



in close proximity to the time of screening. Individuals ingesting supplements containing calcium and/or vitamin D were required to undergo a 2-week “wash out” period prior to starting and to not use these supplements during the study participation.

Thirty participants underwent baseline flexible sigmoidoscopy (unprepped; i.e., without bowel cleansing procedure). Twelve 2.5 mm colonic biopsies were obtained along with two stool specimens from within the sigmoid colon (20 cm above the anus). Tissue and stool samples were saved in 10% formalin, cryopreserved for cultures or snap frozen in liquid nitrogen, and saved at -80°C . Blood was drawn for the complete metabolic panel including liver function/liver injury markers (total albumin, bilirubin, aspartate aminotransferase, alanine aminotransferase, and alkaline phosphatase) and renal function tests (blood urea nitrogen and creatinine).

After baseline sigmoidoscopy, participants were randomized to one of three groups. Ten participants were treated daily for 90 days with Aquamin providing 800 mg of calcium per day. Ten participants received 800 mg of calcium carbonate daily, and ten participants received maltodextrin as placebo. During the interventional period, participants were contacted by study coordinators on a monthly basis to assess study progress/adherence to the study protocol and to identify unwanted side effects. Compliance was assessed by capsule log entries and by counting unused capsules returned at the end of the study.

At the end of the 90-day intervention period (90 ± 5 days), participants again underwent unprepped flexible sigmoidoscopy and eight colonic biopsies along with two stool specimens were collected and stored as at baseline. Blood was also taken for the same serum markers as at baseline. For each of the two

timepoints, one biopsy and one stool specimen from each participant were utilized for microbiome analysis, and one biopsy and one stool specimen were utilized for metabolomic analysis. The remaining tissue samples were retained for backup or were used for histologic, IHC, and proteomic analyses. The results of the latter analyses will be reported separately.

Microbial analysis

DNA was isolated from colon and stool samples using the Qiagen MagAttract PowerMicrobiome DNA/RNA kit. Total bacterial DNA levels were estimated using qPCR with broad-range primers BSF8 (AGAGTTTGATCCTGGCTCAG) and BSR357 (CTGCTGCCTYCCGTA), targeting bacterial 16S rRNA genes (29). Each 10 μL qPCR reaction contained 5 μL PowerUp SYBR Green Master Mix (Applied Biosystems by Thermo Fisher Scientific), 4 μL sample DNA (1:40 dilution), and 0.4 $\mu\text{mol/L}$ of primers. The following cycles were run on a Light Cycler 96 (Roche): 1x(2 minutes at 50°C , 10 minutes at 95°C), 40x(15 seconds at 95°C , 1 minute at 60°C). Each sample was run in duplicate.

Microbial community profiles were generated by Illumina MiSeq sequencing of the V4 region of 16S rRNA-encoding genes after amplifying the extracted colon tissue and stool DNA of all the participants as described previously (30). Samples were amplified, normalized, and sequenced on the MiSeq, and analysis was performed using the MiSeq SOP (31) for the software mothur (v.1.39.0 and v.1.39.5) as described in our earlier study (19). In case of low bacterial biomass, 3 μL of DNA were amplified by touchdown PCR method [1x(2 minutes at 95°C), 20x(20 seconds at 95°C , 15 seconds at annealing

temperature (starts at 60°C, decreases 0.3°C/cycle), 5 minutes at 72°C, 20x(20 seconds at 95°C, 15 seconds at 55°C, 5 minutes at 72°C), 1x(10 minutes at 72°C)]. Thirteen samples, including 8 of 10 postinterventional Aquamin colon samples, required touchdown PCR (with 40 amplification cycles vs. standard PCR with 30 cycles) to achieve sufficient sample amplification for sequencing. After processing, sequences were binned into operational taxonomic units (OTU) based on 3% difference in sequence using the OptiClust method (32). Pre- and post-comparisons between groups included differences in community structure using OTU-based Yue and Clayton distance metric (θ_{YC} ; ref. 33). θ_{YC} distances were visualized using principal coordinates analysis (PCoA). θ_{YC} is a beta diversity metric. Specific OTUs driving community differences were identified by linear discriminant analysis effect size (LEfSe; ref. 34), as was done in our previous study (19). LEfSe utilizes both statistical significance and effect size (linear discriminant analysis score, or LDA) to determine the features (OTUs, in this case) that are differentially abundant between groups (i.e., pre- and postdifferences). FDRs of the significant LEfSe data were corrected using the *p.adjust()* function with the Benjamini-Hochberg algorithm employing *R*, and stated as *q* values. Alpha diversity was assessed by change in the Shannon diversity index from baseline. To evaluate Aquamin-associated microbial community differences in major gut phyla, we sorted the 1,000 most abundant OTUs in each of the baseline and final visit samples into phyla and compared intergroup differences in each of these phyla at endpoint. The DNA isolation and 16S rRNA gene sequencing were done by the University of Michigan Microbial Systems Molecular Biology Laboratory. Sequence data generated in this project are accessible within the NCBI Sequence Read Archive database (BioProject accession number: PRJNA575562).

Metabolomic analysis

Bile acids, SCFAs, and eicosanoid composition were quantified in colon biopsies and stool specimens from a randomly chosen subset of participants ($n = 6$ per group) at each time point (pre- and postintervention). Bile acids were quantified using LC-MS in a two-step solvent extraction (35) as performed in the Regional Comprehensive Metabolomics Resource Cores (RCMRC) at the University of Michigan. Supernatants were combined, dried, and resuspended for LC-MS separation by reverse-phase liquid chromatography (RPLC) and measured by multiple reaction monitoring (MRM). Sample identification was performed by comparison of retention times and mass with an in-house library of bile acids. SCFA analysis was performed by the RCMRC by electron ionization-gas chromatography-mass spectrometry as described recently (36). Compound identification was performed against a library of known SCFAs. Colon tissue was also utilized to analyze a profile of eicosanoids. Eicosanoids were extracted and concentrated using solid phase extraction in colon biopsies. The eluent was dried and resuspended for LC-MS separation by RPLC and measurements by MRM methods (37). Stool specimens were not analyzed for eicosanoids because these metabolites are not bacterial pro-

ducts. All analytes were reported as pmol/mg, after normalization to the sample weight. Metabolomics data acquired in this study are available on the Metabolomics Workbench (Project ID: PR000822; doi: 10.21228/M8F69D) as part of National Metabolomics Data Repository.

Statistical analysis

Pre (baseline) versus post (endpoint) comparisons were conducted for each microbial or metabolomic endpoint for individual participants in each of the three treatment groups (placebo, calcium, Aquamin). Group means and SDs were then calculated for baseline and endpoint measures. Intergroup differences for normally distributed continuous data were made by ANOVA followed by pairwise group comparisons with Bonferroni corrections for multiple comparisons to generate an adjusted *P* value. The two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli was applied to correct for multiple comparisons by controlling the FDR. Microbial community distances were compared by analysis of molecular variance (AMOVA) within the program mothur (v.1.39.0 and v.1.39.5; ref. 38). For continuous data that failed normality testing, intergroup comparisons were made by Kruskal-Wallis ANOVA followed by Mann-Whitney-Wilcoxon pairwise comparison. Statistical analyses were performed using GraphPad Prism (version8) and *R* computing software (*R* version 3.5.3 and RStudio Version 1.1.463). Pearson correlation coefficients were applied to compute correlations among colon and stool samples for θ_{YC} metric. A value of <0.05 was considered significant for both *P* and *q* values for all the analyses. Due to the small cohort size, analyses were not adjusted to any baseline sociodemographic or clinical characteristics or dietary calcium intake.

Results

Participant characteristics

Thirty-six participants were randomly enrolled. Six subjects were lost to follow-up prior to intervention, and a total of 30 participants (10 per arm) completed the study (Fig. 1). This included of 22 female and 8 male participants. Participants were randomized to study arms without regard to age or gender (placebo: 4 males and 6 females; calcium: 2 males and 8 females; Aquamin: 2 males and 8 females). Ages ranged from 20 to 66 years. Compliance (capsule intake) was estimated to be 96% across the three groups. Demographic characteristics are presented in Supplementary Table S2. Based on responses to DHQ II, the average dietary calcium intake (mg/day) values for the three groups at baseline were estimated to be: placebo = 817 ± 245 ; calcium = 964 ± 412 ; and Aquamin = 919 ± 545 (no significant differences found).

Safety and tolerability

Self-reported adverse events over the course of study are shown in Table 1. All adverse events were minor (i.e., headache, gastrointestinal symptoms) and did not preclude any individual from completing the study. The number of individuals

Table 1. Adverse events reported by the study subjects.

Events	Placebo	Calcium	Aquamin
Number of subjects participated	10	10	10
Number of subjects reported events	3	6	3
Total number of adverse events	5	15	7
Upper respiratory events (Flu-like symptoms)	1	1	0
Flu with fever (upper respiratory)	1	0	0
Skin rash	0	0	1
Headache	1	1	0
Gastrointestinal events	2	13	6
Nausea	0	1	0
Constipation	1	1	3
Diarrhea	0	4	1
Flatulence (& bloating)	1	2	0
Borborygmi	0	1	0
Belching	0	1	0
Abdominal pain	0	0	2
Blood in stool	0	3 ^a	0

Note: Adverse events (health issues) reported by subjects over the course of the study. No significant differences were among three groups.

^aOne subject reported three separate incidents of blood in stool over the period of 90 days.

reporting adverse events in the Aquamin group was the same as in the placebo group (3 of 10), whereas 6 of 10 individuals in the calcium group reported one or more events. No serious adverse events (defined as necessitating cessation of study participation or medical intervention/hospitalization) occurred. Serum metabolic markers (liver function/liver injury and renal function) are presented in **Table 2**. No significant change in any individual marker was observed over the course of the study. Likewise, there was no difference among the treatment groups in a panel of metabolic markers and no change in serum calcium levels before and after intervention (**Table 2**).

Table 2. Serum markers of safety and tolerability (Serum metabolic panel).

Group	Total protein (6.0–8.3 g/dl)	Albumin (3.5–4.9 g/dl)	AST (8–30 IU/L)	ALT (≤35 IU/L)	ALKP (30–116 IU/L)	Bilirubin (0.2–1.2 mg/dl)	Glucose (<100 mg/dl)
Placebo pre	7.2 ± 0.4	4.4 ± 0.2	25.6 ± 3.7	28.7 ± 8.3	75.8 ± 9.0	0.46 ± 0.2	91.4 ± 5.3
Placebo post	7.3 ± 0.3	4.5 ± 0.2	26.3 ± 5.5	31.9 ± 7.1	77.9 ± 15.3	0.54 ± 0.2	91.7 ± 6.4
Calcium pre	7.1 ± 0.7	4.4 ± 0.4	28.1 ± 10.1	25.9 ± 12.3	78.1 ± 20.5	0.59 ± 0.6	114.2 ± 58.7
Calcium post	7.2 ± 0.6	4.5 ± 0.4	28.7 ± 9.6	28.8 ± 11.5	80.5 ± 21.0	0.60 ± 0.2	102.0 ± 28.7
Aquamin pre	7.1 ± 0.4	4.5 ± 0.1	24.8 ± 6.8	23.4 ± 11.2	73.7 ± 18.4	0.57 ± 0.2	97.0 ± 22.3
Aquamin post	7.3 ± 0.4	4.6 ± 0.2	30.9 ± 10.9	32.5 ± 19.6	74.4 ± 19.7	0.61 ± 0.4	98.3 ± 22.7
Group	BUN (10–20 mg/dl)	Creatinine (0.6–1.2 mg/dl)	Carbon dioxide (23–30 mmol/L)	Sodium (136–145 mmol/L)	Potassium (3.5–5.2 mmol/L)	Chloride (96–106 mmol/L)	Calcium (8.6–10.3 mg/dl)
Placebo pre	16.9 ± 2.6	0.8 ± 0.1	28.4 ± 1.8	139.9 ± 2.0	4.4 ± 0.3	104.5 ± 1.4	9.5 ± 0.2
Placebo post	17.2 ± 4.5	0.8 ± 0.2	28.5 ± 3.0	140.1 ± 2.1	4.3 ± 0.3	105.2 ± 3.2	9.7 ± 0.3
Calcium pre	14.6 ± 5.1	0.8 ± 0.1	28.2 ± 3.5	140.9 ± 2.0	4.3 ± 0.6	105.8 ± 2.2	9.7 ± 0.5
Calcium post	14.0 ± 5.2	0.8 ± 0.1	28.6 ± 3.7	140.2 ± 1.5	4.2 ± 0.6	104.5 ± 3.5	9.8 ± 0.7
Aquamin pre	12.8 ± 2.8	0.7 ± 0.2	28.1 ± 2.7	140.5 ± 2.4	4.5 ± 0.3	104.6 ± 2.6	9.6 ± 0.4
Aquamin post	12.1 ± 4.1	0.7 ± 0.2	27.8 ± 1.9	140.5 ± 3.5	4.4 ± 0.3	105.2 ± 3.1	9.8 ± 0.4

Note: A comprehensive metabolic panel was done on participant's serum before and after the intervention.

No significant differences found among groups and within a group from the baseline.

Abbreviations: AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; ALKP: Alkaline phosphatase; BUN: Blood Urea Nitrogen.

Effects on bacterial DNA and microbial communities in colon biopsy and stool specimens

qPCR was used to assess bacterial 16S rRNA gene levels in colon biopsy and stool specimens as a way to estimate total bacterial DNA. **Figure 2** demonstrates that in both colon biopsy and stool specimens, posttreatment Aquamin samples had higher cycle quantification (Cq) values, indicating lower amounts of total bacterial DNA than at baseline (Colon: $P = 0.0001/q < 0.0001$; Stool: $P < 0.0001/q < 0.0001$). There was also a decrease in total bacterial DNA in specimens from calcium-treated participants (Colon: $P = 0.032/q = 0.0021$; Stool: $P < 0.0001/q < 0.0001$), although, on average, the decrease was not as great as that seen with Aquamin. Placebo specimens showed no (average) decline in DNA content at endpoint.

Alterations in gut microbial communities

To determine if Aquamin altered the composition of the gut microbiota in the participants, we analyzed bacterial 16S rRNA gene sequences from colon and stool specimens to explore pre-post-intervention differences. After sequence processing and exclusion of 2 samples for low sequence counts (i.e., one placebo stool and one calcium stool), a total of 7,081,672 sequences from 118 samples [median: 59,233 ± 23,618 (SD) sequences/sample; range 2,338–11,4235 sequences/sample] were included in this analysis. When we analyzed the sequence data by samples types for both colon biopsy and stool specimens, there was a total of 3,051,764 sequences from 60 colon samples [median: 49,853 ± 17,198 (SD) sequences/sample; range 12,481–102,310 sequences/sample]. The total number of sequences from 58 stool samples was 4,029,908 [median: 76,797 ± 25,547 (SD) sequences/sample; range: 2,338–11,4235 sequences/sample].

Shifts in gut microbial community composition were assessed by calculating θ_{YC} distances between the pre- and




Figure 2. Decrease in bacterial DNA with Aquamin. qPCR for bacterial DNA in colon (A) and stool (B) samples. The average cycle quantification (Cq) value from duplicate qPCR runs was plotted except for one post-Aquamin colon sample which failed to reach the amplification threshold in one duplicate and was therefore represented by a single Cq value (34.4) rather than the average. Another post-Aquamin colon sample failed to reach the amplification threshold in both duplicates and was, therefore, not plotted or included in the statistical analysis. A decrease in the total bacterial DNA is indicated by a higher Cq value (longer time to reach amplification threshold). There was no difference in pre- vs. postsupplementation for the placebo group. *, $P < 0.05$; ***, $P < 0.001$; and ****, $P < 0.0001$ compared with the corresponding pretreatment value.

postsupplementation communities (beta diversity) for each individual. Individual pre- postdistance values, grouped by intervention, are shown in **Fig. 3A** and **B**. Colonic microbial communities demonstrated a bigger within-participant shift in the Aquamin-supplemented group than was observed for either the placebo or calcium-supplemented group. The difference between Aquamin and calcium in the colon biopsy specimens reached the level of statistical significance ($P = 0.0087/q = 0.0061$). Both figure insets show that the majority of participants that received Aquamin (8 of 10 in colon and 7 of 10 in stool) were above the median value for within-participant

bacterial community change from pre- to postsupplementation for all samples. In colon biopsies, when samples were pooled as shown in the **Fig. 3A** inset, θ_{YC} distance was significantly increased in Aquamin subjects ($n = 10$) as compared with both calcium and placebo subjects ($n = 20$; $P = 0.019$). **Figure 3C** demonstrates a strong correlation between the two specimen types (colon biopsy and stool) for within-participant microbial community shifts (θ_{YC} distances; $P = 0.0041$). The correlation data, thus, suggest that stool specimens could be used as an estimate of the bacterial community in the colonic mucosa. It should be noted, however, that the stool specimens used here




Figure 3. θ_{YC} distances for comparison of pre- vs. postsupplementation values in gut bacterial community composition for colon (A) and stool (B) samples. Higher values reflect greater differences. **, $P = 0.0087$ relative to calcium. The insets show that the majority of the Aquamin (colon and stool) samples were above the median value for all samples. The majority of placebo and calcium samples were at or below the combined median value. **C**, Correlation between θ_{YC} in colon biopsy and stool specimens based on pre- vs. post-supplementation values ($P = 0.0041$).

Downloaded from <http://aacrjournals.org/cancerpreventionresearch/article-pdf/13/1/101/2245322/101.pdf> by guest on 26 January 2025

were obtained during flexible sigmoidoscopy and were not shed specimens. In addition, as part of this analysis, intrasubject versus intersubject variability was determined. θ_{YC} distances between the pre- and postintervention samples for each of the 30 individuals were compared with the θ_{YC} distances between each preintervention sample and all other samples. As seen in Supplementary Fig. S1, intrasubject differences were smaller than intersubject differences in both colon and stool samples ($P \leq 0.0001$).

Figure 4A–D presents PCoA of θ_{YC} distances between samples for all three treatment groups in both colon and stool specimens. At baseline, there were no significant differences in the colon bacterial communities (**Fig. 4A**), but following the 90-day intervention period, microbial communities in the colon biopsies from participants treated with Aquamin segregated from those in the placebo or calcium groups (Aquamin vs. placebo: AMOVA $P = 0.005$ and Aquamin vs. calcium: AMOVA $P = 0.009$; **Fig. 4A**). Further, colonic microbial communities from the posttreatment Aquamin biopsy samples segregated from their own baseline communities (AMOVA, $P = 0.022$; **Fig. 4B**), whereas there was no difference in pre- versus postsupplementation communities for placebo- or calcium-treated participants (**Fig. 4A**). In contrast to these results in colon biopsy material, stool sample microbial communities were not different between or within groups either at baseline or at endpoint (**Fig. 4C and D**).

Alpha diversity was assessed in both colon and stool samples using the Shannon diversity index. In Aquamin-treated participants, there was a decrease in Shannon diversity (**Fig. 4E and F**). For colon biopsy samples, Shannon diversity was significantly decreased ($P = 0.0037/q = 0.0026$) as compared with the placebo.

Alterations in the relative abundance of major gut phyla

First, we determined the total number of phyla and genera identified in each sample type. In colon biopsies, there were 15 different phyla and 224 genera identified. In stool samples, there were 10 phyla and 213 genera found. Overall, there were 7,267 OTUs detected; out of those, 2,573 were present in colon biopsies, 6,642 were present in stool samples, and 1,948 were common in both. A list containing all the phyla and genera identified are presented in Supplementary Tables S3 and S4.

Next, we assessed alterations in the relative abundance of individual OTUs representing the major gut phyla to compare intergroup differences in each of these phyla at endpoint. Bacteroidetes, Firmicutes, and Verrucomicrobia were decreased with Aquamin, whereas Actinobacteria and Proteobacteria were increased (with $P \leq 0.0001/q \leq 0.0001$). Trends were similar in both colon and stool specimens (**Fig. 4G and H**). Firmicutes OTUs also showed a drop with calcium intervention (colon biopsies only; $P < 0.0001/q < 0.0001$), but there was little change in the other phyla. Little change in OTU relative abundance in any of the major phyla was seen with placebo.

Identification of OTUs that differed with Aquamin supplementation

Differentially abundant OTUs were identified by LEfSe to explain the pre- and postsupplementation differences with Aquamin treatment. In the Aquamin-supplemented group, 82 OTUs were identified in colon samples (**Tables 3A and 3B**), and only 9 OTUs were identified in stool samples (**Tables 3C and 3D**) with an LDA score >2 and significance $P < 0.05$ ($q < 0.05$). Overall, more OTUs decreased in relative abundance than increased after supplementation. Differences between pre- and post-Aquamin supplementation were driven principally by an increase in OTUs within the normally less abundant phyla Proteobacteria, and Actinobacteria, along with three OTUs of phylum Firmicutes (**Table 3A**) and a decrease in OTUs within the normally higher abundance phyla Firmicutes and Bacteroidetes (**Table 3B**). Similar trends were also observed in stool samples (**Tables 3C and 3D**).

Effects on bile acid concentrations

Alterations in bile acid levels were assessed in stool samples from Aquamin- and calcium-treated participants in comparison with placebo as shown in **Fig. 5**. Aquamin treatment resulted in a net decline in total bile acids ($P = 0.0375/q = 0.0262$) with the major portion of this change being in unconjugated forms (the major components of the fecal bile acid pool; **Fig. 5A**). The major unconjugated primary bile acids—i.e., cholic acid (CA) and, particularly, chenodeoxycholic acid [CDCA; measured concurrently with deoxycholic acid (DCA)]—were significantly lower in posttreatment Aquamin stool samples ($P = 0.0074/q = 0.0052$ and $P = 0.0310/q = 0.0217$ respectively; **Fig. 5B**). This was not observed with calcium.

Of the measurable secondary bile acids, taurine-conjugated ursodeoxycholic acid (TUDCA), measured concurrently with taurohyodeoxycholic acid (THDCA), was decreased ($P = 0.0287/q = 0.0161$), as were the α and ω muricholic acids (minor secondary bile acid components; $P = 0.0012/q = 0.0009$ and $P = 0.0003/q < 0.0001$ respectively; **Fig. 5C**, inset). Two other unconjugated secondary bile acids [hyocholic acid (HCA) and hyodeoxycholic acid (HDCA)] were significantly decreased with Aquamin as compared with placebo ($P = 0.015/q = 0.010$ and $P < 0.0001/q < 0.0001$ respectively; **Fig. 5C**). HDCA is a byproduct of gut microbial metabolism (39), utilizing HCA and muricholic acids after microbial deconjugation and enzymatic modification. Overall, these Aquamin-associated bile acid changes in stool are consistent with both a decreased bile acid pool and decreased bacterial conversion of primary to secondary bile acids. Of interest, these effects were unique to Aquamin. Calcium supplementation did not show a measurable effect on total fecal bile acids, and only HDCA was significantly decreased (**Fig. 5C**).

Bile acid levels were also assessed in colon biopsy samples from Aquamin- and calcium-treated groups in comparison




Figure 4.

Segregation of gut microbial communities and gut microbial diversity. Biplot figures depicting PCoAs of θ_{VC} distances between colon biopsy and stool samples based on Illumina sequencing of the V4 region of 16S rRNA. θ_{VC} distance is a measure of difference in pre- and posttreatment microbial populations from individual participants with each of the three interventions. Data from colon specimens are shown in **A** and **B**, whereas stool specimen data are presented in **C** and **D**. Some of the OTUs driving the observed segregation between the groups are shown in the biplots. **A**, Differences in colon microbiota between postintervention Aquamin compared with postintervention placebo and calcium were seen: Aquamin samples with AMOVA *P* values of 0.005 (compared with placebo) and 0.009 (compared with calcium). **B**, Postintervention colon samples from Aquamin group were also significantly different relative to preintervention Aquamin samples with an AMOVA *P* value of 0.022. **C** and **D**, There were no significant differences found in stool samples. **E** and **F**, Shannon diversity index. Aquamin intervention reduced gut microbial (alpha) diversity as compared with the placebo in colon biopsy samples (*, *P* < 0.05). No significant change was observed in the gut microbial diversity in stool samples. **G** and **H**, Alterations in the relative abundance of major gut phyla with Aquamin supplementation. The change in the relative abundances of the top 1,000 OTUs (pooled by phyla) and assessed by pre- and postintervention analysis among three interventions in colon and stool specimens. For this analysis, 43 to 48 OTUs across the three treatment groups were pooled for Actinobacteria, 101 to 110 OTUs for Bacteroidetes, 543 to 591 OTUs for Firmicutes, 31 to 39 OTUs for Proteobacteria, and 4 to 6 OTUs for Verrucomicrobia phyla. **, *P* < 0.01; ***, *P* < 0.001; and ****, *P* < 0.0001.

Table 3A. LefSe data for OTU with elevated relative abundance in colon post-Aquamin.^a

OTU	Taxonomic ID	Order (family)	Phylum (class)	LDA score ^b	P value	q value
Otu0034	<i>Ralstonia</i>	Burkholderiales (<i>Burkholderiaceae</i>)	Proteobacteria (Betaproteobacteria)	4.68	0.001	0.024
Otu0052	<i>Comamonadaceae (uc)</i>	Burkholderiales (<i>Comamonadaceae</i>)	Proteobacteria (Betaproteobacteria)	4.48	0.009	0.027
Otu0092	<i>Leifsonia</i>	Actinomycetales (<i>Microbacteriaceae</i>)	Actinobacteria (Actinobacteria)	4.07	0.002	0.024
Otu0082	<i>Stenotrophomonas</i>	Xanthomonadales (<i>Xanthomonadaceae</i>)	Proteobacteria (Gammaproteobacteria)	4.03	0.002	0.024
Otu0095	<i>Rhodococcus</i>	Actinomycetales (<i>Nocardiaceae</i>)	Actinobacteria (Actinobacteria)	3.98	0.013	0.027
Otu0028	<i>Dorea</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.49	0.027	0.040
Otu0067	<i>Lachnospiraceae (uc)</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.36	0.035	0.044
Otu0157	<i>Flavonifractor</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	3.01	0.006	0.027

^aCriteria for inclusion: LDA values greater than 2 with $p/q < 0.05$.

^bLDA values reported as absolute value. OTU: Operational taxonomic unit. uc: unclassified.

Table 3B. LefSe data for OTU with decreased relative abundance in colon post-Aquamin^a.

OTU	Taxonomic ID	Order (family)	Phylum (class)	LDA score ^b	P value	q value
Otu0001	<i>Bacteroides</i>	Bacteroidales (<i>Bacteroidaceae</i>)	Bacteroidetes (Bacteroidia)	4.40	0.016	0.030
Otu0004	<i>Blautia</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	4.15	0.023	0.039
Otu0008	<i>Fusicatenibacter</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	4.00	0.005	0.027
Otu0009	<i>Anaerostipes</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.92	0.032	0.043
Otu0031	<i>Akkermansia</i>	Verrucomicrobiales (<i>Verrucomicrobiaceae</i>)	Verrucomicrobia (Verrucomicrobiae)	3.88	0.021	0.036
Otu0026	<i>Ruminococcus</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.83	0.008	0.027
Otu0016	<i>Parabacteroides</i>	Bacteroidales (<i>Porphyromonadaceae</i>)	Bacteroidetes (Bacteroidia)	3.81	0.013	0.027
Otu0043	<i>Lachnospiraceae (uc)</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.80	0.026	0.040
Otu0029	<i>Bacteroides</i>	Bacteroidales (<i>Bacteroidaceae</i>)	Bacteroidetes (Bacteroidia)	3.68	0.006	0.027
Otu0020	<i>Bacteroides</i>	Bacteroidales (<i>Bacteroidaceae</i>)	Bacteroidetes (Bacteroidia)	3.63	0.029	0.041
Otu0018	<i>Clostridiales (uc)</i>	Clostridiales (<i>Clostridiales_uc</i>)	Firmicutes (Clostridia)	3.60	0.024	0.039
Otu0076	<i>Lachnospiraceae (uc)</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.52	0.003	0.026
Otu0019	<i>Blautia</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.51	0.004	0.027
Otu0010	<i>Collinsella</i>	Coriobacteriales (<i>Coriobacteriaceae</i>)	Actinobacteria (Actinobacteria)	3.33	0.048	0.048
Otu0070	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.29	0.001	0.024
Otu0045	<i>Coprococcus</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.27	0.009	0.027
Otu0025	<i>Streptococcus</i>	Lactobacillales (<i>Streptococcaceae</i>)	Firmicutes (Bacilli)	3.18	0.018	0.035
Otu0110	<i>Parasutterella</i>	Burkholderiales (<i>Sutterellaceae</i>)	Proteobacteria (Betaproteobacteria)	3.16	0.021	0.036
Otu0054	<i>Ruminococcus2</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.15	0.031	0.043
Otu0107	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.14	0.002	0.026
Otu0087	<i>Bacteroides</i>	Bacteroidales (<i>Bacteroidaceae</i>)	Bacteroidetes (Bacteroidia)	3.08	0.019	0.035
Otu0037	<i>Dialister</i>	Selenomonadales (<i>Veillonellaceae</i>)	Firmicutes (Negativicutes)	3.08	0.019	0.035
Otu0069	<i>Coprococcus</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.07	0.045	0.048
Otu0074	<i>Bacteroides</i>	Bacteroidales (<i>Bacteroidaceae</i>)	Bacteroidetes (Bacteroidia)	3.06	0.007	0.027
Otu0068	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.03	0.009	0.027
Otu0091	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	3.03	0.027	0.040
Otu0128	<i>Clostridiales_uc</i>	Clostridiales (<i>Clostridiales</i>)	Firmicutes (Clostridia)	3.01	0.004	0.027
Otu0066	<i>Alistipes</i>	Bacteroidales (<i>Rikenellaceae</i>)	Bacteroidetes (Bacteroidia)	3.00	0.048	0.048
Otu0084	<i>Dorea</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.98	0.048	0.048
Otu0145	<i>Clostridium_IV</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.93	0.012	0.027
Otu0089	<i>Oscillibacter</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.90	0.007	0.027
Otu0148	<i>Clostridiales_uc</i>	Clostridiales (<i>Clostridiales_uc</i>)	Firmicutes (Clostridia)	2.90	0.036	0.044
Otu0113	<i>Clostridium_XIVa(98)</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.85	0.013	0.027
Otu0080	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.85	0.021	0.036
Otu0132	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.81	0.006	0.027
Otu0160	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.80	0.026	0.040
Otu0134	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.80	0.041	0.047
Otu0228	<i>Proteobacteria_uc</i>	Proteobacteria_uc (<i>Proteobacteria_uc</i>)	Proteobacteria_uc (Proteobacteria_uc)	2.78	0.013	0.027
Otu0136	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.76	0.002	0.024
Otu0265	<i>Clostridium_XIVa</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.71	0.029	0.041
Otu0090	<i>Alistipes</i>	Bacteroidales (<i>Rikenellaceae</i>)	Bacteroidetes (Bacteroidia)	2.70	0.026	0.040
Otu0198	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.65	0.015	0.030

(Continued on the following page)

Table 3B. LefSe data for OTU with decreased relative abundance in colon post-Aquamin^a. (Cont'd)

OTU	Taxonomic ID	Order (family)	Phylum (class)	LDA score ^b	P value	q value
Otu0215	<i>Faecalicoccus</i>	Erysipelotrichales (<i>Erysipelotrichaceae</i>)	Firmicutes (Erysipelotrichia)	2.64	0.010	0.027
Otu0227	<i>Clostridium_XIVa</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.64	0.025	0.040
Otu0101	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.63	0.036	0.044
Otu0192	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.61	0.009	0.027
Otu0276	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.61	0.006	0.027
Otu0138	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.56	0.001	0.024
Otu0149	<i>Clostridium_IV</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.55	0.001	0.024
Otu0088	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.55	0.021	0.036
Otu0156	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.53	0.039	0.045
Otu0120	<i>Lactococcus</i>	Lactobacillales (<i>Streptococcaceae</i>)	Firmicutes (Bacilli)	2.52	0.010	0.027
Otu0190	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.50	0.013	0.027
Otu0179	<i>Blautia</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.49	0.039	0.045
Otu0187	<i>Clostridium_XIVb</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.49	0.036	0.044
Otu0141	<i>Blautia</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.42	0.045	0.048
Otu0102	<i>Streptococcus</i>	Lactobacillales (<i>Streptococcaceae</i>)	Firmicutes (Bacilli)	2.41	0.039	0.045
Otu0278	<i>Peptoniphilus</i>	Clostridiales (<i>Peptoniphilaceae</i>)	Firmicutes (Clostridia)	2.36	0.045	0.048
Otu0194	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.31	0.031	0.043
Otu0164	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.29	0.015	0.030
Otu0201	<i>Clostridiales_uc</i>	Clostridiales (<i>Clostridiales_uc</i>)	Firmicutes (Clostridia)	2.28	0.013	0.027
Otu0218	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.25	0.039	0.045
Otu0165	<i>Eggerthella</i>	Coriobacteriales (<i>Coriobacteriaceae</i>)	Actinobacteria (Actinobacteria)	2.24	0.009	0.027
Otu0298	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.21	0.041	0.046
Otu0195	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.19	0.013	0.027
Otu0207	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.19	0.033	0.044
Otu0307	<i>Clostridiales_uc</i>	Clostridiales (<i>Clostridiales_uc</i>)	Firmicutes (Clostridia)	2.19	0.007	0.027
Otu0301	<i>Coriobacteriaceae_uc</i>	Coriobacteriales (<i>Coriobacteriaceae</i>)	Actinobacteria (Actinobacteria)	2.13	0.013	0.027
Otu0185	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.08	0.036	0.044
Otu0224	<i>Clostridium_XVIII</i>	Erysipelotrichales (<i>Erysipelotrichaceae</i>)	Firmicutes (Erysipelotrichia)	2.04	0.013	0.027
Otu0452	<i>Oscillibacter</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.02	0.013	0.027
Otu0341	<i>Holdemania</i>	Erysipelotrichales (<i>Erysipelotrichaceae</i>)	Firmicutes (Erysipelotrichia)	2.02	0.009	0.027
Otu0396	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.02	0.009	0.027
Otu0772	<i>Ruminococcaceae_uc</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.02	0.013	0.027

^aCriteria for inclusion: LDA values greater than 2 with $p/q < 0.05$.

^bLDA values reported as absolute value. OTU: Operational taxonomic unit. uc: unclassified.

Table 3C. LefSe data for OTU with elevated relative abundance in stool post-Aquamin^a.

OTU	Taxonomic ID	Order (family)	Phylum (class)	LDA score ^b	P value	q value
Otu0034	<i>Ralstonia</i>	Burkholderiales (<i>Burkholderiaceae</i>)	Proteobacteria (Betaproteobacteria)	3.15	0.002	0.010
Otu0194	<i>Lachnospiraceae_uc</i>	Clostridiales (<i>Lachnospiraceae</i>)	Firmicutes (Clostridia)	2.75	0.043	0.050
Otu0052	<i>Comamonadaceae (uc)</i>	Burkholderiales (<i>Comamonadaceae</i>)	Proteobacteria (Betaproteobacteria)	2.48	0.002	0.010

^aCriteria for inclusion: LDA values greater than 2 with $p/q < 0.05$.

^bLDA values reported as absolute value. OTU: Operational taxonomic unit. uc: unclassified.

Table 3D. LefSe data for OTU with decreased relative abundance in stool post-Aquamin^a.

OTU	Taxonomic ID	Order (family)	Phylum (class)	LDA score ^b	P value	q value
Otu0149	<i>Clostridium_IV</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	3.02	0.015	0.022
Otu0120	<i>Lactococcus</i>	Lactobacillales (<i>Streptococcaceae</i>)	Firmicutes (Bacilli)	2.73	0.014	0.022
Otu0272	<i>Bacteroides</i>	Bacteroidales (<i>Bacteroidaceae</i>)	Bacteroidetes (Bacteroidia)	2.46	0.014	0.022
Otu0234	<i>Actinomyces</i>	Actinomycetales (<i>Actinomycetaceae</i>)	Actinobacteria (Actinobacteria)	2.43	0.049	0.050
Otu0301	<i>Coriobacteriaceae_uc</i>	Coriobacteriales (<i>Coriobacteriaceae</i>)	Actinobacteria (Actinobacteria)	2.36	0.050	0.050
Otu0354	<i>Faecalibacterium</i>	Clostridiales (<i>Ruminococcaceae</i>)	Firmicutes (Clostridia)	2.08	0.010	0.022

^aCriteria for inclusion: LDA values greater than 2 with $p/q < 0.05$.

^bLDA values reported as absolute value. OTU: Operational taxonomic unit. uc: unclassified.




Figure 5.

Decrease in bile acids and increase in SCFAs in stool specimens. Bile acids and SCFA were assessed as described in the Materials and Methods Section. Values shown represent concentration differences between preintervention and postintervention samples. Asterisks represent statistical significance. **A**, Total bile acids (sum of the total conjugated and total unconjugated bile acids) are shown along with conjugated and unconjugated forms. * reflects decrease with Aquamin at $P = 0.0375$ (total) and at $P = 0.0527$ (unconjugated) versus calcium. **B**, Primary bile acids. CA and CDCA measured concurrently with DCA were significantly decreased with Aquamin at $P = 0.0074$ (CA) and $P = 0.0310$ (DCA/CDCA) versus calcium. **C**, Secondary bile acids. HDCA was significantly decreased with Aquamin and calcium versus placebo with P value < 0.0001 and $= 0.0149$ respectively. HCA and TUDCA, measured concurrently with THDCA were decreased with Aquamin relative to calcium, whereas α and ω muricholic acids were also reduced relative to calcium. For HCA, $P = 0.015$; for TUDCA/THDCA, $P = 0.0287$; for α MCA, $P = 0.0012$; and for ω MCA, $P = 0.0003$. With calcium supplementation, HDCA was also decreased relative to placebo ($P = 0.0149$). Inset: TUDCA/THDCA. **D**, SCFA. Acetate was significantly increased with Aquamin relative to calcium alone ($P < 0.0001$).

with placebo (Supplementary Fig. S2). Similar trends to those seen in stool specimens were observed, but none of the changes with Aquamin or calcium intervention reached statistical significance in comparison with placebo except HDCA which was significantly reduced with Aquamin ($P = 0.0456/q = 0.0319$) as compared with placebo.

Effects on SCFA concentrations

SCFA levels were assessed in both stool and colon samples. Apart from a modest (but statistically significant)

increase in acetate in stool samples from Aquamin-treated individuals relative to calcium ($P < 0.0001/q < 0.0001$), no other statistically significant alterations were observed in either stool (Fig. 5D) or colon (Supplementary Fig. S2) specimens.

As part of the analyses, we attempted to correlate bile acid changes and changes in SCFAs with alterations in microbial parameters. A weak correlation between total bile acid levels and θ YC was observed, but this was not statistically significant (due, in large part, to small sample size).

Downloaded from <http://aacrjournals.org/cancerpreventionresearch/article-pdf/13/1/101/2245322/101.pdf> by guest on 26 January 2025

Effects on colonic eicosanoid concentrations

There was no significant change from baseline detected for any eicosanoid apart from an increase in 13S-hydroxy-octadecadienoic acid (13S-HODE) in calcium-supplemented colon samples (Supplementary Fig. S3).

Discussion

This study demonstrated that 90-day dietary intervention with Aquamin was well-tolerated by healthy individuals. There were no changes in liver injury/function and kidney function markers; there were no serious adverse events; and minor adverse events were no higher with Aquamin than with calcium alone or placebo. These findings are consistent with the previous reports (25–27). Although a much larger database will be required to establish safety, all of the data to date suggest that safety and tolerability will not be issues with Aquamin. At the same time, intervention with Aquamin resulted in measurable changes in the colonic microbial community and the attendant bile acid profile. We are currently carrying out a phase I/II therapeutic trial with Aquamin in ulcerative colitis in remission. The same microbial and metabolomic endpoints assessed here are being measured in the ongoing trial. It is our hope that these microbial and metabolomic findings will prove useful as biomarkers going forward in the interventional trial.

The microbial population present in the colon of an individual is sensitive to environmental influences, including diet (40, 41). Past studies have demonstrated that calcium (18, 19) and Aquamin itself (20) can influence the colonic microbial community in mice. In the present study, we found that Aquamin supplementation in human participants resulted in a decrease in total colonic microbial DNA and an overall decrease in OTUs within the major gut phyla Firmicutes and Bacteroidetes, with a decrease in gut microbial diversity. Thus, the major impact may simply be a decrease in total gut bacteria. The impact of the decrease in bacteria on colonic health is unknown at this time but poses interesting questions for future exploration. Germ-free mice have shown increased resistance to chemical or dietary-induced colon cancers (42). Because our own previous studies have shown that Aquamin-treated mice have decreased incidence of colon polyps (10, 11) as well as fewer liver tumors (43), the possibility that Aquamin-associated microbial decreases may play a mechanistic role in these observations warrants further exploration. Of interest, many of the observations made with samples from Aquamin-treated participants were not seen in samples from individuals ingesting calcium alone. This argues that although calcium is the major mineral component in Aquamin, the effects of Aquamin on the gut microbial community cannot be attributed to calcium alone.

The changes we observed in the post-Aquamin treatment specimens are similar to the effects of broad-spectrum antibiotic treatment, which have been shown to cause a significant reduction in the proportions of Firmicutes and Bacteroidetes and an increase in Proteobacteria (44). As a follow-up to the

observations reported here, it would be informative to assess the potential antimicrobial activity of Aquamin supplementation directly in a minimum inhibitory concentration assay. Metals, particularly cations, have previously been utilized as antimicrobial agents (45). For example, colloidal or nanoparticulate silver and mineral-rich clays are known to have broad-spectrum antimicrobial effects (46, 47). Notably, in our study, the administration of Aquamin did not result in diarrhea and caused only a small drop in bacterial diversity in colon samples; both common adverse effects of broad-spectrum antibiotic use. Thus, the antimicrobial effects of Aquamin appear to be mild in comparison with antibiotics and may allow some of the benefit of reduced gut microbial populations (e.g., reduced potentially carcinogenic secondary bile acids) without inducing symptoms that would render the intervention intolerable.

In parallel with microbial changes, Aquamin administration decreased the levels of total bile acids and selected primary and secondary bile acids. This finding is consistent with the decrease in total bacteria and may be potentially beneficial to colon health. At high concentrations, bile acids are membrane-damaging and cytotoxic (21). In addition, several secondary bile acids, most notably lithocholic acid (LCA) and DCA (22, 23), are known to be carcinogenic. Although these species could not be individually measured in this study for methodological reasons, the combination of DCA and CDCA (the primary bile acid precursor of LCA) was significantly lower in the Aquamin-supplemented group (Fig. 5). This suggests that one beneficial effect of Aquamin supplementation might be through effects on bile acid metabolism. A caveat in the interpretation of bile acid data is that it is not currently known whether the reduction in gut bile acid pools is due to altered bacterial composition or, potentially, to a direct effect on the bile acids, themselves. Previous studies have shown that calcium, the major component of Aquamin, has the capacity to precipitate bile acids (48). Although mineral binding and precipitation might interfere with detection, this could be beneficial if it prevented potentially harmful secondary bile acids from binding to colonic epithelium or entering the circulation. Arguing against direct binding and precipitation is the finding that calcium alone did not mimic the effects of Aquamin on the total bile acid pool.

A decrease in gut bacteria could, potentially, be harmful to colon health if it resulted in decreased concentration of protective bacterial metabolites, such as the colon-protective SCFAs (24). However, in this study there was no decrease in the colonic concentration of butyrate in the Aquamin-treated group and an actual increase in acetate. Thus, even with the apparent decline in total bacteria, the size and/or composition of the microbial pool was sufficient to maintain presupplementation levels of certain SCFAs.

Although the data strongly suggest an Aquamin-associated decrease in gut microbial bacteria, the specific shifts in gut microbial populations are harder to interpret. One complication of low microbial biomass samples during microbial sequencing is greater likelihood of detecting reagent or

laboratory contaminant microbes in the amplification step. This has been documented in low-biomass samples such as glacier ice, air, rocks, etc. (49), but is not likely to be a major issue in the normal high biomass of the colon.

It should be noted that the antimicrobial effect of Aquamin might not be a direct consequence of the mineral components making up the supplement. Our recent studies employing colonoid culture technology demonstrated upregulation of proteins having antimicrobial activity upon treatment with Aquamin (50). These included lactotransferrin (51), natural resistance-associated macrophage protein-2 (52), and Ly6/PLAUR domain-containing protein 8 (53). Thus, it is possible that the antimicrobial effects seen with Aquamin *in vivo* are mediated, in part, through its effects on the colonic mucosa, itself. Finally, decreased bacteria, especially as it relates to colon biopsy specimens, might reflect altered bacterial adhesion. In our colonoid culture study (50), several mucins and CEA-CAMS were altered by Aquamin. Studies by others have noted that alterations in mucosal surface proteins affect microbial interactions with the mucosal wall (15, 16, 53).

In addition to the microbial and metabolomic changes, there are other, direct, beneficial effects of Aquamin on the epithelium of the colonic mucosa. Specifically, Aquamin suppressed colon epithelial proliferation and induced differentiation in colonoid culture. Effects on proliferation were seen, primarily, in colonoids derived from large adenomas (14), whereas improved differentiation—including upregulation of multiple cell–cell and cell–matrix adhesion molecules and barrier proteins—was observed in normal tissue-derived colonoids (50) as well as in adenoma colonoids (14). Direct effects on the colonic mucosa and indirect effects resulting from changes in microbial/metabolomic profiles are, of course, not mutually exclusive. Further, as noted above, the indirect effects (i.e., antimicrobial activity) may reflect changes induced in the colonic mucosa.

The primary limitations of this study were the small sample size and short duration, which were by design, as this was an initial tolerability study rather than an efficacy study. Although the study was too short and too small to provide definitive results with several endpoints, it was sufficiently powered to detect significant differences in several microbial and metabolomic (bile acid) endpoints in Aquamin-treated participants relative to the other two interventions. To better define effects on efficacy-related endpoints, we are currently beginning a 180-day interventional trial with Aquamin in participants with ulcerative colitis in remission. Another limitation was the potential for intrasubject (as well as intersubject) variability that needs to be acknowledged. Pre and postinterventional differences in the microbial community were observed in all subjects, but intrasubject variability was less than variability between individuals (Supplementary Fig. S1).

In summary, this pilot study demonstrated that 90-day dietary intervention with a calcium- and magnesium-rich multimineral supplement was well-tolerated in healthy human volunteers. Adverse events were mild and largely consisted of

gastrointestinal symptoms (constipation, diarrhea, etc.) that did not preclude completion of the study. Reports of adverse events were actually less frequent with Aquamin than with calcium alone. Consistent with this, liver function/liver injury and renal function markers and other serum biochemical markers did not vary significantly with intervention. With respect to microbial and metabolomic findings, Aquamin supplementation resulted in an overall decrease in gut microbial numbers and a decrease in bile acids, including potentially carcinogenic secondary bile acids or their precursors. Despite the antimicrobial-like effects, concentrations of the colon-protective SCFAs were maintained. These findings, along with previous *in vitro* and animal data (10–14, 50) showing a beneficial effect of Aquamin on colonic health, support future longer-term interventional studies in human participants. Finally, the observation that Aquamin had a more pronounced effect on gut microbial populations and bile acid levels than calcium alone supports the view that the beneficial activity of Aquamin (calcium in conjunction with additional trace elements) cannot be attributed to calcium alone. This conclusion supports findings from a recent epidemiologic study suggesting that calcium in combination of additional minerals may be linked with the lower risk of colorectal cancer (54).

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: M.N. Aslam, C.M. Bassis, S.M. Zick, D.K. Turgeon, J. Varani

Development of methodology: M.N. Aslam, C.M. Bassis, S.M. Zick, D.K. Turgeon, J. Varani

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): M.N. Aslam, C.M. Bassis, I.L. Bergin, K. Knuver, D.K. Turgeon, J. Varani

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): M.N. Aslam, C.M. Bassis, I.L. Bergin, K. Knuver, A. Sen, J. Varani

Writing, review, and/or revision of the manuscript: M.N. Aslam, C.M. Bassis, I.L. Bergin, K. Knuver, S.M. Zick, A. Sen, D.K. Turgeon, J. Varani

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): M.N. Aslam, C.M. Bassis, J. Varani

Study supervision: M.N. Aslam, D.K. Turgeon, J. Varani

Acknowledgments

This study was supported by NIH grant CA201782 including supplemental funding through the Office of Dietary Supplements to J. Varani. Metabolomic work was supported by a Pilot/Feasibility Grant from Michigan Nutrition and Obesity Research Center to I.L. Bergin and M.N. Aslam (through 2 P30 DK089503-06 to MNORC). This study also used services at the University of Michigan that were supported by NIH funding (UL1TR002240; KL2TR002241; and TL1TR002242) to the Michigan Institute for Clinical and Health Research (MICH). We thank MICH, the Michigan Clinical Research Unit, the University of Michigan Microbial Systems Molecular Biology Laboratory, and the RCMRC at the University of Michigan for help with various aspects of the study. We thank our study coordinators (Elaine Brady and Deepa Chandrasekhar) for their

assistance with the study. We also thank Marigot LTD. for providing Aquamin as a gift. Most importantly, we are very grateful to the study participants who made it possible for us to carry out this study.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked

advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received June 27, 2019; revised August 2, 2019; accepted November 22, 2019; published first November 26, 2019.

References

- Keum NN, Aune D, Greenwood DC, Ju W, Givannucci EL. Calcium intake and cancer risk: dose-response meta-analysis of prospective observational studies. *Int J Cancer* 2014;135:1940–8.
- Newmark HL, Yang K, Kurihara N, Fan K, Augenlicht LH, Lipkin M. Western-style diet-induced colonic tumors and their modulation by calcium and vitamin D in C57Bl/6 mice: a preclinical model for human sporadic colon cancer. *Carcinogenesis* 2009;1:88–92.
- Mariadason JM, Bordonaro M, Aslam F, Shi L, Kuraguchi M, Velcich A, et al. Down-regulation of beta-catenin TCF signaling is linked to colonic epithelial cell differentiation. *Cancer Res* 2001;61:3465–71.
- Baron JA, Beach M, Mandel JS, van Stolk RU, Haile RW, Sandler RS, et al. Calcium supplements and colorectal adenomas. Polyp Prevention Study Group. *Ann NY Acad Sci* 1999;889:138–45.
- Grau MV, Baron JA, Sandler RS, Haile RW, Beach ML, Church TR, et al. Vitamin D, calcium supplementation, and colorectal adenomas: results of a randomized trial. *J Natl Cancer Inst* 2003;95:1765–71.
- Baron JA, Barry EL, Mott LA, Rees JR, Sandler RS, Snover DC, et al. A trial of calcium and vitamin D for the prevention of colorectal adenomas. *N Engl J Med* 2015;373:1519–30.
- Pommergaard HC, Burcharth J, Rosenberg J, Raskov H. Aspirin, calcitriol, and calcium do not prevent adenoma recurrence in a randomized controlled trial. *Gastroenterology* 2016;150:114–22.
- Crockett SD, Barry EL, Mott LA, Ahnen DJ, Robertson DJ, Anderson JC, et al. Calcium and vitamin D supplementation and increased risk of serrated polyps: results from a randomised clinical trial. *Gut* 2018; pii: gutjnl-2017-315242.
- Bolland MJ, Avenell A, Baron JA, Grey A, MacLennan GS, Gamble GD, et al. Effect of calcium supplements on risk of myocardial infarction and cardiovascular events: meta-analysis. *BMJ* 2010;341: c3691.
- Aslam MN, Paruchuri T, Bhagavathula N, Varani J. A mineral-rich red algae extract inhibits polyp formation and inflammation in the gastrointestinal tract of mice on a high-fat diet. *Integr Cancer Ther* 2010;9:93–9.
- Aslam MN, Bergin I, Naik M, Paruchuri T, Hampton A, Rehman M, et al. A multiminerall natural product from red marine algae reduces colon polyp formation in C57BL/6 mice. *Nutr Cancer* 2012;64:1020–8.
- Aslam MN, Bhagavathula N, Paruchuri T, Hu X, Chakrabarty S, Varani J. Growth-inhibitory effects of a mineralized extract from the red marine algae, *Lithothamnion calcareum*, on Ca²⁺-sensitive and Ca²⁺-resistant human colon carcinoma cells. *Cancer Lett* 2009;283: 186–92.
- Singh N, Aslam MN, Varani J, Chakrabarty S. Induction of calcium sensing receptor in human colon cancer cells by calcium, vitamin D and aquamin: promotion of a more differentiated, less malignant and indolent phenotype. *Mol Carcinog* 2015;54:543–53.
- McClintock SD, Colacino JA, Attili D, Dame MK, Richter A, Reddy AR, et al. Calcium-induced differentiation of human colon adenomas in colonoid culture: calcium alone versus calcium with additional trace elements. *Cancer Prev Res (Phila)* 2018;11:413–28.
- Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. *Cell* 2014;157:121–41.
- Halfvarson J, Brislawn CJ, Lamendella R, Vázquez-Baeza Y, Walters WA, Bramer LM, et al. Dynamics of the human gut microbiome in inflammatory bowel disease. *Nat Microbiol* 2017;2:17004.
- Brennan CA, Garrett WS. Gut microbiota, inflammation, and colorectal cancer. *Annu Rev Microbiol* 2016;70:395–411.
- Chaplin A, Parra P, Laraichi S, Serra F, Palou A. Calcium supplementation modulates gut microbiota in a prebiotic manner in dietary obese mice. *Mol Nutr Food Res* 2016;60:468–80.
- Aslam MN, Bassis CM, Zhang L, Zaidi S, Varani J, Bergin IL. Calcium reduces liver injury in mice on a high-fat diet: alterations in microbial and bile acid profiles. *PLoS One* 2016;11:e0166178.
- Crowley EK, Long-Smith CM, Murphy A, Patterson E, Murphy K, O’Gorman DM, et al. Dietary supplementation with a magnesium-rich marine mineral blend enhances the diversity of gastrointestinal microbiota. *Mar Drugs* 2018;16:pii: E216.
- Mello-Vieira J, Sousa T, Coutinho A, Fedorov A, Lucas SD, Moreira R, et al. Cytotoxic bile acids, but not cytoprotective species, inhibit the ordering effect of cholesterol in model membranes at physiologically active concentrations. *Biochim Biophys Acta* 2013;1828:2152–63.
- Bernstein H, Bernstein C, Payne CM, Dvorakova K, Garewal H. Bile acids as carcinogens in human gastrointestinal cancers. *Mutat Res* 2005;589:47–65.
- Wahlström A, Sayin SI, Marschall HU, Bäckhed F. Intestinal crosstalk between bile acids and microbiota and its impact on host metabolism. *Cell Metab* 2016;24:41–50.
- Ríos-Covián D, Ruas-Madiedo P, Margolles A, Gueimonde M, de Los Reyes-Gavilán CG, Salazar N. Intestinal short chain fatty acids and their link with diet and human health. *Front Microbiol* 2016;7:185.
- Frestedt JL, Walsh M, Kuskowski MA, Zenk JL. A natural mineral supplement provides relief from knee osteoarthritis symptoms: a randomized controlled pilot trial. *Nutr J* 2008;7:9.
- Frestedt JL, Kuskowski MA, Zenk JL. A natural seaweed derived mineral supplement (Aquamin F) for knee osteoarthritis: a randomized, placebo controlled pilot study. *Nutr J* 2009;8:7.
- Slevin M, Allsopp PJ, Magee PJ, Bonham MP, Naughton VR, Strain JJ, et al. Supplementation with calcium and short-chain fructooligosaccharides affects markers of bone turnover but not bone mineral density in postmenopausal women. *J Nutr* 2014;144:297–304.
- National Cancer Institute. Diet History Questionnaire, Version 2.0. Epidemiology and Genomics Research Program. Bethesda, MD: National Cancer Institute; 2010. Available from: <https://epi.grants.cancer.gov/dhq2/>.
- Dollive S, Chen YY, Grunberg S, Bittinger K, Hoffmann C, Vandivier L, et al. Fungi of the murine gut: episodic variation and proliferation during antibiotic treatment. *PLoS One* 2013;8:e71806.
- Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* 2013;79:5112–20.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 2009;75:7537–41.

32. Westcott SL, Schloss PD. OptiClust, an improved method for assigning amplicon-based sequence data to operational taxonomic units. *MSphere* 2017;2:e00073–17.
33. Yue JC, Clayton MK. A similarity measure based on species proportions. *Commun Stat Theory Methods* 2005;34:11, 2123–31, DOI: 10.1080/STA-200066418.
34. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic biomarker discovery and explanation. *Genome Biol* 2011;12:R60.
35. Griffiths WJ, Sjövall J. Bile acids: analysis in biological fluids and tissues. *J Lipid Res* 2010;51:23–41.
36. Demehri FR, Frykman PK, Cheng Z, Ruan C, Wester T, Nordenskjöld A, et al. HAEC Collaborative Research Group. Altered fecal short chain fatty acid composition in children with a history of Hirschsprung-associated enterocolitis. *J Pediatr Surg* 2016;51:81–6.
37. Yang R, Chiang N, Oh SF, Serhan CN. Metabolomics-lipidomics of eicosanoids and docosanoids generated by phagocytes. *Curr Protoc Immunol* 2011;Chapter 14:Unit 14.26.
38. Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 2001;26:32–46.
39. Eysen HJ, De Pauw G, Van Eldere J. Formation of hydoxycholeic acid from muricholic acid and hyocholic acid by an unidentified gram-positive rod termed HDCA 1 isolated from rat intestinal microflora. *Appl Environ Microbiol* 1999;65:3158–63.
40. Louis P, Hold GL, Flint HJ. The gut microbiota, bacterial metabolites and colorectal cancer. *Nat Rev Microbiol* 2014;12:661–72.
41. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 2014;505:559–63.
42. Sobhani I, Amiot A, Le Baleur Y, Levy M, Auriault ML, Van Nhieu JT, et al. Microbial dysbiosis and colon carcinogenesis: could colon cancer be considered a bacteria related disease? *Therap Adv Gastroenterol* 2013;6:215–29.
43. Aslam MN, Bergin I, Naik M, Hampton A, Allen R, Kunkel SL, et al. A multi-mineral natural product inhibits liver tumor formation in C57BL/6 mice. *Biol Trace Elem Res* 2012;147:267–74.
44. Fujisaka S, Ussar S, Clish C, Devkota S, Dreyfuss JM, Sakaguchi M, et al. Antibiotic effects on gut microbiota and metabolism are host dependent. *J Clin Invest* 2016;126:4430–43.
45. Lemire JA, Harrison JJ, Turner RJ. Antimicrobial activity of metals: mechanisms, molecular targets and applications. *Nat Rev Microbiol* 2013;11:371–84.
46. Haydel SE, Remenih CM, Williams LB. Broad-spectrum *in vitro* antibacterial activities of clay minerals against antibiotic-susceptible and antibiotic-resistant bacterial pathogens. *J Antimicrob Chemother* 2008;61:353–61.
47. Morones-Ramirez JR, Winkler JA, Spina CS, Collins JJ. Silver enhances antibiotic activity against gram-negative bacteria. *Sci Transl Med* 2013;5:190ra81.
48. Van der Meer R, Welberg JW, Kuipers F, Kleibeuker JH, Mulder NH, Termont DS, et al. Effects of supplemental dietary calcium on the intestinal association of calcium, phosphate, and bile acids. *Gastroenterology*. 1990;99:1653–9.
49. Eisenhofer R, Minich JJ, Marotz C, Cooper A, Knight R, Weyrich LS. Contamination in low microbial biomass microbiome studies: issues and recommendations. *Trends Microbiol* 2019;27:105–17.
50. Attili D, McClintock SD, Rizvi AH, Pandya S, Rehman H, Nadeem DM, et al. Calcium-induced differentiation in normal human colonic cultures: Cell-cell/cell-matrix adhesion, barrier formation and tissue integrity. *PLoS One* 2019;14:e0215122.
51. Orsi N. The antimicrobial activity of lactoferrin: current status and perspectives. *Biometals* 2004;17:189–96.
52. Cellier MF, Courville P, Campion C. Nramp1 phagocyte intracellular metal withdrawal defense. *Microbes Infect* 2007;9:1662–70.
53. Okumura R, Kurakawa T, Nakano T, Kayama H, Kinoshita M, Motooka D, et al. Lypd8 promotes the segregation of flagellated microbiota and colonic epithelia. *Nature* 2016;532:117.
54. Swaminath S, Um CY, Prizment AE, Lazovich D, Bostick RM. Combined mineral intakes and risk of colorectal cancer in postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2019;28:392–9.

