

Contraction-Mediated Glucose Uptake: A Central Role for Rac1

Leonidas S. Lundell and Anna Krook

Skeletal muscle glucose uptake can be regulated in response to different stimuli, with insulin and contraction being central. While the processes underlying insulin-induced glucose uptake have been extensively characterized, the processes underlying contraction-induced glucose uptake remain elusive. In this issue of *Diabetes*, Sylow et al. (1) provide another piece to this puzzle. They report that Rac1 is essential for contraction-induced GLUT4 translocation.

Contraction induces glucose uptake in skeletal muscle both independently (2) and synergistically (3) with insulin. Several different mechanisms have been proposed, which appear to be activated in parallel following muscle contraction (Fig. 1). Key pathways include activation of the AMP-sensitive protein kinase (AMPK) through changes in the AMP:ATP ratio in response to increasing energy demands (4). Increased Ca^{2+} /calmodulin-dependent protein kinase I (CaMKI) influx during muscle depolarization results in activation of calcium signaling cascades via CaMKI (5) and protein kinase C (PKC) (6). Downstream candidates include the Rab-GTPase-activating protein TBC1D1 (7), and evidence supports a role of increased free radical content in contraction-mediated glucose uptake as a consequence of increased respiratory activity (8).

Rac1 is a small GTPase and member of the Rho family, which has previously been implicated in a wide range of processes including membrane dynamics, cell adhesion, and proliferation (9). Previous work has identified Rac1 as a downstream mediator of insulin-stimulated glucose transport in cultured skeletal muscle, where in vitro siRNA silencing leads to abrogated actin remodeling and prevents insulin-induced GLUT4 translocation (10). In the current study, Sylow et al. (1) provide evidence for a role for Rac1 in mediating contraction-induced GLUT4 membrane translocation, demonstrating that in vivo, Rac1 is activated in response to exercise in both human and mouse skeletal muscle. Consistent with a role for Rac1 in contraction-induced glucose uptake, ex vivo contraction of mouse muscle lacking Rac1 displayed impaired contraction-induced glucose transport. Furthermore, pharmacological inhibition of Rac1 in vitro reduced insulin-stimulated glucose uptake. Interestingly, the authors present two lines of evidence that contraction-mediated Rac1 activation is AMPK-independent. Mice with a targeted deletion of

AMPK kinase activity displayed a normal Rac1 activation. Additionally AICAR, a pharmacological AMPK activator, did not lead to increased Rac1 activation (1).

While providing new molecular insight into regulation of GLUT4 traffic, a number of key questions remain to be resolved. Since the authors exclude a role for AMPK as an upstream regulator of Rac1, what then are the remaining contenders for this role? Possible candidates include TBC1D1 with its calcium-binding motifs, as well as several PKC isoforms that are activated by an AMPK-independent mechanism (Fig. 1). Furthermore, Sylow et al. show that exercise leads to Rac1 phosphorylation and GTP loading. The molecular pathways regulating these events are imprecisely established and could take very different forms, for example kinase-mediated phosphorylation (11), GDP to GTP exchange by stimulation of guanine nucleotide exchange factor activation (12), decreased GTP hydrolysis by inhibition of GTPase activating proteins (13), or inhibition of GDP dissociation by guanosine nucleotide dissociation inhibitors (9). In addition, the effectors lying downstream of Rac1 involved in actin remodeling remain to be characterized, and future studies to address this mechanism are warranted (10). The direct effect of exercise in Rac1-depleted mice will also be interesting in further studies, both with respect to glucose uptake and in relationship to exercise performance and muscle ergonomics.

While evidence from rodent models of type 2 diabetes indicates that insulin-stimulated but not exercise-stimulated glucose transport is impaired in skeletal muscle from diabetic rodents (14), the data from type 2 diabetic patients is less clear. Indeed, AICAR-stimulated and hypoxia-stimulated glucose uptake is reduced in skeletal muscle from type 2 diabetic patients (4,15). Furthermore, upon a careful examination of the limited number of subjects that have been studied, contraction-induced GLUT4 translocation may also be reduced in skeletal muscle from type 2 diabetic patients (16). This has led to the proposal that insulin resistance in human skeletal muscle involves defective GLUT4 traffic and targeting (17). A broader challenge in dissecting the disease mechanism for skeletal muscle insulin resistance in type 2 diabetes is to understand whether Rac1 activation is altered in the context of human pathophysiology, and to determine whether Rac1 represents a convergence point linking insulin-induced and contraction-induced signals to the control of GLUT4 translocation. Answers to these questions will potentially increase the molecular understanding of contraction-induced glucose uptake and provide novel entry points for possible therapeutic interventions in type 2 diabetes.

ACKNOWLEDGMENTS

No potential conflicts of interest relevant to this article were reported.

From the Department of Physiology and Pharmacology, Section for Integrative Physiology, Karolinska Institutet, Stockholm, Sweden.

Corresponding author: Anna Krook, anna.krook@ki.se.

DOI: 10.2337/db12-1789

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

See accompanying original article, p. 1139.




FIG. 1. Possible upstream and direct regulators of Rac1 activity. Contraction stimulates GLUT4 exocytosis by activating a range of different molecules, including AMPK, PKC, and TBCD1. Evidence presented by Sylow et al. (1) indicates that AMPK is not involved in Rac1 activation. Other possible pathways include TBCD1 and PKC. Rac1 cycling between active, GTP bound state and inactive, GDP bound state is another possible node by which Rac1 regulates GLUT4 exocytosis.

REFERENCES

- Sylow L, Jensen TE, Kleinert M, et al. Rac1 is a novel regulator of contraction-stimulated glucose uptake in skeletal muscle. *Diabetes* 2013;62:1139–1151
- Wallberg-Henriksson H, Holloszy JO. Contractile activity increases glucose uptake by muscle in severely diabetic rats. *J Appl Physiol* 1984;57:1045–1049
- Wallberg-Henriksson H, Constable SH, Young DA, Holloszy JO. Glucose transport into rat skeletal muscle: interaction between exercise and insulin. *J Appl Physiol* 1988;65:909–913
- Koistinen HA, Galuska D, Chibalin AV, et al. 5-amino-imidazole carboxamide riboside increases glucose transport and cell-surface GLUT4 content in skeletal muscle from subjects with type 2 diabetes. *Diabetes* 2003;52:1066–1072
- Hawley SA, Selbert MA, Goldstein EG, Edelman AM, Carling D, Hardie DG. 5'-AMP activates the AMP-activated protein kinase cascade, and Ca²⁺/calmodulin activates the calmodulin-dependent protein kinase I cascade, via three independent mechanisms. *J Biol Chem* 1995;270:27186–27191
- Niu W, Bilan PJ, Yu J, et al. PKC ϵ regulates contraction-stimulated GLUT4 traffic in skeletal muscle cells. *J Cell Physiol* 2011;226:173–180
- Jessen N, An D, Lihn AS, et al. Exercise increases TBC1D1 phosphorylation in human skeletal muscle. *Am J Physiol Endocrinol Metab* 2011;301:E164–E171
- Merry TL, Steinberg GR, Lynch GS, McConell GK. Skeletal muscle glucose uptake during contraction is regulated by nitric oxide and ROS independently of AMPK. *Am J Physiol Endocrinol Metab* 2010;298:E577–E585
- Bustelo XR, Ojeda V, Barreira M, Sauzeau V, Castro-Castro A. Rac-ing to the plasma membrane: the long and complex work commute of Rac1 during cell signaling. *Small GTPases* 2012;3:60–66
- Chiu TT, Jensen TE, Sylow L, Richter EA, Klip A. Rac1 signalling towards GLUT4/glucose uptake in skeletal muscle. *Cell Signal* 2011;23:1546–1554
- Kwon T, Kwon DY, Chun J, Kim JH, Kang SS. Akt protein kinase inhibits Rac1-GTP binding through phosphorylation at serine 71 of Rac1. *J Biol Chem* 2000;275:423–428
- Ueda S, Kataoka T, Satoh T. Activation of the small GTPase Rac1 by a specific guanine-nucleotide-exchange factor suffices to induce glucose uptake into skeletal-muscle cells. *Biol Cell* 2008;100:645–657
- Tatsis N, Lannigan DA, Macara IG. The function of the p190 Rho GTPase-activating protein is controlled by its N-terminal GTP binding domain. *J Biol Chem* 1998;273:34631–34638
- Coderre L, Kandror KV, Vallega G, Pilch PF. Identification and characterization of an exercise-sensitive pool of glucose transporters in skeletal muscle. *J Biol Chem* 1995;270:27584–27588
- Ryder JW, Yang J, Galuska D, et al. Use of a novel impermeable biotinylated photolabeling reagent to assess insulin- and hypoxia-stimulated cell surface GLUT4 content in skeletal muscle from type 2 diabetic patients. *Diabetes* 2000;49:647–654
- Kennedy JW, Hirshman MF, Gervino EV, et al. Acute exercise induces GLUT4 translocation in skeletal muscle of normal human subjects and subjects with type 2 diabetes. *Diabetes* 1999;48:1192–1197
- Garvey WT, Maianu L, Zhu JH, Brechtel-Hook G, Wallace P, Baron AD. Evidence for defects in the trafficking and translocation of GLUT4 glucose transporters in skeletal muscle as a cause of human insulin resistance. *J Clin Invest* 1998;101:2377–2386