

# Effect of Intensive Diabetes Treatment on Carotid Artery Wall Thickness in the Epidemiology of Diabetes Interventions and Complications

Epidemiology of Diabetes Interventions and Complications (EDIC) Research Group

The Epidemiology of Diabetes Interventions and Complications (EDIC) is a multicenter longitudinal observational study of the Diabetes Control and Complications Trial (DCCT) cohort. One of the major objectives of EDIC is to study the development and progression of atherosclerotic cardiovascular disease in type 1 diabetes. In this study, we evaluated the role of cardiovascular risk factors and antecedent therapy in the DCCT on carotid intima-media wall thickness (IMT) in type 1 diabetes. At ~18 months after the end of the DCCT, high-resolution B-mode ultrasonography was used to assess the carotid arteries of 1,325 patients with type 1 diabetes, 19–51 years of age, with duration of diabetes ranging from 6.3 to 26.1 years. An age- and sex-matched nondiabetic population ( $n = 153$ ) was studied with the same protocol. The ultrasound protocol was carried out in 28 EDIC clinics by centrally trained and certified sonographers using one of three scanning systems. Determination of IMT from videotaped images was performed by a single reader at the Central Ultrasound Reading Unit. Univariate associations with greater IMT were strongest for older age and longer diabetes duration, greater waist-to-hip ratio (men only), higher blood pressure, higher LDL cholesterol, and smoking. The DCCT therapy group (intensive versus conventional) and  $HbA_{1c}$  measured at the time of the ultrasound or the mean  $HbA_{1c}$  during the DCCT, were not significantly related to IMT. Multivariate analyses suggest that age, height, smoking, and BMI were the major predictors of common carotid IMT, whereas age, smoking, and LDL cholesterol predicted internal carotid IMT. There were significant differences between the IMT values of the internal carotid artery in the EDIC male cohort and similarly aged male nondiabetic control subjects. There were no significant differences between the IMT values in the EDIC female cohort and similarly aged female nondiabetic control

subjects. At this point in the planned 10-year follow-up of the DCCT cohort, neither intensive therapy nor  $HbA_{1c}$  level appears to influence the early signs of atherosclerosis. Traditional risk factors, including age, smoking, and LDL cholesterol, were related to IMT. As the cohort is only now entering the age interval during which rapid progression and clinical expression of atherosclerosis are expected, further follow-up will help to determine the role of hyperglycemia, and its interaction with other risk factors, on the development of atherosclerosis. *Diabetes* 48:383–390, 1999

Type 1 diabetes is well recognized as a risk factor for early cardiovascular disease (CVD), leading to a more than 10-fold increase in risk in young adults (1–3) and greatly reducing the sex differential in CVD seen in the general population. The mechanism that underlies this effect of diabetes is unclear, and whether type 1 diabetes initiates the atherosclerotic process early or merely hastens the process once started is controversial (2,4). Furthermore, few studies have delineated the risk factors for CVD in type 1 diabetes, although data confirm a strong relationship with renal disease (5), particularly in men (6). Other risk factors of interest include the lipoprotein profile, blood pressure (BP), fibrinogen level, waist-to-hip ratio (WHR), and depressive symptomatology, particularly in women (6).

Carotid ultrasonography, measuring both the presence of stenosis and intima-media wall thickness (IMT), has provided a powerful noninvasive technique to determine atherosclerosis (7–13). IMT has been extensively used as an outcome measure in clinical trials (14–20). Strong correlations between IMT and cardiovascular risk factors and coronary artery disease (CAD) have been demonstrated in the general population (21,22). However, few studies have been performed in diabetes, particularly in type 1 diabetic populations. The Atherosclerosis Risk in Communities Study demonstrated greater IMT in both diabetic subjects (mainly those with type 2 diabetes) and nondiabetic participants with moderate hyperglycemia compared with subjects with normal glucose levels (23). Similarly, IMT was demonstrated to be increased in type 1 and type 2 diabetic Japanese subjects compared with those without diabetes (24). Two studies from Italy (25,26) have also reported increased IMT and a higher presence of carotid plaques and stenoses in type 2 diabetes (25). More relevant to the current study, young type 1 diabetic subjects (10–25 years of age) in Japan have increased IMT compared with control subjects (27).

The EDIC Research Group (see APPENDIX) is sponsored by the Division of Diabetes, Endocrinology, and Metabolic Diseases of the National Institute of Diabetes and Digestive and Kidney Diseases through research contracts and the General Research Center Program, National Center for Research Resources, the National Institutes of Health.

Address correspondence and reprint requests to EDIC Research Group, Box NDIC/DCCT, Bethesda, MD 20892.

Received for publication 27 March 1998 and accepted in revised form 17 September 1998.

AER, albumin excretion rate; BP, blood pressure; CAD, coronary artery disease; CVD, cardiovascular disease; dBp, diastolic blood pressure; DCCT, Diabetes Control and Complications Trial; ECG, electrocardiogram; EDIC, Epidemiology of Diabetes Interventions and Complications; GFR, glomerular filtration rate; IMT, intima-media wall thickness; sBP, systolic blood pressure; WHR, waist-to-hip ratio.

Data concerning determinants of IMT in type 1 diabetes are also limited. In a Japanese population (27), lipoproteins, BP, and HbA<sub>1c</sub> were not related to IMT in 105 type 1 diabetic subjects. In contrast, in an even smaller study of type 1 diabetes ( $n = 31$ ), age and HbA<sub>1c</sub> predicted IMT (28). The role of hyperglycemia (or glycemic control) as a risk factor for CVD or atherosclerosis in type 1 diabetes is poorly understood. Recent data in type 1 diabetes suggest little, if any, effect of HbA<sub>1c</sub> on CVD (6,8,29). In long-term follow-up of the original 102 subjects in the Stockholm Diabetes Intervention Study, the patients originally assigned to intensive therapy had decreased IMT of the left but not the right common carotid artery, compared with the patients originally assigned to conventional treatment (30).

To address the limited studies and contradictory results with regard to the course of CVD in type 1 diabetes and the role of risk factors for atherosclerosis in type 1 diabetes, carotid ultrasonography was performed on prior participants of the Diabetes Control and Complications Trial (DCCT) (31) who are taking part in a longitudinal follow-up study called the Epidemiology of Diabetes Interventions and Complications (EDIC) (31a). The DCCT cohort (31), which excluded patients with hypertension, hyperlipidemia, and known CAD at baseline, provides an excellent opportunity to examine a large population of patients with type 1 diabetes without obvious CVD risk at baseline who have had careful prospective measurements of many of the putative risk factors for CVD previously established in type 2 diabetes and in nondiabetic populations. The DCCT cohort also provides the opportunity to examine the effects of diabetes treatment on CVD. The randomized interventions during the DCCT might influence the development of CVD, either directly by altering

glucose levels or indirectly by altering lipid levels, the development of nephropathy or other risk factors. In addition, other side effects of intensive therapy documented in the DCCT, such as weight gain and increased rates of hypoglycemia, might also affect CVD risk.

Here we present the results of carotid arterial sonographic measurements of 1,325 patients during the baseline data collection of the EDIC study. The primary aims of this baseline analysis are to 1) determine the feasibility of implementing the protocol in 28 sites; 2) compare the EDIC cohort results with normative data from individuals matched for age and sex; 3) examine the intercorrelations of IMT with demographic, clinical, and biochemical covariates that have been reported previously by other investigators; and 4) determine whether the antecedent DCCT therapy (intensive versus conventional) and the different levels of glycemia achieved resulted in differences in IMT.

## RESEARCH DESIGN AND METHODS

**Subject population.** The original cohort of the DCCT consisted of 1,441 men and women who were 13–40 years of age and had type 1 diabetes for 1–15 years at entry (31). They entered the DCCT between 1983 and 1989 and were studied for an average of 6.5 years. A total of 730 patients were randomly assigned to conventional diabetes treatment and 711 to intensive diabetes treatment. In 1993, the DCCT was stopped because of evidence of a powerful salutary effect of intensive therapy on retinal, renal, and neurological long-term complications (31). At study close-out, DCCT subjects were informed of and invited to join EDIC, a multicenter longitudinal observational study. Of the 1,425 living members of the original cohort, 1,388 (96%) elected to participate in some or all aspects of EDIC. The carotid ultrasound protocol was carried out in 1,325 patients (92% of the original DCCT cohort) as part of the EDIC baseline examination. Table 1 presents the clinical characteristics of these 1,325 patients at EDIC baseline, stratified by original treatment group and sex.

TABLE 1  
Baseline clinical characteristics of EDIC participants by sex and original treatment group assignment during the DCCT

	Women			Men		
	Intensive	Conventional	<i>P</i> value	Intensive	Conventional	<i>P</i> value
<i>n</i>	321	313	—	340	351	—
Attained age (years)	35 ± 7	34 ± 7	0.042	36 ± 7	36 ± 7	0.912
Attained duration of type 1 diabetes (months)	168 ± 58	170 ± 61	0.722	167 ± 59	159 ± 55	0.096
Height (cm)	165 ± 6	165 ± 6	0.682	178 ± 7	179 ± 7	0.076
BMI (kg/m <sup>2</sup> )	26.5 ± 4.6	25.1 ± 3.6	<.001	26.8 ± 4.2	25.9 ± 3.2	0.016
BMI >27 (%)	39.7	27.5	0.001	41.2	34.5	0.069
Natural WHR	0.76 ± 0.07	0.76 ± 0.07	0.731	0.88 ± 0.08	0.87 ± 0.09	0.101
sBP (mmHg)	114 ± 12	114 ± 13	0.739	119 ± 11	120 ± 12	0.186
dBp (mmHg)	74 ± 9	72 ± 10	0.101	77 ± 9	77 ± 8	0.446
Hypertensive (%)	9.7	12.9	0.210	21.2	17.6	0.240
Total cholesterol (mg/dl)	188 ± 36	188 ± 39	0.495	187 ± 35	184 ± 36	0.166
HDL cholesterol (mg/dl)	59 ± 14	59 ± 14	0.737	49 ± 13	50 ± 12	0.742
LDL cholesterol (mg/dl)	113 ± 29	113 ± 32	0.542	119 ± 30	115 ± 31	0.145
Triglycerides (mg/dl)	84 ± 74	82 ± 73	0.427	98 ± 77	97 ± 78	0.756
AER (mg/24 h)	21.1 ± 63.2	63.5 ± 313.9	0.003	28.6 ± 112.9	46.1 ± 123.2	0.038
AER 40 mg/24 h (%)	6.9	16.0	<.001	7.4	17.9	<.001
GFR	117.5 ± 23.5	119.3 ± 26.1	0.069	115.3 ± 19.7	116.1 ± 24.8	0.335
Insulin dose (U · kg <sup>-1</sup> · day <sup>-1</sup> )	0.63 ± 0.21	0.64 ± 0.19	0.801	0.67 ± 0.24	0.65 ± 0.20	0.269
Cigarette smoking (current) (%)	19.7	18.1	0.605	19.3	18.4	0.750
HbA <sub>1c</sub>	7.9 ± 1.4	8.2 ± 1.5	0.009	7.9 ± 1.2	8.4 ± 1.3	<.001
Mean HbA <sub>1c</sub> during DCCT	7.3 ± 0.9	9.0 ± 1.4	<.001	7.2 ± 0.9	9.0 ± 1.1	<.001
Framingham score	0.018 ± 0.025	0.017 ± 0.028	0.037	0.039 ± 0.038	0.037 ± 0.035	0.704

Data are means ± SD, *n*, or %. Baseline is that at EDIC baseline (1994–1995) after an average of 6.5 years of intensive treatment in DCCT. Hypertensive is defined as sitting sBP ≥ 140 mmHg and/or dBp ≥ 90 mmHg or the use of antihypertensive medication. Framingham score is defined as described by Anderson et al. (An Updated Coronary Risk Profile. *Circulation* 83:355–365, 1991).

**Ultrasonography and image analysis.** Carotid ultrasonography was performed between June 1994 and April 1995 (1–2 years after the close of the DCCT) at 28 clinical centers using one of three machines (Toshiba, American Medical Systems, Tustin, CA; ATL Ultra Mark 9; Advanced Technology Laboratory, Bothell, WA; and Acuson XT 128; Mt. View, CA). The criteria used to select these machines included the following:

- Accurate delineation of near and far wall boundaries,
- Accurate plaque detection and sizing,
- Simultaneous Doppler, preferably with color,
- Detection and quantification of early subintimal change,
- Accurately reproduced images on videotape using S-VHS recorders.

The EDIC Ultrasound Scanning Protocol was adapted from procedures used in the Cardiovascular Health Study (7), the Insulin Resistance Atherosclerosis Study (32), and the Mexican-American Heart Study (33).

The imaging protocol involved obtaining a single longitudinal lateral view of the distal 10 mm of the right and left common carotid arteries and three longitudinal views in different imaging planes of each internal carotid artery. The internal carotid artery was defined as including both the carotid bulb, identified by the loss of the parallel wall present in the common carotid artery, and the 10-mm segment of the internal carotid artery distal to the tip of the flow divider that separates the external and internal carotid arteries (Fig. 1).

Centrally trained and certified sonographers conducted the studies. Studies were recorded on S-VHS tapes and sent weekly to the Central Ultrasound Reading Unit. The maximum IMT of the common carotid artery was defined as the mean of the maximum IMT for near and far walls in both right and left sides. The maximum IMT of the internal carotid artery was defined in the same way, and the results from the three scans were averaged (anterior, lateral, and posterior views on both sides).

**Normative data.** During the first year of the EDIC, each of the 28 clinics performed six carotid ultrasounds on nondiabetic subjects who were between the ages of 20 and 50 years with no history of CVD, hypertension, or stroke ( $n = 153$ ). The mean age ( $35.3 \pm 8.5$ ) and percentage of women (49.7%) were not significantly different than those of the EDIC cohort.

**Other procedures.** On the anniversary of enrolling in the DCCT, each EDIC subject has a standardized annual history and physical examination, including a detailed evaluation of overall health, diabetes management, occurrence of diabetic complications, development of new disease, and medications used. Annual evaluations also include resting electrocardiograms (ECGs), Doppler ultrasound measurements of ankle/arm BP ratios, and arm BPs. Serum creatinine and HbA<sub>1c</sub> are determined as they were in the DCCT (34). Lipid profiles and 4-h urine collections for measurement of albumin excretion rate and creatinine clearance are obtained in alternate years using the same methods as in the DCCT (35).

**Statistical analysis.** Quality scores for carotid ultrasound scans were based on the number of lines visualized from the eight views. The proportion of lines with quality scores of good or excellent were compared across the 28 clinics.

To evaluate the possible association with other covariates, multiple linear regression models were fit to the average maximum IMTs of the common and internal carotid arteries. Both models were fit by ordinary least squares, but a reciprocal transformation had to be applied to internal IMT to obtain a homoscedastic and approximately normal residual distribution. The semipar-

tial  $R^2$  for each variable measures the increase in  $R^2$  obtained by introducing that variable into a model already including all the others.


Wilcoxon rank-sum tests were used to make separate comparisons of the distributions of common IMT and internal IMT between EDIC patients and nondiabetic control subjects within strata defined by age and sex. Separate linear models for each segment were used to adjust for covariance with age before testing; EDIC patients and control subjects were adjusted jointly but stratified by sex. Comparison of IMTs between the DCCT randomized treatment groups was made using the Wei-Lachin test of stochastic ordering (36), which has greatest power against alternatives in which both segments tend to be thicker in the same treatment group. Tests were stratified by sex and decade of age, and an overall test adjusted for between-strata differences (36) was also performed. Based on the results of the multiple linear regression models, the EDIC patients' IMT values were first adjusted for covariance with age and cumulative pack-years of cigarette smoking. All  $P$  values are reported at their nominal levels, without adjustment for multiple comparisons.

## RESULTS

**Data quality.** Baseline reproducibility analyses of 140 replicate measures resulted in absolute mean differences of 0.04 mm for both the common carotid and the intimal carotid. Intraclass correlation between original and re-readings of maximum wall thickness were 0.71 and 0.89 for the common and for internal carotid arteries, respectively.

Carotid artery data by clinic were generally of uniform high quality. The percentage of scans of the common carotid artery in which all six lines were legible and graded as good or excellent ranged from 90 to 100% across clinics. Visualization of the internal carotid artery was less uniform, with the percentage of scan with all six lines legible ranging from 15 to 94% of scans. Scans that did not meet the criteria (mainly segments of the internal carotid) had only four or five lines legible. A process for continuous quality control was instituted so that information was fed back to each center in an effort to improve scan quality.

**Comparison of nondiabetic control subjects with EDIC subjects.** Table 2 presents the IMT data including the average maximum of the common and internal carotid arteries stratified by sex and age decades for nondiabetic and type 1 diabetic subjects. A Wilcoxon test of the difference between sexes for both diabetic and nondiabetic groups indicates highly significant ( $P = 0.0001$ ) differences, with men having greater IMT mean thickness. Adjusting for covariance with height reduces but does not eliminate the significance ( $P =$



**FIG. 1.** Schematic drawing of the carotid artery: the bifurcation of the common into the internal and external carotid arteries, the location of the segments scanned in the EDIC, and the six lines to be measured in the sonographic images.

TABLE 2  
Comparison of IMT in nondiabetic subjects with EDIC type 1 diabetes cohort

Age (years)	Women					Men				
	Nondiabetic		Type 1 diabetes			Nondiabetic		Type 1 diabetes		
	<i>n</i>	IMT (mm)	<i>n</i>	IMT (mm)	<i>P</i>	<i>n</i>	IMT (mm)	<i>n</i>	IMT (mm)	<i>P</i>
<b>Common carotid artery</b>										
20–29	25	0.603 ± 0.062	172	0.616 ± 0.073	0.471	21	0.648 ± 0.098	125	0.636 ± 0.059	0.971
30–39	27	0.663 ± 0.074	278	0.657 ± 0.081	0.453	25	0.657 ± 0.076	350	0.684 ± 0.083	0.103
40–49	25	0.704 ± 0.078	178	0.696 ± 0.079	0.622	30	0.741 ± 0.094	211	0.745 ± 0.104	0.925
<b>Internal carotid artery</b>										
20–29	25	0.547 ± 0.053	172	0.583 ± 0.092	0.065	21	0.588 ± 0.085	125	0.629 ± 0.083	0.032
30–39	27	0.632 ± 0.153	278	0.632 ± 0.147	0.859	25	0.645 ± 0.104	350	0.684 ± 0.114	0.071
40–49	25	0.655 ± 0.074	178	0.719 ± 0.226	0.704	30	0.741 ± 0.147	211	0.806 ± 0.261	0.437

Data are means ± SD or *n*. Nondiabetic data were obtained from six normal subjects without diabetes at each of the 29 clinics using the EDIC ultrasound scanning protocol and read at the Central Ultrasound Reading Unit. Data exclude six EDIC patients who were 19 years old when scanned and five who were aged 50 or 51 years old. *P* values are from Wilcoxon rank-sum test after linear adjustment for covariance with age. *P* < 0.0001 for both common and internal IMT, type 1 men versus type 1 women; *P* = 0.59 for common IMT, *P* = 0.004 for internal IMT, type 1 men versus nondiabetic men; *P* = 0.89 for common IMT, *P* = 0.11 for internal IMT, type 1 women versus nondiabetic women.

0.0007). A test of trend of the increasing maximum wall thickness over the age decades is significant in all strata (e.g., nondiabetic men, type 1 diabetic men, etc.).

Comparing the nondiabetic population with the sex-matched control subjects revealed that overall the type 1 diabetic men had greater IMT thickness for the internal carotid (0.713 ± 0.184 vs. 0.664 ± 0.130 mm, *P* = 0.004) but not for the common carotid (0.694 ± 0.10 vs. 0.686 ± 0.10 mm, *P* = 0.59). Within the different age-groups, only the youngest group of men had significantly different internal carotid IMT than their age-matched control subjects (0.629 vs. 0.588, *P* = 0.03). The type 1 diabetic women's mean IMT thickness was similar to the nondiabetic women's for both the common and internal carotid.

**Univariate correlation.** Table 3 presents univariate Spearman correlations of various clinical characteristics with average maximum wall thickness for the common and internal carotid arteries. Attained age and Framingham risk scores were the variables most strongly associated with IMT in both the common and internal carotid arteries. Attained duration of type 1 diabetes had a weaker, but statistically significant, association with wall thickness that, in men, persisted after adjusting for age. Current HbA<sub>1c</sub> was not correlated with wall thickness; however, HbA<sub>1c</sub> at the time of carotid studies did not reflect glycemic exposure during the DCCT, since all subjects were encouraged to adopt intensive therapy at the end of the DCCT. Therefore, correlations between mean HbA<sub>1c</sub> during the DCCT and IMTs were also computed. Mean HbA<sub>1c</sub> during the DCCT was not significantly correlated with either common or internal carotid IMT among men or with internal IMT among women. The correlation of mean HbA<sub>1c</sub> with common IMT in women, while nominally significant, was weak (*r* = -0.11, *P* = 0.008) and indicated an inverse relationship. Total and LDL cholesterol levels were correlated with wall thickness, but HDL cholesterol and triglyceride levels were not. Systolic blood pressure (sBP) was associated with IMT in both men and women, and diastolic blood pressure (dBp) was associated with IMT in men only. WHR cor-

related with IMT in men, but not women, while total pack-years of cigarette smoking was correlated with both common and internal carotid IMT in both sexes. Correlations were similar in the two treatment groups.

**Multivariate analyses.** Table 4 shows the multiple linear regression analysis. Apart from age, the major predictors of common carotid IMT in women were height, glomerular filtration rate (GFR), BMI, and smoking, while in men, height, smoking, BMI, sBP, duration of type 1 diabetes, and average LDL cholesterol during the DCCT were independent predictors. Overall, 29 and 20% of the variance was explained in men and women, respectively. For internal carotid IMT, age, sBP, and LDL were the major predictors in both sexes, and smoking had a significant but reduced effect compared with the model for the common carotid, especially in men. GFR was also a predictor in men. Again, more of the variance was explained in men (24%) than in women (17%).

**Tests for treatment group difference.** Table 5 presents IMT by sex, age, and treatment group. There are no strata in which intensive treatment group patients had significantly different age- and smoking-adjusted wall thicknesses than conventional treatment group patients. The difference between intensive and conventional treatment from the N-weighted test of stochastic ordering for the combined strata was not statistically significant (*P* = 0.39).

**DISCUSSION**

The current report confirms the feasibility of using carotid ultrasonography in a large multicenter study with excellent reproducibility for common carotid and acceptable reproducibility for internal carotid IMT measurements (37,38). The quality of scans was good or excellent for the vast majority of centers, especially for the common carotid artery.

There was no significant difference in mean IMT between EDIC subjects and age- and sex-matched control subjects, except for internal carotid IMT among men. Because two major determinants of IMT are age and BP, the similarity in IMT may reflect the young age of the EDIC subjects and the

TABLE 3  
Spearman correlation coefficients of clinical characteristics with average maximum IMT

	Sex	Common carotid artery		Internal carotid artery	
		Correlation	Pvalue	Correlation	Pvalue
Attained age (years)	M	0.46	<0.001	0.38	<0.001
	F	0.42	<0.001	0.35	<0.001
Attained duration of type 1 diabetes (months)	M	0.13	<0.001	0.12	0.002
	F	0.08	0.04	0.12	0.004
Height (cm)	M	0.17	<0.001	0.05	0.19
	F	0.12	0.003	-0.04	0.36
BMI (kg/m <sup>2</sup> )	M	0.15	<0.001	0.09	0.02
	F	0.06	0.12	0.02	0.56
WHR	M	0.22	<0.001	0.14	<0.001
	F	0.01	0.89	0.01	0.85
sBP (mmHg)	M	0.21	<0.001	0.14	<0.001
	F	0.14	<0.001	0.11	0.007
dBP (mmHg)	M	0.11	0.004	0.13	<0.008
	F	0.07	0.08	0.07	0.09
Total cholesterol (mg/dl)	M	0.21	<0.001	0.23	<0.001
	F	0.09	0.04	0.13	0.002
HDL cholesterol (mg/dl)	M	-0.03	0.37	0.00	0.92
	F	0.00	0.96	0.03	0.41
LDL cholesterol (mg/dl)	M	0.24	<0.001	0.26	<0.001
	F	0.08	0.06	0.13	0.001
Triglycerides (mg/dl)	M	0.07	0.06	0.05	0.19
	F	0.06	0.14	0.00	0.94
Smoking (total pack-years)	M	0.21	<0.001	0.09	0.02
	F	0.15	<0.001	0.16	<0.001
Insulin dose (U · kg <sup>-1</sup> · day <sup>-1</sup> )	M	-0.06	0.10	-0.10	0.02
	F	-0.08	0.06	-0.05	0.20
HbA <sub>1c</sub>	M	0.07	0.07	0.02	0.56
	F	-0.02	0.62	0.02	0.68
Mean HbA <sub>1c</sub> during the DCCT	M	0.06	0.12	-0.02	0.57
	F	-0.11	0.008	-0.07	0.07
Framingham risk score	M	0.47	<0.001	0.38	<0.001
	F	0.42	<0.001	0.35	<0.001
AER (mg/24 h)	M	0.10	0.01	0.07	0.09
	F	-0.01	0.76	0.01	0.85
GFR	M	-0.01	0.76	0.02	0.54
	F	-0.05	0.25	-0.02	0.65

initial exclusion of hypertensive subjects in the DCCT. These data are in contrast to those of Yamasaki et al. (27), who reported significantly greater IMT for 10- to 25-year-old type 1 diabetic subjects compared with nonmatched control subjects, despite the smaller number of type 1 diabetic patients and control subjects (<10% and 30%, respectively) in the Japanese study. The discrepancy between the Japanese and EDIC studies may be secondary to the unmatched control group in the Japanese study, racial differences in the pathogenesis of atherosclerosis, or other factors. The finding of increased IMT in type 2 diabetic subjects (23,26) may be secondary to older age or the high prevalence of other CVD risk factors found in type 2 diabetes that were not present in the EDIC population.

Another possible explanation of the lack of a marked difference in IMT in our type 1 diabetic subjects is the possibility that IMT does not reflect the aspects, or stages, of atherosclerosis that are enhanced or exacerbated by type 1 diabetes. Although IMT is thought to be an intermediate biological marker of atherosclerosis and correlates with the presence of plaque or clinical events (39), it does not provide

a direct measure of occlusive disease or plaque stability. Nor does it directly measure the hemostatic (and possibly inflammatory) disturbances that may play a vital role in atherosclerotic disease and that are disturbed in diabetes, such as abnormal fibrinogen levels (40) and platelet function (41). The true predictive power of IMT for future events is uncertain, and the data supporting a correlation with existent disease, while encouraging, are still limited (21,22,39,42,43). These potential limitations, however, do not negate the value of carotid ultrasonography in diabetes, which may throw more light on the pathogenesis of atherosclerotic disease generally, as well as on the specific enhanced risk in diabetes. Only careful follow-up with repeated sonography and assessment of clinical outcomes will allow definitive determination of the effect of diabetes on the overall natural history of atherosclerotic CVD.

The current study has limitations, the major one being that the DCCT cohort was a selected trial population and not necessarily representative of type 1 subjects in the general population. However, the DCCT conventional treatment group was found to be generally comparable to a subset of a

TABLE 4  
Summary of multiple linear regression models for common carotid IMT and 1/internal carotid IMT

	Common			1/Internal		
	Slope	P	Semipartial R <sup>2</sup> (%)	Slope	P	Semipartial R <sup>2</sup> (%)
<b>Women</b>						
Age	0.00435	<0.001	11.95	-0.01176	<0.001	7.02
Type 1 diabetes duration (months)	0.00006	0.267	0.21	-0.00034	0.078	0.54
Height	0.00131	0.011	1.11	0.00188	0.315	0.18
Smoking (pack-years)	0.00076	0.062	0.60	-0.00367	0.013	1.07
BMI	0.00155	0.050	0.67	0.00122	0.670	0.03
sBP	0.00028	0.307	0.18	-0.00217	0.027	0.85
LDL cholesterol (DCCT average)	0.00009	0.507	0.08	-0.00172	<0.001	2.00
GFR	0.00025	0.053	0.65	-0.00056	0.230	0.25
HbA <sub>1c</sub> (DCCT average)	-0.00310	0.274	0.21	0.01192	0.247	0.23
Intensive therapy	-0.00522	0.515	0.07	0.02407	0.407	0.12
<b>Men</b>						
Age	0.00451	<0.001	9.35	-0.01167	<0.001	7.19
Type 1 diabetes duration (months)	0.00018	0.001	1.53	-0.00046	0.006	1.13
Height	0.00161	<0.001	2.15	-0.00051	0.684	0.03
Smoking (pack-years)	0.00149	<0.001	2.52	-0.00263	0.015	0.89
BMI	0.00322	<0.001	2.06	-0.00131	0.612	0.04
sBP	0.00085	0.002	1.38	-0.00201	0.016	0.88
LDL cholesterol (DCCT average)	0.00033	0.009	1.03	-0.00220	<0.001	5.07
GFR	0.00025	0.072	0.49	-0.00105	0.012	0.94
HbA <sub>1c</sub> (DCCT average)	0.00226	0.481	0.08	0.00894	0.350	0.13
Intensive therapy	-0.00985	0.241	0.21	0.01190	0.635	0.03

Total women in the common group is 20.5%; total women in the 1/internal group is 17.4%; total men in the common group is 29.4%; total men in the 1/internal group is 24.3%.

population-based type 1 diabetes cohort (44). A further limitation to the study was the exclusion at baseline of patients with hypertension and hyperlipidemia. As noted, the relatively young age of the EDIC cohort and the exclusions noted above may have reduced the occurrence of CVD and the ability to detect risk factor associations and differences in IMT at the current time. It should also be noted that the current analysis is essentially cross-sectional (albeit with a well-documented and randomized historical prospective measure of

glycemic exposure) and is subject to the limitations of these studies, in particular, the absence of baseline (carotid) studies. Cardiovascular risk profile, however, did not differ by DCCT treatment group at baseline (31).

An increased number of CVD events is very likely to occur in the DCCT cohort during the 10 years of the EDIC. By the end of the 10-year follow-up, the mean age of the study population, the major predictor of CAD, will approach 43 years, and type 1 diabetes duration will average 22 years. Based on

TABLE 5  
Average maximum IMTs by age and randomized treatment assignment

Sex	Age (years)	Common carotid artery				Internal carotid artery		P
		n	Intensive	n	Conventional	Intensive	Conventional	
<b>Women</b>								
	20–29	78	0.626 ± 0.078	94	0.608 ± 0.068	0.576 ± 0.071	0.589 ± 0.107	0.809
	30–39	137	0.651 ± 0.086	141	0.663 ± 0.076	0.628 ± 0.129	0.636 ± 0.164	0.437
	40–49	102	0.698 ± 0.079	76	0.693 ± 0.080	0.704 ± 0.207	0.738 ± 0.250	0.954
<b>Men</b>								
	20–29	65	0.637 ± 0.062	60	0.635 ± 0.056	0.636 ± 0.091	0.622 ± 0.072	0.743
	30–39	167	0.680 ± 0.079	183	0.687 ± 0.086	0.684 ± 0.107	0.685 ± 0.120	0.807
	40–49	104	0.728 ± 0.084	107	0.761 ± 0.118	0.791 ± 0.211	0.820 ± 0.302	0.149
<b>Combined (stratified-adjusted)</b>								0.388

Data are means ± SD or n. P values are from the Wei-Lachin test of stochastic ordering after adjustment of IMT values for covariance with age and pack-years of cigarette smoking. The “combined” test aggregates the within-stratum results while accounting for the differences between age-groups and sex. Means and SDs were calculated from the unadjusted IMTs. The table excludes 11 patients who were <20 or >49 years old when tested.

estimates derived from previous studies (1–6), the prevalence of CAD, as manifested clinically and/or as detected by ECG or exercise tolerance testing, is likely to be 15–40%. It seems likely that we shall have sufficient power to determine the relationship of IMT to CVD events occurring over the subsequent 10 years.

Documented risk factors for increased IMT include age, hypertension, lipid profile, and smoking. Our data suggest that these factors relate to IMT in type 1 diabetes. Among established lipid risk factors, only LDL level correlated with IMT. Some risk factors appeared to affect IMT in the common carotid, while others affected the internal carotid wall. For example, in multivariate analyses, smoking was more strongly related to common carotid IMT than to internal carotid, while LDL cholesterol was stronger for the internal carotid.

The large sample size in the current study allowed the multivariate analyses to establish a variety of factors including smoking, sBP, GFR, BMI, and LDL cholesterol as predictors of IMT. Some studies in type 2 diabetes have confirmed some of these associations, e.g., BP (24–26) and hyperlipidemia (24), while other studies in type 2 diabetes have failed to find such associations (27,28), possibly owing to their small sample size. The relationship between GFR and IMT may reflect the recognized association between renal disease and CVD in type 1 diabetes (5,6).

The role of glycemic control (or glycemic level) is of particular interest. The few data available relating glycemia to CVD in type 1 diabetes are controversial. The DCCT results suggested a borderline salutary effect of intensive treatment on combined macrovascular events (45). A follow-up report by the Stockholm Diabetes Intervention Study suggested that intensive treatment leads to reduced IMT thickness and less stiffening of the carotid arteries (30), although the results were not uniformly supportive. The current study does not support an association between HbA<sub>1c</sub> at the time of carotid ultrasonography or the mean HbA<sub>1c</sub> during the 6.5 years of the DCCT and IMT. In addition, treatment assignment during the DCCT did not have an apparent effect on IMT. It may be argued that the absence of a major difference in HbA<sub>1c</sub> at the time of ultrasonography between the two groups might explain the absence of an effect, and that 1.5 years of similar control after the DCCT may obliterate any benefit that 6.5 years of better control may have had. However, as much of the adverse effect of hyperglycemia in atherosclerosis is thought to be a chronic effect, e.g., cumulative advanced glycosylation end product formation (46), some differential effect of DCCT exposure might reasonably be expected. However, the current findings do not support an obvious association between glycemia and CVD in type 1 diabetes, consistent with several previous studies (3,6,28). Further follow-up will permit more definitive assessments as age, IMT, atherosclerosis, and clinical events all increase. Future studies will also help determine whether recent American Diabetes Association guidelines on the management of diabetic dyslipidemia in type 2 diabetes (47) should apply to type 1 diabetic subjects as well. The predictive power of LDL cholesterol for IMT in the current study provides some further support for aggressive cholesterol lowering (47). Intensive glycemic control in type 1 diabetes, nonetheless, remains the central component of diabetes management in view of its undoubted benefit in delaying or preventing microvascular complications and improving the CVD risk profile.

## ACKNOWLEDGMENTS

The Office of Research on Women's Health at the National Institutes of Health funded the carotid ultrasounds.

## APPENDIX: THE EDIC RESEARCH GROUP

Study chairmen: S. Genuth, D. Nathan; Albert Einstein College of Medicine: H. Shamoon, H. Duffy; Case Western Reserve University: B. Dahms, L. Mayer; Cornell University Medical Center: D. Brillion, M. Lackaye; Henry Ford Health System: F. Whitehouse, D. Kruger; International Diabetes Center: R. Bergenstal, M. Johnson; Joslin Diabetes Center: A. Jacobson, J. Doyle, M. Waters; Massachusetts General Hospital: D. Nathan, S. Crowell, J. Godine, C. McKittrick; Mayo Foundation: J. Service, G. Ziegler; Medical University of South Carolina: J. Colwell, D. Wood, R. Mayfield; Northwestern University: M. Molitch, B. Schaefer; University of California, San Diego: O. Kolterman, G. Lorenzi; University of Iowa: W. Sivitz, M. Bayless; University of Maryland School of Medicine: D. Counts, A. Kowarski (past member), D. Ostrowski; University of Michigan: D. Greene, C. Martin; University of Minnesota: J. Bantle, B. Rogness; University of Missouri: D. Goldstein, A. Smith; University of New Mexico: D. Schade, C. Johannes; University of Pennsylvania: S. Schwartz, B.J. Maschak-Carey; University of Pittsburgh: T. Orchard, N. Silvers; University of South Florida: J. Malone, A. Mangione; University of Tennessee: A. Kitabchi, M.B. Murphy; University of Texas Southwestern University Medical Center: P. Raskin, S. Strowig; University of Toronto: B. Zinman, A. Barnie; University of Washington: J. Palmer, J. Ginsberg; University of Western Ontario: J. Dupre, J. Harth; Vanderbilt University: R. Lorenz, J. Lipps; Washington University, St. Louis: N. White, J. Santiago (deceased), L. Levandoski; Yale University School of Medicine: W. Tamborlane, P. Gatcomb; clinical coordinating center (Case Western Reserve University): B. Dahms, P. Corcoran, J. Quin; data coordinating center (The George Washington University, Biostatistics Center): J. Lachin, P. Cleary, D. Kenny, L. Diminick, D. Lamas; National Institute of Diabetes and Digestive and Kidney Diseases program office: C. Cowie, R. Eastman; central fundus photograph reading center (University of Wisconsin): M. Davis, L. Hubbard, P. Geithman, J. Brickbauer, L. Kastorff, M. Neider; central biochemistry laboratory (University of Minnesota): M. Steffes, J. Buckska, B. Chavers; central carotid ultrasound unit (New England Medical Center): D. O'Leary; central electrocardiogram reading unit (University of Minnesota): R. Crow, C. O'Donnell; external advisory committee: G. Weir (chairman), C. Clark, R. D'Agostino, M. Espeland, B. Klein, H. Jacobson, T. Manolio, L. Rand, D. Singer, M. Stern; writing team: T. Orchard, P. Cleary, B. Zinman, D. O'Leary; and editor, EDIC publications: D. Nathan.

## REFERENCES

1. Dorman JS, LaPorte RE, Kuller LH, Cruickshanks KJ, Orchard TJ, Wagener DK, Becker DJ, Cavender DE, Drash AL: The Pittsburgh insulin-dependent diabetes mellitus (type 1 diabetes) morbidity and mortality study: mortality results. *Diabetes* 33:271–277, 1984
2. Krolewski AS, Kosinski EJ, Warram JH, Leland OS, Busick EJ, Asmal AC, Rand LI, Christlieb AR, Bradley RF, Kahn CR: Magnitude and determinants of coronary artery disease in juvenile-onset insulin-dependent diabetes mellitus. *Am J Cardiol* 59:750–755, 1987
3. Maser RE, Wolfson SK, Stein EA, Drash AL, Becker DJ, Dorman JS, Ellis D, Orchard TJ: Cardiovascular disease and arterial calcification in insulin-dependent diabetes mellitus: interrelationships and risk factor profiles: Pittsburgh Epidemiology of Diabetes Complications Study-VI. *Arterioscler Thromb* 11:958–965, 1991
4. Orchard TJ: Summary and comment: magnitude and determinants of coronary

- artery disease in juvenile-onset insulin-dependent diabetes mellitus. *Diabetes Spectrum* 4:344–345, 1991
5. Borch-Johnsen K, Andersen PK, Deckert T: The effect of proteinuria on relative mortality in type 1 (insulin-dependent) diabetes mellitus. *Diabetologia* 28:590–596, 1985
  6. Lloyd CE, Kuller LH, Becker DJ, Ellis D, Wing RR, Orchard TJ: Coronary artery disease in type 1 diabetes: gender differences in risk factors, but not risk. *Arterioscler Thromb Vasc Biol* 16:720–726, 1996
  7. Fried LP, Borhani NO, Enright P, Furberg CD, Gardin JM, Kronmal RA, Kuller LH, Manolio TA, Mittelmark MB, Newman A, O'Leary DH, Psaty B, Rautaharju P, Tracy RP, Weiler PG: The Cardiovascular Health Study: design and rationale. *Ann Epidemiol* 1:263–276, 1991
  8. The ARIC Investigators: The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. *Am J Epidemiol* 129:687–702, 1989
  9. Salonen R, Salonen JT: Progression of carotid atherosclerosis and its determinants: a population-based ultrasonography study. *Atherosclerosis* 81:33–40, 1990
  10. Burke GL, Evans GW, Riley WA, Sharrett AR, Howard G, Barnes RW, Rosamond W, Crow RS, Rautaharju PM, Geiss G, for the ARIC Study Group: Arterial wall thickness is associated with prevalent cardiovascular disease in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Stroke* 26:386–391, 1995
  11. Allan PL, Mowbray PI, Lee AJ, Fowkes GR: Relationship between carotid intima-media thickness and symptomatic and asymptomatic peripheral arterial disease the Edinburgh Study. *Stroke* 28:348–353, 1997
  12. Bots ML, Hoes AW, Koudstaal PJ, Hofman A, Grobbee DE: Common carotid artery intima-media thickness and risk of stroke and myocardial infarction: the Rotterdam Study. *Circulation* 96:1432–1437, 1997
  13. Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR, Clegg X: Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) Study, 1987–1993. *Am J Epidemiol* 146:483–494, 1997
  14. ACAPS Group: Rationale and design for the Asymptomatic Carotid Artery Plaque Study (ACAPS). *Control Clin Trials* 13:293–314, 1992
  15. Crouse JR, Byington RP, Bond MG, Espeland MA, Sprinkle JW, McGovern M, Furberg CD: Pravastatin, lipids, and atherosclerosis in the carotid arteries: design features of a clinical trial with carotid atherosclerosis outcome. *Control Clin Trials* 13:495–506, 1992
  16. Hennerici M, Klephas W, Gries FA: Regression of carotid plaques during low-density lipoprotein cholesterol elimination. *Stroke* 22:989–992, 1991
  17. Furberg CD, Borhani NO, Byington RP, Gibbons ME, Sowers JR: Calcium antagonists and atherosclerosis: the multicenter Isradipine/Diuretic Atherosclerosis Study. *Am J Hypertens* 6:245–259, 1993
  18. Blankenhorn DH, Selzer RH, Crawford DW, Barth JD, Liu C-R, Liu C-H, Mack WJ, Alaupovic P: Beneficial effects of colestipol-niacin therapy on the common carotid artery: two- and four-year reduction of intima-media thickness measured by ultrasound. *Circulation* 88:20–28, 1993
  19. Hodis HN, Mack WJ, LaBree L, Selzer RH, Liu C, Liu C, Alaupovic P, Kwong-Fu H, Azen SP: Reduction in carotid arterial-wall thickness using lovastatin and dietary therapy: a randomized controlled clinical trial. *Ann Intern Med* 124:548–556, 1996
  20. Furberg CD, Adams HP, Applegate WB, Byington RP, Espeland MA, Hartwell T, Hunninghake DB, Lefkowitz DS, Probstfield J, Riley WA, Young B for the Asymptomatic Carotid Artery Progression Study (ACAPS) Research Group: Effect of lovastatin on early carotid atherosclerosis and cardiovascular events. *Circulation* 90:1679–1687, 1994
  21. Heiss G, Sharrett AR, Barnes R, Chambless LE, Szklo M, Alzola C, ARIC Investigators: Carotid atherosclerosis measured by B-mode ultrasound in populations: associations with cardiovascular risk factors in the ARIC Study. *Am J Epidemiol* 134:250–256, 1991
  22. O'Leary DH, Polak JF, Kronmal RA, Kittner SJ, Bond G, Wolfson SK Jr, Bommer W, Price TR, Gardin JM, Savage PJ: Distribution and correlates of sonographically detected carotid artery disease in the Cardiovascular Health Study. *Stroke* 23:1752–1760, 1992
  23. Folsom AR, Eckfeldt JH, Weitzman S, Ma J, Chambless LE, Barnes RW, Cram KB, Hutchinson RG: Relation of carotid artery wall thickness to diabetes mellitus, fasting glucose and insulin, body size, and physical activity. *Stroke* 25:66–73, 1994
  24. Kawamori R, Yamasaki Y, Matsushima H, Nishizawa H, Nao K, Hougaku H, Maeda H, Handa N, Matsumoto M, Kamada T: Prevalence of carotid atherosclerosis in diabetic patients. *Diabetes Care* 15:1290–1294, 1992
  25. Pujia A, Gnasso A, Irace C, Colonna A, Mattioli PL: Common carotid arterial wall thickness in type 1 diabetes subjects. *Diabetes Care* 17:1330–1336, 1994
  26. Bonora E, Tessari R, Micciolo R, Zenere M, Targher G, Padovani R, Falezza G, Muggeo M: Intimal-medial thickness of the carotid artery in nondiabetic and type 1 diabetes patients. *Diabetes Care* 20:627–631, 1997
  27. Yamasaki Y, Kawamori R, Matsushima H, Nishizawa H, Kodama M, Kajimoto Y, Morishima T, Kamada T: Atherosclerosis in carotid artery of young type 1 diabetes patients monitored by ultrasound high-resolution B-mode imaging. *Diabetes* 43:634–639, 1994
  28. Kanters SDJM, Algra A, Banga J: Carotid intima-media thickness in hyperlipidemic type 1 and type 2 diabetic patients. *Diabetes Care* 20:276–280, 1997
  29. Koivisto VA, Stevens LK, Mattock MB, Ebeling P, Stephenson J, Idzior-Walus B, EURODIAB Type 1 Diabetes Complications Study Group: Cardiovascular disease and its risk factors in type 1 diabetes in Europe. *Diabetes Care* 19:689–697, 1996
  30. Jensen-Urstad KJ, Reichard PG, Rosfors JJS, Lindblad LEL, Jensen-Urstad MT: Early atherosclerosis is retarded by improved long-term blood glucose control in patients with type 1 diabetes. *Diabetes* 45:1253–1258, 1996
  31. The Diabetes Control and Complications Trial Research Group: The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 329:977–986, 1993
  - 31a. EDIC Research Group: Epidemiology of diabetes interventions and complications (EDIC): design, implementation, and preliminary results of a long-term follow-up of the Diabetes Control and Complications Trial cohort. *Diabetes Care* 22:99–111, 1999
  32. D'Agostino RB, Burke G, O'Leary D, Rewers M, Selby J, Savage PJ, Saad MF, Bergman RN, Howard G, Wagenknecht L, Haffner SM: Ethnic differences in carotid wall thickness: the Insulin Resistance Atherosclerosis Study. *Stroke* 27:1744–1749, 1996
  33. Wei M, Gonzalez C, Haffner SM, O'Leary DH, Stern MP: Ultrasonography-assessed maximum carotid wall thickness in Mexico City residents and Mexican-Americans in San Antonio, TX: association with diabetes and cardiovascular risk factors. *Atheroscler Thromb Vasc Biol* 16:1184–1188, 1996
  34. DCCT Research Group: The Diabetes Control and Complications Trial (DCCT): design and methodologic considerations for the feasibility phase. *Diabetes* 35:530–545, 1986
  35. DCCT Research Group: *DCCT Manual of Operations*. Springfield, VA, National Technical Information Service 93-183382, 1993
  36. Lachin JM: Some large-sample distribution-free estimation tests for multivariate partially incomplete data from the populations. *Stat Med* 11:1151–1170, 1992
  37. Crouse JR: B-mode ultrasound in clinical trials. *Circulation* 88:319–320, 1993
  38. Espeland MA, Craven TE, Riley WA, Corson J, Romont A, Furberg CD: Reliability of longitudinal ultrasonographic measurements of carotid intimal-medial thickness. *Stroke* 27:480–485, 1996
  39. Burke GL, Evans GW, Riley WA, Sharrett AR, Howard G, Barnes RW, Rosamond W, Crow RS, Rautaharju PM, Heiss G: Arterial wall thickness is associated with prevalent cardiovascular disease in middle-aged adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Stroke* 26:386–391, 1995
  40. Ganda OP, Arkin CF: Hyperfibrinogenemia: an important risk factor for vascular complications in diabetes. *Diabetes Care* 15:1245–1250, 1992
  41. Mustard JF, Packham MA: Platelets and diabetes mellitus. *N Engl J Med* 297:1345–1347, 1977
  42. Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR, Clegg LX: Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) Study, 1987–1993. *Am J Epidemiol* 146:483–494, 1997
  43. Hodis HN, Mack WJ, LaBree L, Selzer RH, Liu C, Liu C, Azen SP: The role of carotid arterial intima-media thickness in predicting clinical coronary events. *Ann Intern Med* 128:262–269, 1998
  44. Klein R, Santiago J, Cleary P, Moss S, Nathan D: A comparison of the study populations in the Diabetes Control and Complications Trial (DCCT) and the Wisconsin Epidemiologic Study of Diabetic Retinopathy (WESDR). *Arch Intern Med* 155:745–754, 1995
  45. Diabetes Control and Complications Trial Research Group: Effect of intensive diabetes management on macrovascular events and risk factors in the Diabetes Control and Complications Trial. *Am J Cardiol* 75:894–903, 1995
  46. Lyons TJ, Lopes-Virella MF: *Glycosylation-Related Mechanisms in Diabetes and Atherosclerosis: Molecular Basis and Clinical Aspects*. Drazmiz B, Eckel RH, Eds. New York, Elsevier, 1993, p. 191–212
  47. American Diabetes Association: Management of dyslipidemia in adults with diabetes (Position Statement). *Diabetes Care* 21 (Suppl. 1):S36–S44, 1998



Author Queries (please see Q in margin and underlined text)

Q1: Is “*w*” the correct symbol for equations such as “ $w = 153$ ”?

Does it need explanation at first mention?

Q2: The last line of the first footnote has been moved to Acknowledgments.

Q2a: Please supply a running title of 40 characters.

Q3: In Table 1, please supply title of article by Anderson et al.

Q4: Please list the name of the machine from Acuson.

Q5: In Table 2, last line of footnote,  $P = 11$  correct?

Q6: Should CHD be changed to CAD or CVD?

Q7: Stockholm Diabetes Intervention “Society” in discussion vs. “Study” in introduction. OK as is or change needed?

Q8: Appendix—The term “(past)” appears after “A. Kowarski”; does this mean “past member”?

Q9: For ref. 33, please list volume, page range, and year of publication.

Q10: For ref. 35, please list location of publisher.

Table 1: First column, row six: should the number after  $\pm$  begin with a decimal point (0.07)?