

Comparative Genetics of Type 1 Diabetes and Autoimmune Disease

Common Loci, Common Pathways?

Kevin G. Becker

Genome-scale analysis in type 1 diabetes has resulted in a number of non-major histocompatibility complex loci of varying levels of statistical significance. In no case has a specific gene been proven to be the source of genetic linkage at any candidate locus. Comparative analysis of the position of loci for type 1 diabetes with candidate loci from other autoimmune/inflammatory diseases shows considerable overlap. This supports a hypothesis that the underlying genetic susceptibility to type 1 diabetes may be shared with other clinically distinct autoimmune diseases such as systemic lupus erythematosus, multiple sclerosis, and Crohn's Disease. *Diabetes* 48:1353–1358, 1999

Type 1 diabetes is an autoimmune disease thought to arise through a complex interaction of genetic and environmental factors. Like many autoimmune diseases, type 1 diabetes is characterized by immune dysregulation that precedes overt clinical onset of the disease. These abnormalities include antibody disturbances as well as dysregulation in cellular immune responses (1–4).

Type 1 diabetes in humans and corresponding animal models have long been associated with specific alleles of the major histocompatibility complex (MHC) (2). This genetic region found at human chromosome 6p21.3 covers ~8.5 cM (5) and has been linked or associated with both disease susceptibility and resistance in many, if not most, autoimmune diseases. The MHC codes for a cluster of at least 30 (5) related class I and class II HLA genes. The MHC also contains many genes that are not related to HLA surface antigens. Specific alleles of genes found within the MHC have also been associated with susceptibility and resistance to infectious diseases in humans and in animal models (6).

The relationship of the MHC with immune/inflammatory diseases is thought to be due to the essential requirement of

HLA molecules in normal antigen recognition, processing, and presentation, ultimately leading to activation and progression of the immune system. This process is somehow perturbed in the disease state and has a combinatorial effect with other genetic modifiers and environmental triggers, leading to disease in some individuals. The genes, protein products, and biological pathways of the MHC associated with type 1 diabetes are central to normal immune function, are found in normal and affected individuals, and are not uniquely disease specific. Recently, a number of non-MHC loci have been linked to type 1 diabetes using a genome-wide scanning strategy. These loci have been given disease-specific provisional designations: *IDDM1* through *-15* (7). As described here, the genetic location of many of those provisional loci co-localize or overlap with loci from other autoimmune/inflammatory diseases, suggesting shared pathways in the etiology of clinically distinct autoimmune diseases.

APPROACHES IN THE STUDY OF TYPE 1 DIABETES

Molecular approaches in the study of type 1 diabetes can be categorized in two general ways: a traditional candidate gene approach and the more recent reverse-genetic approach. The candidate gene strategy is highly focused and logical. This is the classical hypothesis-driven approach and is usually focused on a small number of genes, proteins, and biological pathways of high information content and current critical importance. In the case of type 1 diabetes, this has often involved either immune specific regulatory proteins (4,8) or autoantigens specifically expressed on or related to the pancreas, such as insulin itself, islet cell autoantibody-69 (ICA69), GAD65, and IA-2 (2,9). This straightforward approach is well suited for the analysis of later-stage active disease processes. However, this strategy is based somewhat on prior knowledge and suffers from a significant lack of information (>95%) concerning the human genome sequence and complex regulatory pathways. It is inherently limited and subjective because of the critical need to focus on specific proteins or genes.

The more recent reverse-genetic approach in the study of type 1 diabetes typically involves genome-wide scans using affected sib pairs or multiplex families studied with a large set of polymorphic markers covering the entire human genome. This approach is objective, broadly focused, and statistically rigorous. Genome scanning methods have been used in the analysis of common complex multigenic diseases, including many autoimmune and infectious diseases, psychiatric dis-

From the Gerontology Research Center, National Institute on Aging, National Institutes of Health, Baltimore, Maryland.

Address correspondence and reprint requests to Kevin G. Becker, PhD, Gerontology Research Center, Room 4-101, 5600 Nathan Shock Dr., National Institute on Aging, National Institutes of Health, Baltimore, MD 21224. E-mail: beckerk@gre.nia.nih.gov.

Received for publication 26 March 1999 and accepted in revised form 5 April 1999.

LOD, logarithm of odds; MHC, major histocompatibility complex; SLE, systemic lupus erythematosus.

orders such as schizophrenia and bipolar illness, and hypertension, as well as animal models of complex human diseases. Limitations of genome-wide scans when applied to complex autoimmune diseases include: heterogeneity in disease phenotypes, population and ethnic differences, imperfect statistical and analytical models, and the use of different polymorphic marker panels and genetic maps. This has resulted, in some cases, in a lack of replication and confirmation at different loci among different groups studying the same disease. The typical result of a successful first-stage genome-wide scan ends with linkage to one or more genetic intervals of ~10–30 cM genetic distance. This is a huge distance at a molecular level, quite often with few biological clues as to the exact molecular basis of the linkage. This makes the next task of identifying and cloning the linked gene in that interval quite difficult.

Using the genome-wide scan approach, many groups have identified a number of candidate susceptibility loci for type 1 diabetes having varying levels of statistical significance: from suggestive to highly significant (10). These loci generally include the MHC, as well as numerous non-MHC candidate loci. Many of the candidate loci have suggestive significance values less than the traditional values (logarithm of odds [LOD] > 3.0) observed in monogenic diseases displaying Mendelian inheritance. Many of these candidate loci have been given a disease-specific designation (i.e., *IDDM1* through *-15*) as a temporary operational designation. While some of these loci have been confirmed, a number of loci have not been replicated (11).

Similar studies have been performed on other complex autoimmune/inflammatory diseases including multiple sclerosis (12–15), Crohn's disease (16,17), rheumatoid arthritis (18,19), psoriasis (20,21), systemic lupus erythematosus (22–25), and asthma (26), among others. Similar results have been found in these disease studies as well: multiple loci (5–10), most with suggestive levels of statistical significance (LOD 1–3), disease-specific designations (*IBD1*, *-2*, *-3*; *SLE1*, *-2*, *-3*, etc.), and a lack of replication of some loci in subsequent studies. In no case for the candidate loci identified using the genome-wide scan approach in the study of common autoimmune diseases, including type 1 diabetes, has a specific non-MHC gene been shown to be the source of the genetic linkage.

CO-LOCALIZATION OF AUTOIMMUNE DISEASE LOCI: A PATTERN IS EMERGING

Recently, it has been noted that the position of provisional loci found in type 1 diabetes co-localize or overlap with loci found in different autoimmune/inflammatory diseases (27–30). This is consistent with a hypothesis that, like the MHC, some of these provisional loci may involve common susceptibility genes or biochemical pathways that are central to normal immune function. These genes or pathways may contribute to immune dysregulation shown to be present in different autoimmune diseases, possibly before the onset of overt clinical symptoms (2). Table 1 shows the names and chromosomal locations of the candidate loci identified for type 1 diabetes. This includes loci given provisional disease-specific designations (*IDDM1*, *-2*, *-3*, *-4*, etc.), as well as other loci linked to type 1 diabetes. Also shown are loci from other human autoimmune/inflammatory disease genome-wide scans that co-localize, overlap, or are within at least 10 cM of a type 1 diabetes locus. Ten centi-

morgans is the approximate limit of resolution of a typical first-stage genome-wide scan. For example, *IDDM2* found at 11p15.5 (31) is found at the exact same position as loci for systemic lupus erythematosus (SLE) (23), ankylosing spondylitis (32), asthma (26), and multiple sclerosis (14). All four disease loci have been defined by the same polymorphic marker, D11S922, at 0.323 cM position on human chromosome 11 (5). A candidate gene at 11p15.5 is the insulin gene itself. Variable number of tandem repeat (VNTR) polymorphisms at the 5' end of the insulin gene have been linked to type 1 diabetes (32,33). One interpretation of this genetic linkage to insulin as a candidate gene is that insulin acts as an autoantigen (34). Another interpretation is that insulin or the tightly linked IGF-2 acts as a growth factor expressed in the thymus, promoting the immune response (35). Co-localization of multiple autoimmune diseases at this location suggests that whatever the exact gene is that is ultimately found at the locus *IDDM2*, it may play a broader role in autoimmune development.

As shown in Table 1, most (15 of 17, or 88%) of the provisional loci for type 1 diabetes identified to date have co-localized or overlapping loci found in other autoimmune or inflammatory diseases. In some cases, different autoimmune diseases are defined by the same genetic marker, including the loci *IDDM1*, *IDDM2*, *IDDM8*, *IDDM12*, *Xp11.4*, and *Xp11.1*. There has been no specific effort to study different autoimmune/inflammatory diseases using a standard panel of polymorphic markers to allow direct comparisons. This general pattern of locus co-localization is not found in human nonautoimmune disease (28). Three loci for type 1 diabetes (*IDDM5*, *-10*, *-15*) do not appear to have overlapping loci from other autoimmune studies. Data for *IDDM14* are not publicly available (7). This pattern of co-localization of loci from related immune diseases has been found in both human and animal studies and has been shown to be statistically significant (27,28). It has been proposed that co-localization of autoimmune loci may be due to common biological pathways shared among related autoimmune/inflammatory diseases in both human disease (28,36) and animal models (27,29,30,37).

THE IMPORTANCE OF SUBPHENOTYPING

In some cases, using genome-wide scans in animal models of autoimmune disease, the general disease phenotype has been broken down into individual loci having discrete subphenotypes. This has generally not been possible in human studies, and emphasizes the importance of animal models in autoimmune studies. This subphenotyping at specific loci can potentially provide clues to the functional significance of candidate genes at each locus (37). For example, in murine experimental allergic encephalomyelitis, Butterfield et al. (38) define independent quantitative trait loci with linkages to susceptibility, onset, severity, and duration of disease. In murine SLE, Morel et al. (39) identified independent loci for glomerulonephritis and anti-dsDNA antibody production. This subphenotyping in animal models allows greater functional characterization of contributing loci in complex autoimmune phenotypes and may provide useful biological information in the search for candidate genes at specific human type 1 diabetes loci.

Recently, Concannon et al. (7) identified a novel locus for type 1 diabetes (maximum LOD score = 3.31) at human chro-

TABLE 1
Names and chromosomal locations of candidate loci identified for type 1 diabetes

Loci	Chromosomal location	Marker	Reference	Overlapping autoimmune loci	Marker	Reference
<i>IDDM1</i>	6p21	D6S426	7	MS	D6S273	14
				MS	D6S273	13
				Asthma	D6S276	26
				CD	D6S276	17
				Ankylosing spondylitis	D6S276	32
				SLE	D6S426	23
				Coeliac disease	HLA-DQ	63
<i>IDDM2</i>	11p15.5	D11S922	31	SLE	D11S922	23
				Ankylosing spondylitis	D11S922	32
				Asthma	D11S96	26
				Multiple sclerosis	D11S922	14
<i>IDDM3</i>	15q26	D15S107	57	SLE	D15S127	23
				Ankylosing spondylitis	D15S127	32
				Coeliac disease	D15S642	64
				Coeliac disease	D15S207	63
<i>IDDM4</i>	11q13	FGF3	31	Asthma	FCER1B	26
		FGF3	57	EAE	(D7Mit37)	65
		FGF3	58			
		D11S1296	11			
<i>IDDM5</i>	6q25	D6S290	11	None		
<i>IDDM6</i>	18q21	D18S39	59	Rheumatoid arthritis	D18S57, D18S474	18
		D18S64	31			
<i>IDDM7</i>	2q31–33	D2S152	60	SLE	D2S1391	22
<i>IDDM8</i>	6q27	D6S264	31	SLE	D6S1027	22
		D6S281	11	Ankylosing spondylitis	D6S281	32
<i>IDDM9</i>	3q21	D3S1303	31	Rheumatoid arthritis	D3S1267	18
		(kdp1)	61	Multiple sclerosis	D3S1309	12
<i>IDDM10</i>	10p11–q11	D10S193	62	None		
<i>IDDM11</i>	14q24.3	D14S67	57	SLE	D14S74	23
				Graves' disease	D14S81	66
				Antibody response	(D12Mit27)	67
				SLE	D2S1391	22
<i>IDDM12</i>	2q33	D2S152	60	Thyroiditis	CTLA4	68
		CTLA4	7			
<i>IDDM13</i>	2q36	D2S301	7	Rheumatoid arthritis	D2S377, D2S2354	18
				Ankylosing spondylitis	D2S126	32
<i>IDDM14</i>	NA	NA		NA		
<i>IDDM15</i> <i>1q42</i>	6q21 1q42	D6S283	7	none		
		AGT	7	SLE	D1S103	40
		D1S1644		SLE	D1S3462	22
				SLE	D1S235	23
				Ankylosing spondylitis	D1S229	32
<i>Xp11.4</i>	Xp11.4	DXS1068	69	Rheumatoid arthritis	DXS1068	18
				Multiple sclerosis	DXS1068	12
<i>Xp11.1</i>	Xp11.1	DXS991	31	Multiple sclerosis	DXS991	13

All chromosomal positions are from Location Database (5). Identical markers are in bold. NA, not available.

mosome 1q at marker D1S1617. This locus co-localizes with loci for SLE (40) and ankylosing spondylitis (32). In human SLE, this locus is linked to serum levels of anti-chromatin antibody, and in mouse SLE to both anti-chromatin and anti-DNA antibody production (24,40–42). The co-localization of these three autoimmune diseases at this position suggests that this locus may be involved in a pathway that affects the quantitative regulation of antibody levels, and that this may contribute to the ultimate disease phenotype in all three diseases. Interestingly, the immune disorder Chediak-Higashi syndrome maps to 1q42 as well (43). The mouse equivalent of Chediak-Higashi is the beige locus. Mutations at the beige locus have been associated with severe immune abnormali-

ties (44). Comparative analysis across related diseases and across species, especially in the context of subphenotyping, may help in identifying the biological basis of each locus.

AN ANALOGY TO TUMOR BIOLOGY PATHWAYS

The process of tumorigenesis is thought to arise from a complex combination of environmental and genetic events. Generally, this involves familial or somatic mutations in sets of genes essential to critical pathways related in part to cellular growth control. Mutations in ubiquitously expressed genes combine with alterations in tissue-specific pathways, resulting in the overall phenotype in a specific tumor type. This combinatorial process quite often involves

shared general factors in basic cell cycle control (CDK4, p16, Rb1, WAF1), signal transduction (MET, ras, B-catenin, src), DNA repair (p53, ATM, MSH2, MLH1), and apoptosis (BAX, BCL2) (45). Different combinations of these general factors in concert with more tissue- and tumor-specific factors, such as BRCA1 and -2, PTC, WT1, NF1, and PML, lead to tumorigenesis in a given cell type (45). Each tumor is unique. It is the interaction and sequential expression of these factors in a given tumor that help define the tumor specific characteristics of aggressiveness, drug resistance, and metastatic potential.

Complex autoimmune diseases can be thought of as similar to cancer in that they are thought to involve environmental triggers in the context of complex sets of interacting susceptibility genes. The end-stage phenotype of a specific autoimmune disease may be clinically distinct and/or organ specific, while the etiology of many autoimmune diseases may involve shared processes of immune regulation (46). It is likely that basic pathways affecting proinflammatory/anti-inflammatory cytokine ratios, apoptosis, complex antibody regulation, effector T-cell populations, and hormonal control of the immune system are involved in these related diseases (46). Overlapping of provisional loci between type 1 diabetes and other autoimmune/inflammatory diseases suggests that, in some cases, common pathways may be involved, in part, in the etiology of clinically distinct autoimmune diseases. Environmental factors, a subset of genetic loci, or epigenetic mechanisms may contribute to disease or target-tissue specification.

SUMMARY AND IMPLICATIONS FOR THE STUDY OF TYPE 1 DIABETES

Genome-scale analysis of the genetics of type 1 diabetes has identified more than 15 provisional loci of varying levels of statistical significance. The next task is to assign biological significance to each genetic interval. This quite often involves painstaking analysis of polymorphisms in large populations, quite often with conflicting results. The association of the MHC with type 1 diabetes has been known for more than 2 decades (47), and although much has been learned about the biology of the MHC, the exact biological basis for the MHC linkage to type 1 diabetes is still unknown. This suggests that establishing the exact biological nature of genetic linkage to provisional loci for type 1 diabetes will be quite difficult.

The specific basis for co-localization of loci between multiple autoimmune diseases at each locus could be due to 1) the same allele of the same gene, 2) different alleles of the same gene, or 3) different members of a tightly linked complex of functionally related genes. Of course, for any given locus, co-localization of loci from different autoimmune diseases could be coincidental. Familial association of different autoimmune diseases in the same pedigree (48–50), co-occurrence of different autoimmune diseases, including type 1 diabetes, in the same individual (51–55), and shared clinical parameters of different autoimmune diseases (46,56) suggest a common biological basis in autoimmune/inflammatory disease. Co-localization and overlapping of candidate loci in autoimmune diseases, including type 1 diabetes, suggests that in some cases, common biological pathways may be involved in the etiology of type 1 diabetes and other clinically distinct autoimmune diseases.

ACKNOWLEDGMENTS

The author would like to thank Drs. L.W. Whitney, W. Biddison, and C. Sasaki for helpful comments.

REFERENCES

- Schranz DB, Lernmark A: Immunology in diabetes: an update. *Diabetes Metab Rev* 14:3–29, 1998
- Gottlieb PA, Eisenbarth GS: Diagnosis and treatment of pre-insulin dependent diabetes. *Annu Rev Med* 49:391–405, 1998
- Roep BO: T-cell responses to autoantigens in IDDM: the search for the Holy Grail. *Diabetes* 45:1147–1156, 1996
- Rabinovitch A, Suarez-Pinzon WL: Cytokines and their roles in pancreatic islet beta-cell destruction and insulin-dependent diabetes mellitus. *Biochem Pharmacol* 55:1139–1149, 1998
- Collins A, Frezal J, Teague J, Morton NE: A metric map of humans: 23,500 loci in 850 bands. *Proc Natl Acad Sci U S A* 93:14771–14775, 1996
- Hill AV: The immunogenetics of human infectious diseases. *Annu Rev Immunol* 16:593–617, 1998
- Concannon P, Gogolin-Ewens KJ, Hinds DA, Wapelhorst B, Morrison VA, Stirling B, Mitra M, Farmer J, Williams SR, Cox NJ, Bell GI, Risch N, Spielman RS: A second-generation screen of the human genome for susceptibility to insulin-dependent diabetes mellitus. *Nat Genet* 19:292–296, 1998
- Falcone M, Sarvetnick N: The effect of local production of cytokines in the pathogenesis of insulin-dependent diabetes mellitus. *Clin Immunol* 90:2–9, 1999
- Bach JF, Koutouzov S, van Endert PM: Are there unique autoantigens triggering autoimmune diseases? *Immunol Rev* 164:139–155, 1998
- Lander E, Kruglyak L: Genetic dissection of complex traits: guidelines for interpreting and reporting linkage results. *Nature Genet* 11:241–247, 1995
- Luo DF, Buzzetti R, Rotter JI, Maclaren NK, Raffel LJ, Nistico L, Giovannini C, Pozzilli P, Thomson G, She JX: Confirmation of three susceptibility genes to insulin-dependent diabetes mellitus: IDDM4, IDDM5 and IDDM8. *Hum Mol Genet* 5:693–698, 1996
- Ebers GC, Kukay K, Bulman DE, Sadovnick AD, Rice G, Anderson C, Armstrong H, Cousin K, Bell RB, Hader W, Paty DW, Hashimoto S, Oger J, Duquette P, Warren S, et al.: A full genome search in multiple sclerosis. *Nature Genet* 13:472–476, 1996
- Sawcer S, Jones HB, Feakes R, Gray J, Smaldon N, Chataway J, Robertson N, Clayton D, Goodfellow P, Compston A: A genome screen in multiple sclerosis reveals susceptibility loci on chromosome 6p21 and 17q22. *Nature Genet* 13:464–468, 1996
- Haines JL, Ter-Minassian M, Bazyk A, Gusella JF, Kim DJ, Terwedow H, Pericak-Vance MA, Rimmler JB, Haynes CS, Roses AD, Lee A, Shaner B, Menold M, Seboun E, Fitoussi RP, et al.: A complete genomic screen for multiple sclerosis underscores a role for the major histocompatibility complex. *Nature Genet* 13:469–471, 1996
- Kuokkanen S, Sundvall M, Terwilliger JD, Tienari PJ, Wikstrom J, Holmdahl R, Pettersson U, Peltonen L: A putative vulnerability locus to multiple sclerosis maps to 5p14-p12 in a region syntenic to the murine locus Eae2. *Nature Genet* 13:477–480, 1996
- Hugot J-P, Laurent-Puig P, Gower-Rousseau C, Olson JM, Lee JC, Beaugier L, Naom I, Dupas JL, Van Gossum A, Orholm M, Bonaiti-Pellie C, Weissenbach J, Mathew CG, Lennard-Jones JE, Cortot A, Colombel JF, Thomas G: Mapping of a susceptibility locus for Crohn's disease on chromosome 16. *Nature* 379:821–823, 1996
- Satsangi J, Parker M, Louis E, Hashimoto L, Kato N, Welsh K, Terwilliger JD, Lathrop GM, Bell JI, Powell D: Two stage genome-wide search in inflammatory bowel disease provides evidence for susceptibility loci on chromosomes 3, 7, and 12. *Nat Genet* 14:199–202, 1996
- Cornelis F, Faure S, Martinez M, Prud'homme JF, Fritz P, Dib C, Alves H, Barrera P, de Vries N, Balsa A, Pascual-Salcedo D, Maenaut K, Westhovens R, Migliorini P, Tran TH, et al.: New susceptibility locus for rheumatoid arthritis suggested by a genome-wide linkage study. *Proc Natl Acad Sci U S A* 95:10746–10750, 1998
- Remmers EF, Longman RE, Du Y, O'Hare A, Cannon GW, Griffiths MM, Wilder R: A genome scan localizes five non-MHC loci controlling collagen-induced arthritis in rats. *Nat Genet* 14:82–85, 1996
- Matthews D, Fry L, Powels A, Weber J, McCarthy M, Fisher E, Davies K, Williamson R: Evidence that a locus for familial psoriasis maps to chromosome 4q. *Nat Genet* 14:231–233, 1996
- Tomfohrde J, Silverman A, Barnes R, Fernandez-Vina MA, Young M, Lory D, Morris L, Wuepper KD, Stastny P, Menter A, Bowcock A: Gene for familial psoriasis susceptibility mapped to the distal end of human chromosome 17q. *Science* 264:1141–1145, 1994

22. Moser KL, Neas BR, Salmon JE, Yu H, Gray-McGuire C, Asundi N, Bruner GR, Fox J, Kelly J, Henshall S, Bacino D, Dietz M, Hogue R, Koelsch G, Nightingale L, et al.: Genome scan of human systemic lupus erythematosus: evidence for linkage on chromosome 1q in African-American pedigrees. *Proc Natl Acad Sci U S A* 95:14869–14874, 1998
23. Gaffney PM, Kearns GM, Shark KB, Ortmann WA, Selby SA, Malmgren ML, Rohlf KE, Ockenden TC, Messner RP, King RA, Rich SS, Behrens TW: A genome-wide search for susceptibility genes in human systemic lupus erythematosus sib-pair families. *Proc Natl Acad Sci U S A* 95:14875–14879, 1998
24. Morel L, Rudofsky UH, Longmate JA, Schiffenbauer J, Wakeland EK: Polygenic control of susceptibility to murine systemic lupus erythematosus. *Immunity* 1:219–229, 1995
25. Kono DH, Burlingame RW, Owens DG, Kuramochi A, Balderas RS, Balomenos D, Theofilopoulos AN: Lupus susceptibility loci in New Zealand mice. *Proc Natl Acad Sci U S A* 91:10168–10172, 1994
26. Daniels SE, Bhattacharyya S, James A, Leaves NI, Young A, Hill MR, Faux JA, Ryan GF, le Souf PN, Lathrop GM, Musk AW, Cookson WO: A genome-wide search for quantitative trait loci underlying asthma. *Nature* 383:247–250, 1996
27. Vyse TJ, Todd JA: Genetic analysis of autoimmune disease. *Cell* 85:311–318, 1996
28. Becker KG, Simon RM, Bailey-Wilson JE, Freidlin B, Biddison WE, McFarland HF, Trent JM: Clustering of non-major histocompatibility complex susceptibility candidate loci in human autoimmune diseases. *Proc Natl Acad Sci U S A* 95:9979–9984, 1998
29. Wu J, Longmate JA, Adamus G, Hargrave PA, Wakeland EK: Interval mapping of quantitative trait loci controlling humoral immunity to exogenous antigens: evidence that non-MHC immune response genes may also influence susceptibility to autoimmunity. *J Immunol* 157:2498–2505, 1996
30. Teuscher C, Rhein DM, Livingstone KD, Paynter RA, Doerge RW, Nicholson SM, Melvold RW: Evidence that Tmevd2 and eae3 may represent either a common locus or members of a gene complex controlling susceptibility to immunologically mediated demyelination in mice. *J Immunol* 159:4930–4934, 1997
31. Davies JL, Kawaguchi Y, Bennett ST, Copeman JB, Cordell HJ, Pritchard LE, Reed PW, Gough SC, Jenkins SC, Palmer SM, Ballfour KM, Rowe BR, Farrall M, Barnett AH, Bain SC, Todd JA: A genome-wide search for human type 1 diabetes susceptibility genes. *Nature* 371:130–136, 1994
32. Brown MA, Pile KD, Kennedy LG, Campbell D, Andrew L, March R, Shatford JL, Weeks DE, Calin A, Wordsworth BP: A genome-wide screen for susceptibility loci in ankylosing spondylitis. *Arthritis Rheum* 41:588–595, 1998
33. She JX, Marron MP: Genetic susceptibility factors in type 1 diabetes: linkage, disequilibrium and functional analyses. *Curr Opin Immunol* 10:682–689, 1998
34. Sarugeri E, Dozio N, Belloni C, Meschi F, Pastore MR, Bonifacio E: Autoimmune responses to the beta cell autoantigen, insulin, and the INS VNTR-IDD2 locus. *Clin Exp Immunol* 114:370–376, 1998
35. Vaffiadis P, Bennett ST, Todd JA, Grabs R, Polychronakos C: Divergence between genetic determinants of IGF2 transcription levels in leukocytes and of IDD2-encoded susceptibility to type 1 diabetes. *J Clin Endocrinol Metab* 83:2933–2939, 1998
36. Wu J, Edberg JC, Redecha PB, Bansal V, Guyre PM, Coleman K, Salmon JE, Kimberly RP: A novel polymorphism of FcγRIIIa (CD16) alters receptor function and predisposes to autoimmune disease. *J Clin Invest* 100:1059–1070, 1997
37. Teuscher C, Hickey WF, Grafer CM, Tung KS: A common immunoregulatory locus controls susceptibility to actively induced experimental allergic encephalomyelitis and experimental allergic orchitis in BALB/c mice. *J Immunol* 160:2751–2756, 1998
38. Butterfield RJ, Sudweeks JD, Blankenhorn EP, Korngold R, Marini JC, Todd JA, Roper RJ, Teuscher C: New genetic loci that control susceptibility and symptoms of experimental allergic encephalomyelitis in inbred mice. *J Immunol* 161:1860–1867, 1998
39. Morel L, Mohan C, Yu Y, Schiffenbauer J, Rudofsky UH, Tian N, Longmate JA, Wakeland EK: Multiplex inheritance of component phenotypes in a murine model of lupus. *Mamm Genome* 10:176–181, 1999
40. Tsao BP, Cantor RM, Kalunian KC, Chen CJ, Badsha H, Singh R, Wallace DJ, Kitridou RC, Chen SL, Shen N, Song YW, Isenberg DA, Yu CL, Hahn BH, Roter JI: Evidence for linkage of a candidate chromosome 1 region to human systemic lupus erythematosus. *J Clin Invest* 99:725–731, 1997
41. Morel L, Rudofsky UH, Longmate JA, Schiffenbauer J, Wakeland EK: Polygenic control of susceptibility to murine systemic lupus erythematosus. *Immunity* 1:219–229, 1994
42. Drake CG, Rozzo SJ, Hirschfeld HF, Smaroworawong NP, Palmer E, Kotzin BL: Analysis of the New Zealand Black contribution to lupus-like renal disease. Multiple genes that operate in a threshold manner. *J Immunol* 154:2441–2447, 1995
43. Barrat FJ, Auloge L, Pastural E, Lagelouse RD, Vilmer E, Cant AJ, Weissenbach J, Le Paslier D, Fischer A, de Saint Basile: Genetic and physical mapping of the Chediak-Higashi syndrome on chromosome 1q42–43. *Am J Hum Genet* 59:625–632, 1996
44. Barbosa MD, Barrat FJ, Tchernev VT, Nguyen QA, Mishra VS, Colman SD, Pastural E, Dufourcq-Lagelouse R, Fischer A, Holcombe RF: Identification of mutations in two major mRNA isoforms of the Chediak-Higashi syndrome gene in human and mouse. *Hum Mol Genet* 6:1091–1098, 1997
45. Fearon ER: Human cancer syndromes: clues to the origin and nature of cancer. *Science* 278:1043–1050, 1997
46. Rose NR: Autoimmune diseases-tracing the shared threads. *Hospital Practice* 32:147–154, 1997
47. Nerup J, Platz P, Andersen OO, Christy M, Lyngsøe J, Poulsen JE, Ryder LP, Nielsen LS, Thomsen M, Svejgaard A: HL-A antigens and diabetes mellitus. *Lancet* 2:864–866, 1974
48. Bias WB, Reveille JD, Beaty TH, Meyers DA, Arnett FC: Evidence that autoimmunity in man is a Mendelian dominant trait. *Am J Hum Genet* 39:584–602, 1986
49. Ginn LR, Lin JP, Plotz PH, Bale SJ, Wilder RL, Mbayu A, Miller FW: Familial autoimmunity in pedigrees of idiopathic inflammatory myopathy patients suggests common genetic risk factors for many autoimmune diseases. *Arthritis Rheum* 41:400–405, 1998
50. Grennan DM, Sanders PA, Thomson W, Dyer PA: Rheumatoid arthritis: inheritance and association with other autoimmune diseases. *Dis Markers* 4:157–162, 1986
51. Yamato E, Ikegami H, Kawaguchi Y, Fujisawa T, Hamada Y, Ueda H, Shintani M, Ogihara T: Insulin-dependent diabetes mellitus associated with autoimmune thyroiditis and rheumatoid arthritis. *Am J Med Sci* 313:64–66, 1997
52. Sattar MA, Al-Sughyer AA, Siboo R: Coexistence of rheumatoid arthritis, ankylosing spondylitis and dermatomyositis in a patient with diabetes mellitus and the associated linked HLA antigens. *Br J Rheumatol* 27:146–149, 1988
53. Lorini R, d'Annunzio G, Vitali L, Scaramuzza A: IDDM and autoimmune thyroid disease in the pediatric age group. *J Pediatr Endocrinol Metab* 9 (Suppl. 1):89–94, 1996
54. Midgard R, Gronning M, Riise T, Kvale G, Nyland H: Multiple sclerosis and chronic inflammatory diseases: a case-control study. *Acta Neurol Scand* 93:322–328, 1996
55. Warren SA: Why study relationships between diseases? Clues to the etiology of multiple sclerosis (MS) derived from family illness history. *Can J Public Health* 77 (Suppl. 1):92–97, 1986
56. Martin S, Kardorf J, Schulte B, Lampeter EF, Gries FA, Melchers I, Wagner R, Bertrams J, Roep BO, Pflutzner A: Autoantibodies to the islet antigen ICA69 occur in IDDM and in rheumatoid arthritis. *Diabetologia* 38:351–355, 1995
57. Field LL, Tobias R, Thomson G, Plon S: Susceptibility to insulin-dependent diabetes mellitus maps to a locus (IDDM1) on human chromosome 14q24.3-q31. *Genomics* 33:1–8, 1996
58. Hashimoto L, Habita C, Beressi JP, Delepine M, Besse C, Cambon-Thomsen A, Deschamps I, Rotter JI, Djoulah S, James MR, Froguel P, Weissenbach J, Lathrop GM, Juller C: Genetic mapping of a susceptibility locus for insulin-dependent diabetes mellitus on chromosome 11q. *Nature* 371:161–164, 1994
59. Merriman T, Twells R, Merriman M, Eaves I, Cox R, Cucca F, McKinney P, Shield J, Baum D, Bosi E: Evidence by allelic association-dependent methods for a type 1 diabetes polygene (IDDM6) on chromosome 18q21. *Hum Mol Genet* 7:1003–1010, 1997
60. Luo DF, Maclaren NK, Huang HS, Muir A, She JX: Intrafamilial and case-control association analysis of D2S152 in insulin-dependent diabetes. *Autoimmunity* 2:143–147, 1995
61. Yokoi N, Kanazawa M, Kitada K, Tanaka A, Kanazawa Y, Suda S, Ito H, Serikawa T, Komeda K: A non-MHC locus essential for autoimmune type 1 diabetes in the Komeda Diabetes-Prone rat. *J Clin Invest* 100:2015–2021, 1997
62. Reed P, Cucca F, Jenkins S, Merriman M, Wilson A, McKinney P, Bosi E, Joner G, Ronning K, Thorsby E: Evidence for a type 1 diabetes susceptibility locus (IDDM10) on human chromosome 10p11-q11. *Hum Mol Genet* 7:1011–1016, 1997
63. Zhong F, McCombs CC, Olson JM, Elston RC, Stevens FM, McCarthy CF, Michalski JP: An autosomal screen for genes that predispose to celiac disease in the western counties of Ireland. *Nat Genet* 14:329–333, 1996
64. Houlston RS, Tomlinson IP, Ford D, Seal S, Marossy AM, Ferguson A, Holmes GK, Hosie KB, Howdle PD, Jewell DP: Linkage analysis of candidate regions for coeliac disease genes. *Hum Mol Genet* 8:1335–1339, 1997
65. Baker D, Rosenwasser OA, O'Neill, Turk J: Genetic analysis of experimental allergic encephalomyelitis in mice. *J Immunol* 155:4046–4051, 1995
66. Tomer Y, Barbesino G, Greenberg DA, Concepcion E, Davies TF, and the ICGATD: Linkage analysis of candidate genes in autoimmune thyroid disease III: detailed analysis of chromosome 14 localizes Graves disease-1 (GD-1) close to multinodular goiter-1 (MNG-1). *J Clin Endocrinol Metab* 83:

- 4321–4327, 1998
67. Puel A, Groot PC, Lathrop MG, Demant P, Mouton D: Mapping of genes controlling quantitative antibody production in Biozzi mice. *J Immunol* 154: 5799–5805, 1995
68. Awata T, Kurihara S, Iitaka M, Takei S, Inoue I, Ishii C, Negishi K, Izumida T, Yoshida Y, Hagura R, Kuzuya N, Kanazawa Y, Katayama S: Association of *CTLA-4* gene A-G polymorphism (*IDDM12* locus) with acute-onset and insulin-depleted IDDM as well as autoimmune thyroid disease (Graves' disease and Hashimoto's thyroiditis) in the Japanese population. *Diabetes* 47:128–129, 1998
69. Cucca F, Goy JV, Kawaguchi Y, Esposito L, Merriman ME, Wilson AJ, Cordell HJ, Bain SC, Todd JA: A male-female bias in type 1 diabetes and linkage to chromosome Xp in MHC HLA-DR3-positive patients. *Nat Genet* 19:301–302, 1998