

# Thirty Years of Computational Autopoiesis: A Review

---

Barry McMullin

Artificial Life Laboratory  
Research Institute for Networks  
and Communications  
Engineering (RINCE)  
Dublin City University  
Dublin 9  
Ireland  
Barry.McMullin@dcu.ie

**Abstract** Computational autopoiesis—the realization of autopoietic entities in computational media—holds an important and distinctive role within the field of artificial life. Its earliest formulation by Francisco Varela, Humberto Maturana, and Ricardo Uribe was seminal in demonstrating the use of an artificial, computational medium to explore the most basic question of the abstract nature of living systems—over a decade in advance of the first Santa Fe Workshop on Artificial Life. The research program it originated has generated substantive demonstrations of progressively richer, lifelike phenomena. It has also sharply illuminated both conceptual and methodological problems in the field. This article provides an integrative overview of the sometimes disparate work in this area, and argues that computational autopoiesis continues to provide an effective framework for addressing key open problems in artificial life.

---

## Keywords

Autopoiesis, self-maintenance, self-repair, individuality

---

## I Preamble: A Personal Retrospect

I must begin by acknowledging my debt, both personal and intellectual, to Francisco Varela.

I met Francisco for the first time in 1991—at the very first European Conference on Artificial Life (ECAL). As co-organizer of the conference, he was, of course, rather busy, so I was quite diffident in approaching him. Yet he was immediately welcoming, and happy to take the time to sit and discuss with me my initial tentative exploration of the concept of autopoiesis.

Subsequently we corresponded at some length, and in 1992 he accepted an invitation to Dublin to participate in the workshop *Autopoiesis and Perception*, which I co-organized with my colleague Noel Murphy [32]. We continued to correspond and meet at intervals over the following years. I particularly remember a discussion over dinner one evening during the third ECAL (Grenada, Spain, in 1995), where he dazzled me not only with his ability to maintain three simultaneous conversations with different people, but to do so in three (or more?) different languages, switching continually between them.

In any case, that particular conversation laid the basis for a more intense correspondence over the following two years that culminated in an exhaustive (and exhausting?) re-examination of the original, computational model of autopoiesis [46]. This will be discussed in more detail below; for now it is enough to note that it ultimately resolved a number of difficult questions regarding the earlier work—but that this resolution was possible only because of Francisco's unfailing patience with my questions, and cheerful willingness to dredge both his memory and his files to locate critical contemporary records of the work.

Francisco Varela was a brilliant and original scientist. But my enduring memory is of Francisco the man: his enthusiasm, his infectious good humor, his idealism, and his sheer appetite for life. He is sorely missed.

## 2 The Idea of Autopoiesis

What does it mean for something to be alive? Why are some things alive, and some not? Is there only one way of being alive, or many different and distinct ways?

These are the most fundamental questions that should define biology—and thus, in turn, artificial life—as a distinct field of enquiry. Yet the answers are not straightforward, and certainly not the subject of clear consensus. With the demise of vitalism, it was recognized that living systems are made of the same *kinds* of things as the non-living. It follows that the distinctiveness of living systems must arise from the particular manner or modes of organization that they embody. But it has proved surprisingly difficult to articulate what this mode (or these modes) of living organization consist of.

The project of autopoiesis was motivated *inter alia* by the attempt to answer just such questions. It proposes that, at some level, and in some way, all living things share a common organization; and that this is characterized by two intimately entwined closure properties:

- Closure in *production*: the system is composed of components that give rise to processes of production that, in turn, collectively produce more of those same components.
- Closure in *space*: the self-construction of a boundary between the living self and the ambience in which it is embedded, yet from which it distinguishes itself.

The central claim of autopoietic theory is that autopoietic organization is necessary (though hardly sufficient) for the emergence of living phenomenology. If that claim is taken seriously, it naturally defines a particular program within artificial life: the attempt to realize autopoietic organization in artificial, especially computational, media. Indeed, the attempt to do just this has been an integral part of work on autopoiesis from its very inception. Accordingly, we will turn now to the historical setting and context for the formulation of autopoietic theory and its seminal casting in a specifically computational context.

## 3 Origins

... the simulation rapidly provided the results our intuition had led us to expect: the spontaneous emergence in this artificial bi-dimensional world of units which self-distinguished by means of the formation of a ‘membrane’, and which showed a capacity of self-repair. ... It is important to mention this article [46] here because it was the first publication on the idea of autopoiesis in English for an international public, which led the international community to take charge of the idea. In addition it anticipated what twenty years later would become the explosive field now called artificial life and cellular automata.  
—Varela [45, p. 414]

As recounted by Varela, the concrete idea of autopoiesis arose only slowly, and in particular circumstances of time and place. The development seems to have originated around 1967 [45]. Varela was then a student (or “apprentice”) under Humberto Maturana in the Department of Sciences at the University of Chile. Maturana was already well

known, with an international reputation for his work on the neurophysiology of vision (see, e.g., [13]); but he had started to develop a basic dissatisfaction with the idea of “information” as being key to understanding brain and cognition.

The problems here are profound, and are still by no means resolved. Nervous systems somehow contrive to imbue mere “information” with meaning and significance. Despite the continuing dominance of the information-processing paradigm of cognition, such signification cannot be intrinsic to the raw information. Indeed, the very basis of Shannon’s information theory is, precisely, the separation of information from meaning. So signification is something that is somehow generated through the autonomous, self-referring dynamics of a nervous system, constrained by its embedding in, and coupling with, the world it comes to know.

These ideas of autonomous circular organization in nervous systems were developed by Maturana over the following several years. While Varela’s personal trajectory took him to the USA at this time, the two continued to meet and collaborate, with each other and with Heinz von Foerster, in discussing these ideas. This culminated in the publication of Maturana’s article “Neurophysiology of cognition” [22]. Varela retrospectively identified this as a critical step (to which, as Maturana acknowledged, both Varela and von Foerster had contributed)—“. . . the indisputable beginning of a turn in a new direction” [45, p. 412]. In particular, it made the first connection between these issues of self-reference in nervous systems and corresponding, but even more basic, problems regarding the molecular constitution of the biological cell—which is to say, of life itself.

Varela returned to Chile in 1970, now as a colleague of Maturana in the Biology Department at the University of Chile. They deliberately focused on exploring the nature of the organization of the living organism. At this point the problem was becoming clear as that of understanding or characterizing the demarcation between living and nonliving systems—but to do so in a way that was abstracted away from the specific biochemical contingencies of any particular form of life. So we see already here exactly the distinction between “life as it is” and “life as it could be” that eventually provided the intellectual foundation for the field of artificial life [17].

By mid-1971, the word *autopoiesis* itself had been coined, and Varela and Maturana had outlined a *minimal* model or exemplar. By this was meant an imaginary, abstract, chemical world, incorporating just those species and reactions that would suffice to allow the constitution of a minimal autopoietic system. This would therefore serve to isolate, in a very precise and concrete way, just what was, and was not, being pointed at by the new concept.

Thus, the minimal model served an important expository purpose—to give concrete form to what is otherwise a rather difficult and unfamiliar idea, but to do so in a way that strips away the many complex distractions that any real biochemical system would additionally present. Moreover, then as now, no real living cell is yet described in sufficient detail for us to be able to definitively identify its constitution as autopoietic, in any case.

Yet, the minimal model served another, potentially much more important purpose: as a *test* of what had now become, not just a concept, but a *theory* of autopoiesis. Maturana and Varela conjectured that autopoietic organization was capable of giving rise to characteristically biological phenomenology—in particular, to that most primitive cellular phenomenon of “self-repair,” of the conservation of composite, macroscopic, cellular organization even as the microscopic constituents of the cell are continuously turned over.

Ideally, of course, such a test case might have been implemented in real, wet chemistry. But this is technically challenging and, in any case, runs the risk of reintroducing many distracting complications that are not relevant to the restricted question being

asked. So, instead, they hit on the idea of using a computer simulation and, in particular, the idea of a low-level, fine-grained (“individual-” or “agent-based”) simulation.

This methodological idea was undoubtedly already in the air. Varela himself described it as an “obvious step” [45, p. 414]. It drew explicitly on the pioneering work of John von Neumann—work carried out in the early 1950s, but properly published only a few years prior to Varela and Maturana’s work [47, 29]—and on the seminal development of the game of life by Conway et al. (in Gardner), reported just as Varela and Maturana were already working along these lines [12].

On the other hand, some distinctive and highly original innovations were also required. The basic idea of a discrete two-dimensional space, with local interactions, clearly derives from von Neumann. However, Varela and Maturana introduced the idea of directly implementing (quasi-)stable *particles* that can move across the lattice, rather than relying on the relatively complex and fragile emergence of patterns of underlying node states with this property (e.g., in the manner of gliders in the game of life). They also incorporated a deliberately stochastic dynamics, and inter-particle transformations or reactions—essential to anything motivated by the idea of realizing an abstract chemistry (as opposed to an abstract physics). From a contemporary perspective, writing in the pages of the *Artificial Life Journal*, these are now essentially standard techniques; but it is important to recognize the originality and insight that they represented in that much earlier era.

Next, the assistance of Ricardo Uribe of the School of Engineering at the University of Chile was enlisted, and, as reported in the opening quotation of this section, a computer simulation of the minimal model was created, which rapidly confirmed the qualitative expectations of Varela and Maturana.

By the end of 1971, then, the theory of autopoiesis in its most basic form—as a theory of the molecular organization of living cells—had been quite fully articulated, and tested in the form of a highly simplified (and *therefore* highly demanding) computer simulation. Two texts had been written: a comprehensive technical account, and a draft of a shorter summary article (including the simulation results).

Publication proved difficult. The extended version was rejected by at least five publishers and journals; the short article was submitted to several journals, including *Science* and *Nature*, with similar results. However, following a visit to Chile by Heinz von Foerster in mid-1973, the article was further revised and submitted to *BioSystems* (of which von Foerster was a member of the editorial board), and eventually published in mid-1974 [46]. Much more comprehensive publications have, of course, followed [44, 23, 24]; but, as the first English language publication of the theory of autopoiesis for an international audience, the 1974 *BioSystems* article remains a key reference, which continues to be widely read and cited. Further, as the first application of agent-based computer modeling demarcating the living from the nonliving state, it occupies a seminal position in the prehistory of what we now call artificial life.

#### 4 The Minimal Model

The original computational model of autopoiesis was presented in the form of a detailed, natural-language algorithm [46, Appendix]. This algorithm is reviewed and critiqued, in detail, in [27]. I will return to that critique in due course, but first it is important to have a general, qualitative view of the model and the phenomenology it is intended to exhibit.

The chemistry takes places in a discrete two-dimensional space. Each position in the space is either empty or occupied by a single particle. Particles generally move in random walks in space. There are three distinct particle types, engaging in three distinct reactions (see Figure 1):

- *Production*: Two substrate ( $S$ ) particles may react, in the presence of a catalyst ( $K$ ) particle, to form a link ( $L$ ) particle.
- *Bonding*:  $L$  particles may bond to other  $L$  particles. Each  $L$  particle can form (at most) two bonds, thus allowing the formation of indefinitely long chains, which may close to form membranes. Bonded  $L$  particles become immobile.
- *Disintegration*: An  $L$  particle may spontaneously disintegrate, yielding two  $S$  particles. When this occurs, any bonds associated with the  $L$  particle are destroyed also.

Chains of  $L$  particles are permeable to  $S$  particles but impermeable to  $K$  and  $L$  particles. Thus a closed chain, or membrane, that encloses  $K$  or  $L$  particles effectively traps such particles.

The basic autopoietic phenomenon predicted for this system is the possibility of realizing dynamic cell-like structures that, on an ongoing basis, produce the conditions for their own maintenance. Such a system would consist of a closed chain (membrane) of  $L$  particles enclosing one or more  $K$  particles. Because  $S$  particles can permeate the membrane, there can be ongoing production of  $L$  particles. Since these cannot escape from the membrane, this will result in the buildup of a high concentration of  $L$  particles. On an ongoing basis, the membrane will rupture as a result of disintegration of component  $L$  particles. Because of the high concentration of  $L$  particles inside the membrane, there should be a high probability that one of these will drift to the rupture site and effect a repair *before* the  $K$  particle(s) escape, thus reestablishing precisely the conditions allowing the buildup and maintenance of that high concentration of  $L$  particles.

A secondary phenomenon that may arise is the spontaneous establishment of an autopoietic system from a randomized initial arrangement of the particles.

Both these phenomena were reported and illustrated in the original article.

One specific weakness of this original model deserves mention here. It is that the  $K$  component is not itself produced by any reaction in the putatively autopoietic system. *Prima facie* this appears to violate the demand for closure in the processes of production. The article itself is somewhat confusing on this point. A “six-point key” is provided for determining whether any given entity is autopoietic. Point 6 of this at first appears to require that all components should, indeed, be produced by interactions among the components; but it immediately equivocates by allowing some exceptions if the relevant components “participate as necessary permanent constitutive components in the production of other components” [46, p. 193]. This does appear somewhat clumsy. On the other hand, there must surely be *some* exceptions allowed (specifically, covering the case of the  $S$  particles that are simply harvested from the environment). In any case, as will be discussed later, more recent elaborations of this original model have specifically allowed for production of the  $K$  particles [1], so there is no *fundamental* difficulty here.

## 5 Elaboration (I)

Autopoiesis grew and developed in many different directions after 1974. The *computational* approach was specifically pursued by Milan Zeleny, in a series of publications between 1975 and 1978 [53, 54, 48, 49].

Zeleny first re-implemented the original model in the programming language APL, and then explored a number of different elaborations of it. In particular, he reported such phenomena as growth, change in shape, oscillation in chemical activity, and self-reproduction of autopoietic entities [48, 49].

Prima facie these represent important demonstrations of progressively richer, lifelike phenomena within the framework of computational autopoiesis. However, there are some serious difficulties in assessing the significance of these results. The articles cited above do not detail the exact modifications of the original algorithm that gave rise to the new phenomena; but the qualitative descriptions suggest that, in at least some cases, the original constraints of local interaction, random motion of particles, and time-independent particle dynamics may have been rather arbitrarily relaxed. If that were so, the interest of these results would be seriously diminished—in the sense that the more complex “macroscopic” phenomena may have been, in effect, implicitly programmed into individual microscopic particles.

The ideal way to address these questions would be, of course, to directly examine the software responsible. Indeed, this software was made available for use by interested researchers at the time [50, Note 11], although it is not entirely clear whether this was in a form that allowed inspection of the source code. In any case, and unfortunately, it appears that no copies of this software have survived.<sup>1</sup> It is therefore now impossible to come to any more definitive conclusions. However, this very impasse clearly suggests a lesson for the methodology of scientific research where the phenomena under investigation have their primary realization in the form of computers executing specific programs. I will return to this important methodological question below.

## 6 Diversification

Maturana and Varela themselves continued to explore the general paradigm of autopoiesis, of course. Milestones included the publication of *Autopoiesis and cognition: The realization of the living* in 1980 [23] and the more popular and accessible account, *The tree of knowledge: The biological roots of human understanding* in 1987 [24]. Also significant was the appearance, in the early 1980s, of two collections of articles by diverse authors, both edited by Milan Zeleny [51, 52]. These demonstrated a broadening interest in autopoiesis from researchers in a range of fields.

In the late 1980s there was also a return to the question of a minimal model of (molecular) autopoiesis—but this time not via computation but through real, wet chemistry. This direction of research was initially introduced in 1988, in a collaboration between Varela and Luisi [20], and quite quickly led to the exhibition of concrete chemical models [19].

Over that same period, from about 1988 to 1992, a strong philosophical interest in and critique of autopoietic ideas emerged in the literature of general systems theory. This was particularly associated with the work of Fleischaker [4–6]. A special issue of the *Journal of General Systems*, edited by Fleischaker in 1992 [7], provided a forum for a robust debate, from a variety of specialists, over the scope and, indeed, the precise meaning of autopoiesis.

For my purposes here, the most significant aspect of this discussion is the question of possible spaces, or *domains*, in which autopoiesis can be realized—and especially the status of purely computational implementations. Fleischaker has the merit of taking a very clear position on this: she holds that, as least insofar as autopoiesis is taken to be definitional or criterial for “life,” it must be interpreted as restricted “entirely to the physical domain” [5, p. 43]. In turn I take her to mean that computational models of autopoiesis must be regarded only as simulations, and not realizations, of life.

I am not fully persuaded by this position. In particular, it seems to me that we should distinguish between a purely formal *description* of a computation (e.g., the original algorithm of Varela et al. [46], or even a specific program implementing this algorithm)

<sup>1</sup> Milan Zeleny, personal communication, 1994.

and a concrete, physical, *computer* that is actually executing such an algorithm. The latter is clearly a very different kind of system from a conventional chemical or molecular one; but, equally, it is certainly still a *physical* one. If it hosts autopoietic entities—albeit removed by a significant number of levels from the underlying hardware—then those entities are surely also still *physical*. Thus, I would find it hard to categorize such an entity as “not living” *solely* on a putative basis of its being “not physical.”

On the other hand, I am also unconvinced that debate on this issue will be particularly fruitful in any case. I would argue that—within the field of artificial life at least—the substantive issue at the current time is not whether a computationally based autopoietic entity is “really” alive, but rather whether we can exhibit computationally based autopoietic entities with substantially richer, more lifelike phenomenology. The early work of Zeleny was arguably developing in this direction; but, as we have seen, the concrete implementations of that work are no longer available, and it is now very difficult to appraise its substance critically.

I will therefore now return to consider a second phase of (attempted) elaboration of the phenomenology of computational autopoiesis.

## 7 Conjecture and Refutation

As far as I am aware, following Zeleny’s initial explorations, little further substantive development of *computational* autopoiesis was published between c. 1980 and the early 1990s.

As recounted in the preamble, my personal interest began around 1991. I was then completing a detailed reevaluation of von Neumann’s original work on “self-reproducing” automata. My conclusion from that study was that this could be best understood as focused not on self-reproduction per se, but on self-reproduction as a mechanism for the evolutionary growth of complexity [29]. Indeed, von Neumann had demonstrated how the architecture of a passive genetic *description tape* coupled with a general purpose, programmable *constructor* would give rise to a dense network of mutational pathways connecting relatively simple automata to arbitrarily complex ones. This is a very significant result; but it does not, in itself, suffice to realize the open-ended evolutionary growth of complexity in artificial systems.

The critical obstacle (one recognized clearly by von Neumann himself) is that his automata are extremely *fragile*. While they are logically capable of self-reproduction, they can do so only in a completely empty environment, with no disturbance to their operation. In practice, of course, Darwinian evolution could only happen within ecosystems where automata interact both with each other and with a more or less complex environment. Thus, to have a serious prospect of open-ended evolutionary growth of complexity, one requires automata that not only have a von Neumann style genetic architecture but are also capable of surviving—maintaining and repairing themselves—in the face of environmental perturbation. Indeed, they should do this even while being open to turnover of their components (the latter being necessary to harvest materials for self-reproduction, if for no other reason).

Now von Neumann had exhibited—in detail—how to realize automata with his genetic architecture within a comparatively simple discrete two-dimensional space. Similarly, Varela et al. had exhibited how to realize automata capable of basic self-repair and turnover of components in a similar space (or, at least, one explicitly inspired by von Neumann’s). The question therefore arises whether one can engineer a single artificial universe, hopefully of not much greater inherent complexity, in which these two kinds of automaton can be unified.

This then was the problem I started to discuss with Varela in the early 1990s [25]. My starting point was the original computational model of autopoiesis. The research

program was, initially at least, rather simple: reimplement that model, demonstrating the original minimal autopoietic phenomenology; and then explore elaborations (more particle types, more reactions, etc.) in the general direction of trying to unite the autopoietic functionality with something like a von Neumann genetic system.

However, this program failed at the very first hurdle: recreating the original, minimal, autopoietic phenomenology. It took some time, and several different attempts (which became intermittent, over a number of years), but it gradually became clear that the difficulties were more than superficial.

Eventually, in the autumn of 1996, I had the opportunity (with support from Chris Langton at the Santa Fe Institute) to make a concentrated study of the problem. This resulted in a detailed analysis of a number of anomalies in the original *BioSystems* article [46]. These encompassed both difficulties in interpreting the algorithm, and apparent inconsistencies between the algorithm and the presented experimental results.

While a significant number of detailed problems with the early computational model were uncovered in this process, the great majority of these were relatively minor, and could have been ignored or easily corrected. However, there was one phenomenon that was more serious and was responsible for the fundamental difficulty in recreating the original autopoietic phenomena. That was the problem of *premature bonding* of the  $L$  particles. According to the original algorithm, once  $L$  particles form, they can spontaneously bond together; and further, once bonded, they become immobile. But this is a fatal flaw for the supposed autopoietic closure of the reaction network: the free  $L$  particles are effectively sequestered by the side reaction of spontaneous bonding. This makes them essentially unavailable for membrane repair, and also clogs up the interior of the cell, eventually blocking further production.

Figures 1 and 2 show an experimental example of this. Figure 1 shows the initial configuration of an externally created cell. Figure 2 shows the state of the cell after 110 time steps. The membrane has not yet suffered any decay. However, the interior of the membrane is now completely clogged with bonded—and thus immobile— $L$  particles. Only two open positions remain inside the membrane, one occupied by the  $K$  particle. Since the production reaction requires two  $S$  particles adjacent to each other and to the  $K$  particle, there is no longer any available site for further production within the membrane, and further production of  $L$  particles is impossible. It follows that, whenever the membrane does eventually rupture, there will be no mobile  $L$  particles available to effect a repair.

This is a very robust failure. It is not sensitive to any precise details of the implementation. Indeed, the same problem was retrospectively identified in two other, independent attempts at re-implementation [15, 34], and may well have been present in Zeleny's work also (but superficially masked there through the introduction of nonlocal interactions, etc.).

This then provided the basis for a more extended and detailed discussion with Francisco Varela. But, of course, there had been a long lapse of time since the original experimental work had been carried out, c. 1971. Further, it should be remembered that the work had taken place against a backdrop of great turbulence in Chile. Varela had been a prominent supporter of the Allende government. Following the *coup d'état* and the coming to power of Pinochet in September 1973, Varela, like many others, was effectively forced into exile [45]. These events coincided with the development of the *BioSystems* article. Circumstances had therefore combined to make it very difficult to recall or reconstruct technical details of that work some 25 years later.

This was the situation of the discussion in late 1996 when chance intervened. Varela was now based in Paris, and there he happened to receive a shipment of papers and files, apparently belonging to him, which had just been located back at the University of Chile. Among these papers, he identified several contemporary, albeit fragmentary,




Figure 1. Initial configuration of autopoietic cell in 2D computational model. Small hollow squares are substrate (S); solid disk is catalyst (K); large hollow squares with bonds are membrane link (L) particles.

records of the early computational exploration of autopoiesis. Critically, these included an early discursive description of the model and a listing of a version of the actual computer code (in FORTRAN IV).

With the help of these documents it was finally possible to identify, with reasonable confidence, exactly the causes of the discrepancies that had been identified in the *BioSystems* article.<sup>2</sup> In particular, an explanation was found for the control of premature bonding in the original model. It transpired that the model had actually included an additional, undocumented interaction: chain-based bond inhibition. The effect of this was that free L particles in the immediate vicinity of the membrane (or, indeed, any chain of L particles) were inhibited from spontaneously forming bonds with each other. *Provided* the cell is not very big, this is sufficient to prevent premature bonding and ensure a continuing supply of free, mobile L particles to repair membrane ruptures as and when they arise. Once this interaction was reintroduced, the original autopoietic phenomenology of ongoing self-repair, as reported in [46], was easily realized again. Detailed experiments documenting and demonstrating this were presented at ECAL 1997 [33].

The implicit limitation of the chain-based bond inhibition mechanism to small cells should be carefully noted: it reveals, in particular, that this original form of autopoietic

<sup>2</sup> A comprehensive account, including the original critique, copies of the rediscovered contemporary documents and source code, and the discrepancy analysis, has been made available through the *Working Papers* series of the Santa Fe Institute [27]. A more informal account of the unraveling of this puzzle is also available [26].




Figure 2. Illustration of premature bonding.

cell will *not* easily support growth, or, more particularly, cell reproduction by fission. This is obviously rather important for any projected extension to realizing Darwinian evolution of such cells. We will return to this limitation below.

While this eventual recreation of the original autopoietic phenomenology ultimately hinged on a rather mundane technical refinement, I would argue that it was still a useful, if modest, contribution, for two distinct reasons:

- As already noted, the 1974 *BioSystems* article occupies a pivotal position in the history of the theory of autopoiesis, and continues to be widely referenced. Within that article, the computational model is critical, both to the exposition of the concept and to testing the conjecture that the phenomenon of dynamic self-maintenance of organization (in the face of component turnover), can be achieved by even a minimal autopoietic system. Defects in the reporting of that model—and thus in the reproducibility of the phenomena described—could therefore undermine the credibility of the core theory. Given that the main problem was actually the result of a relatively minor technical oversight—and *not* indicative of any substantive flaw in the core theory—it is important to clearly correct the historical record.
- Perhaps more importantly, to the extent that the technical basis for the phenomenology of the original model was obscure, or flawed, this presented a significant barrier to further study or elaboration of this approach.

It should be admitted that, in contrast to the other interactions in this model, chain-based bond inhibition has a distinctly ad hoc flavor. That being the case, it can reasonably be asked why this interaction should be considered as being acceptable—that is, as not detracting from the significance of the reported phenomena. This is particularly moot in that certain other (conjectured) interactions in the later models of Zeleny were treated more critically above (Section 5). However, it seems to me that a distinction should be drawn here between interactions that are ad hoc simply in the sense of “unfamiliar” or “unphysical,” and interactions that are nonlocal (in some relevant space) or that have preprogrammed time variation. As already noted, the latter have the clear potential to allow both aggregate and time-structured phenomena to be effectively programmed in (either in single agents, or in the environment as a whole). For example, apparently macroscopic cellular organization might, in effect, be an artifact constructed and coordinated by a single central control agent; in which case its “emergence” would presumably be of little interest.

By contrast, chain-based bond inhibition is still a perfectly local and time-independent interaction, occurring on a smaller scale than, and uninfluenced by, macroscopic cellular organization. It is for this reason that, in my view, the identification of this additional interaction does not undermine the significance of the dynamic, macroscopic, cellular organization originally reported for this model.

Of course, the question at stake here—of choosing appropriate abstractions or axiomatizations of primitive agents—is a difficult one, which has been at the heart of the field of artificial life right since its inception. Indeed, in a lecture originally dating from 1949, von Neumann captured this tension explicitly and succinctly:

... one may define parts in such numbers, and each of them so large and involved, that one has defined the whole problem away. If you choose to define as elementary objects things which are analogous to whole living organisms, then you obviously have killed the problem, because you would have to attribute to these parts just those functions of the living organism which you would like to describe or to understand. So, by choosing the parts too large, by attributing too many and too complex functions to them, you lose the problem at the moment of defining it. [47, p. 76]

This continues to be a matter of debate in the field, and I do not suggest that the particular demarcation I have suggested above—based specifically on locality and time independence—offers anything more than a very rough, heuristic guide. Particular models and their associated phenomenology must still be critically evaluated on their individual merits.

In closing this discussion of the clarification of the original model of computational autopoiesis it is, perhaps, equally important to comment on something that this work did *not* demonstrate: namely, it certainly did not show that there was, or is, any *obstacle of principle* in realizing autopoietic organization in an artificial computational medium. This must be stated clearly, as precisely an opposite interpretation has recently been attributed to this work:

... The non-computability of Autopoietic systems, as advanced here, apparently collides with the simulation results involving tessellation automatons [*sic*] [46]. But new versions of this simulation show that the original report of computational autopoiesis was flawed, as it used a non-documented feature involving chain-based bond inhibition [33]. Thus the closure exhibited by tessellation automatons is not a consequence of the “network” of simulated processes, but rather an artifact of coding procedures [33]. —Letelier et al. [18, p. 270]

For context, the overarching conjecture being advanced by Letelier is that autopoiesis is, in some fundamental sense, a “non-computable” process. By this they mean that while it can (obviously?) be realized in real physico-chemical form, there is some essential aspect of this process that cannot, even in principle, be realized in a strictly computational system. It amounts to a specific denial of the Church-Turing thesis (regarding the ultimate computability of the entire physical universe). But clearly there is a logical conflict between such a conjecture and the original illustration of the very concept of autopoiesis via a perfectly computational model. The quotation above is an attempt to resolve this conflict.

However, this is a most peculiar analysis.

It is true, of course, that the original report of computational autopoiesis was “flawed,” and, indeed, that this was primarily because “it used a non-documented feature involving chain-based bond inhibition.” However, the citation for the last sentence in this quotation is simply wrong: there is no reference whatsoever in [33] to any distinction between “simulated processes” and “coding procedures.” Nor can I conceive, even now, in retrospect, of any basis for assigning a different status to “chain-based bond inhibition” from that assigned to any other interaction in that particular artificial chemistry (production, bonding, absorption, etc.).

Thus, there is no basis whatever for supposing that this clarification of the “original report” undermines the essential result. The only flaw was in enumerating the processes or interactions underlying them: one necessary interaction was omitted from this description. The phenomena originally reported were accurate, and ultimately reproducible in perfectly computable form. If it is accepted that the phenomena *reported* in the original article constituted autopoiesis—as it apparently is by Letelier—then they *still* constitute autopoiesis. This was stated most explicitly in the conclusion of [33]:

It should be emphasised that the substantive point of this paper is to correct the historical record. . . . However, this correction does not add to, or modify, the original conceptual foundation of autopoiesis in any significant way.

I conclude that the overall thesis of Letelier et al.—of the non-computability of autopoietic systems—should be taken as *refuted*, rather than corroborated, by the results of [33].

## 8 Lessons for Methodology?

I have critically reviewed two distinct bodies of work in computational autopoiesis above—the original model of Varela et al., and the developments of that by Zeleny. There are certain similarities in my comments. In both cases I have identified anomalies or difficulties of interpretation of the published results. However, the final conclusions are quite different. In the case of the original model, it proved possible to locate the source code (or, at least, a close relative) on which the results were based; as a result, the substantive difficulties were fully resolved. For Zeleny’s work, unfortunately, the source code is apparently lost; definitive resolution of questions relating to it, at this stage, is therefore extremely difficult, if not impossible.

I should emphasize that these differing outcomes are largely due to chance—the fortuitous rediscovery of the Varela documents—rather than design. Regardless of that, however, they suggest a methodological lesson worth digesting. At the time of the original publications considered above, the technological facilities were not generally available to support easy distribution or access to accompanying code—but this is no longer the case. I would suggest therefore that as a general principle, published reports on computer models of ALife should be accompanied by access to the program code for

the models on the World Wide Web; and indeed, that practice has now been followed in the cases of the surviving documents from Varela et al. [27] and the successful modern reimplementation [28, 33].

## 9 Elaboration (II)

### 9.1 Individuation

As already explained, a natural research program in computational autopoiesis is to attempt to realize Darwinian evolution among lineages of autopoietic cells. This requires—among other things—the realization of cells that are at least capable of self-reproduction through simple growth and fission.

In pursuing this program, an initial problem that arises then is *individuation*.

My own exploration of this problem [30] begins with a conceptual comparison of autopoiesis and Stuart Kauffman's idea of a *collectively autocatalytic set* [16]. I conclude that there is a critical distinction between these, precisely in that autopoiesis requires self-generated *individuation*. This is classically captured by stating that the autopoietic network of processes must give rise to a *boundary*; however, it turns out that it is not entirely clear what should qualify, in general, as a boundary. In [30] I suggest a heuristic test for this, which essentially asks whether two putatively individual cells, in direct contact with each other, can reliably maintain their separate identities.

This test is, of course, motivated directly by the issue of self-reproduction. In the situation of self-reproduction of an autopoietic cell by growth and fission, the parent and offspring will clearly initially be in direct proximity to each other. If they fail to maintain their separate identities—if they can as easily merge back into one cell—this will hardly qualify as self-reproduction in the functional sense required for evolution, as such merging would undermine the very Malthusian population growth required for Darwinian selection.<sup>3</sup> This then is precisely the acid test for individuation: whether an entity can maintain its self-identity when it interacts with another entity of exactly the same material components and organization.

The test was first applied to a variety of classical artificial life systems: the  $\alpha$ -universes [14], Alchemy [8, 9], and Tierra [42]. In these cases, the conclusion was as expected: it served to make more precise the a priori judgment that none of these should be considered as realizing computational autopoiesis.

The surprise came when the test was applied to the original (re-implemented) model of computational autopoiesis [46, 33]. Though experimental results were not presented in detail, it appears that the chain-based bond inhibition reaction, while essential to the self-repair of a single isolated cell, also has the unintended side effect of inhibiting maintenance of the bounding membrane when two membranes are adjacent to each other. This means that, if anything, adjacent agents tend positively to merge rather than to maintain their individuality. In turn, if the proposed heuristic test is accepted as an operational test for autopoietic individuation, we are forced to the somewhat controversial conclusion that the original minimal model, which was intended precisely to be an exemplar of the concept, is not in fact properly autopoietic.

However, to forestall misinterpretation, I would emphasize that this result still does not argue for any obstacle *in principle* to computational autopoiesis. It merely demonstrates that robust autopoietic organization is not easy or trivial to achieve.

<sup>3</sup> It may be noted that there is an interesting "bootstrapping" problem here. Darwinian evolution requires fairly well defined individuals; but the development of suitable individuation mechanisms may need an extended evolutionary process. The resolution of this is likely to involve incremental and mutually reinforcing improvements in individuation and evolvability. An excellent discussion of the evolution of individuality in conventional biology is provided by Buss [2].

## 9.2 Direct Descendants

Two direct elaborations of the original model of computational autopoiesis have been published more recently.

Breyer et al. [1] first relaxed the original restrictions on the motion of bound  $L$  particles to allow the formation of flexible chains and, indeed, membranes. Allowing multiple, doubly bonded  $L$  particles to occupy a single lattice site further enhanced membrane motion. Adopting more sophisticated bond rearrangement interactions then allows chain fragments formed within the cell to be dynamically integrated into the membrane. These mechanisms together obviate the need for the chain-based bond inhibition mechanisms—since now, even if  $L$  particles spontaneously bond in the interior of the cells, the oligomers so formed remain mobile and, through bond rearrangement, can still successfully function to repair membrane ruptures. In this way, the model is reported as successfully supporting cell growth.

The authors separately consider mechanisms for production of further  $K$  particles within a cell. As previously noted, this possibility was absent in the original model, but is, of course, a necessary condition for any development toward self-reproduction. However, the full achievement of self-reproduction is not reported.

Rather similar developments were independently reported by McMullin and Groß (Chu) [31]. Again, bound  $L$  particles were allowed to move, facilitating flexible membranes; a somewhat more complex bond rearrangement interaction was introduced to facilitate cell growth; and the chain-based bond inhibition interaction was discarded. In this case, however, instead of simply permitting spontaneous  $L$  bonding within the cell (and then relying on mobility of these fragments to sustain repair), chain *initiation* (i.e., bonding between two free  $L$  particles) was disabled. This meant that the interior of the cell could still sustain a relatively large concentration of highly mobile free  $L$  particles. Other refinements included “smart” repair of the membrane (where membrane particles were replaced as soon as they started to decay—even before the membrane was actually ruptured) and “membrane affinity” of free  $L$  particles. The latter meant that the membrane tended to take a bilayer form—the outer layer particles being bonded, and the inner layer providing a stock of free  $L$  particles to effect very efficient repair.

These modifications together permitted the establishment of autopoietic cells with much greater dynamic stability (longer lifetimes) and/or cells capable of sustained growth. However, it should be noted that, with chain initiation disabled, this particular model would not be capable of exhibiting spontaneous cell generation *de novo*. Again, this article identifies the enriched phenomenology as potential steps toward the realization of self-reproduction of autopoietic cells, through growth and fission; but does not report the achievement of that.

## 9.3 Lattice Artificial Chemistry

A significantly different approach to computational autopoiesis, termed *lattice artificial chemistry* (LAC), has been recently pioneered in a series of studies by Ono and Ikegami.

LAC differs from the original model of Varela et al. [46] (and its direct descendants above) in adopting a *coarse-grained* approach. In the original model, each lattice site can generally be occupied by at most one particle: it is fine-grained to the single-molecule level. In LAC, by contrast, each single lattice site is permitted to host many particles, of many distinct molecular species.

Using this approach, Ono and Ikegami have progressively reported:

- Bounded, self-maintaining (autopoietic) cells in one dimension; reproduction of these cells through two distinct mechanisms [37].

- Bounded, self-maintaining (autopoietic) cells in two dimensions; reproduction of these cells [36, 38]. A particular innovation here was the introduction of anisotropic repulsion among one particle species. This is critical to the formation of thin films, which can function as membranes.
- Variant strains of two-dimensional cells are introduced, based on a variety of individually autocatalytic particles, with different rates of membrane particle generation [39]. Mutation between the varieties of autocatalysts is possible. Experiments are reported in which there is selection at the proto-cell level, favoring cells with the higher-productivity autocatalysts, and an optimized cell size.
- Most recently, a generalization to a three-dimensional space has been described (with Madina [21]).

Because LAC uses the coarse-grained approach, direct comparisons with the fine-grained models are difficult. For example, when multiple cells are present in the same space, they normally share membranes, so that a population of cells exhibits a honeycomb structure. It is difficult to assess the degree of individuation on the part of single cells in this case (as multiple cells contribute to the maintenance of any given boundary).

The exhibition of selection at the cell level seems particularly significant [39]. At this point, this is still limited to selection among a predesigned variety of cell strains—there is nothing like a von Neumann style genetic architecture, or programmable constructor. Nonetheless, the LAC methodology clearly presents a very fruitful avenue of research.

## 10 Related Developments

While this article has concentrated on the development of computational autopoiesis *per se*, this is clearly related to several other, parallel developments.

The *chemoton* of Tibor Gánti [10], though apparently developed quite independently, is clearly related to cellular autopoiesis; indeed, this first (English) report on the chemoton concept appeared in the same journal, *BioSystems*, as the first presentation of autopoiesis [46], but in the following year (1975). These parallels—and links to a wide variety of much earlier work—have been more recently reviewed by Gánti [11].

The chemoton is again a proposal for an abstract, minimal cell, consisting of a collectively autocatalytic network of reactions that is enclosed within a membrane that is also generated and maintained by the reaction network. It differs from the minimal autopoiesis model in explicitly including a *genetic subsystem*. It is also rather more detailed in its analysis of the required chemical dynamics (kinetics, etc.), and aims at supporting self-reproduction by growth and fission even in the minimal version.

While the provision of a genetic system is significant, it should be emphasized that, in the basic chemoton, this is limited to replication of varieties of *polymers*, which then have direct chemical effects. The polymers do *not* participate in a translation process; thus, this subsystem is again limited in scope compared to a von Neumann style genetic architecture, or programmable constructor.

The chemoton has been subject to various computer simulation studies, particularly by Csendes [3]. However, these were based essentially on an ordinary differential equation approach, rather than being molecular or agent-based. This means that, in particular, the distributed spatial dynamics of membrane growth, fission, and individuation were not substantively modeled.

Other computational systems with some connection or similarity to autopoiesis include *Alchemy* [8, 9], the  $\alpha$ -universes [14], *Coreworld* [41], and *Tierra* [42]. However,

as already mentioned, while these do involve closed, self-sustaining reaction networks, they do not have mechanisms for the self-generation of spatial boundaries.

Another very interesting line of work involves much more physico-chemically realistic computer models of minimal cellular structures. Rasmussen and colleagues, in particular, have recently exhibited spontaneous emergence of such hierarchical levels of organization [40]. This line of attack poses some scaling difficulties, as the computational demands of physico-chemical realism rise very rapidly. On the other hand, the cost of computation continues to fall, so this may well be a very fruitful domain of further research in the immediate future.

Finally, we should note that there are clear conceptual connections between the idea of autopoiesis and Robert Rosen's metabolism-repair systems and "closure under efficient causation" [43]. This connection has recently been made explicit by Letelier et al. [18], with the concrete suggestion that Rosen's work may provide an appropriate *formal* framework for the understanding of autopoiesis. A similar conjecture has also been recently presented by Nomura [35].

This approach immediately encounters the difficulty that Rosen explicitly argued that his form of closure could not, even in principle, be embedded within a purely computational system. By contrast, computer realization has been an explicit exemplar of autopoietic closure from the very start. As already discussed, Letelier et al. [18] have attempted to resolve this difficulty by interpreting [33] as showing that computational autopoiesis was fundamentally flawed from the beginning; but this is not an interpretation I can share. Accordingly, at this point, while I would agree that Rosen's work presents an interesting avenue for further exploration and contrast with autopoiesis, it is certainly not yet demonstrated that it can contribute a substantive, formal, *theory* to underpin it.

## II Conclusion

The idea of autopoiesis, and especially its computational realization, has proven very fruitful in its development and elaboration over the last thirty years. As we have seen, it represents a core thread in the history of what we now call artificial life, linking back directly to von Neumann's founding work on evolutionary automata.

We now have a comparatively clear formulation of several key open problems. One very general challenge is to articulate the relationship between the fine-grained and coarse-grained models of computational autopoiesis, and the relationship between both of these and work in wet artificial life. More specific, concrete challenges revolve around the exhibition of substantive Darwinian evolutionary phenomena among lineages of artificial (computational) autopoietic individuals. Phenomena of particular interest would include the evolution of individuality itself [2], the exhibition of a molecular-agent computational model of the chemoton [11], and the evolution of von Neumann style genetic architecture based on a programmable constructor [29].

Francisco Varela was a pivotal figure in the development of our field to this point. His influence and intellectual legacy provide a strong foundation to tackle these difficult and profound problems that we still face in properly understanding the organization of the living, its origin, and its evolution.

## Acknowledgments

Financial support for the work has been provided by the Research Institute for Networks and Communications Engineering (RINCE) at Dublin City University.

## References

1. Breyer, J., Ackermann, J., & McCaskill, J. (1998). Evolving reaction-diffusion ecosystems with self-assembling structures in thin films. *Artificial Life*, 4, 25–40.
2. Buss, L. (1987). *The evolution of individuality*. Princeton, NJ: Princeton University Press.
3. Csendes, T. (1984). A simulation study on the chemoton. *Kybernetes*, 13(2), 79–85.
4. Fleischaker, G. R. (1988)a. *Autopoiesis: System logic and the origin of life*. Ph.D. dissertation, Boston University, Boston, MA.
5. Fleischaker, G. R. (1988)b. Autopoiesis: The status of its system logic. *BioSystems*, 22, 37–49.
6. Fleischaker, G. R. (1990). Origins of life: An operational definition. *Origins of Life and Evolution of the Biosphere*, 20, 127–137.
7. Fleischaker, G. R. (Ed.). (1992). *Autopoiesis—A debate: Controversy over physical, biological, and social systems*. *International Journal of General Systems*, Volume 21.
8. Fontana, W., & Buss, L. W. (1994a). “The arrival of the fittest”: Toward a theory of biological organization. *Bulletin of Mathematical Biology*, 56(1), 1–64.
9. Fontana, W., & Buss, L. W. (1994b). What would be conserved if “the tape were played twice”? *Proceedings of the National Academy of Sciences of the USA*, 91, 757–761.
10. Gánti, T. (1975). Organization of chemical reactions into dividing and metabolizing units: The chemotons. *BioSystems*, 7, 15–21.
11. Gánti, T. (1997). Biogenesis itself. *Journal of Theoretical Biology*, 187, 583–593.
12. Gardner, H. (1971). On cellular automata, self-reproduction, the Garden of Eden, and the game “life.” *Scientific American*, 224, 112–117.
13. Maturana, H. R., Lettvin, J. Y., McCulloch, W. S., & Pitts, W. H. (1960). Anatomy and physiology of vision in the frog. *Journal of General Physiology*, 43, 129–175.
14. Holland, J. H. (1976). Studies of the spontaneous emergence of self-replicating systems using cellular automata and formal grammars. In A. Lindenmayer & G. Rozenberg (Eds.), *Automata, languages, development* (pp. 385–404). New York: North-Holland.
15. Jullien, F., & McMullin, B. (1995). FRJ’s simple autopoiesis program. Program source in Pascal, for MS-DOS platform. [http://www.eeng.dcu.ie/~alife/frj\\_sap/](http://www.eeng.dcu.ie/~alife/frj_sap/).
16. Kauffman, S. A. (1993). *The origins of order: Self-organization and selection in evolution*. Oxford, UK: Oxford University Press.
17. Langton, C. G. (1989). Artificial life. In C. G. Langton (Ed.), *Artificial Life* (pp. 1–47), Vol. VI of *Sante Fe Institute Studies in the Sciences of Complexity*. Redwood City, CA: Addison-Wesley.
18. Letelier, J. C., Marín, G., & Mpodozis, J. (2003). Autopoietic and (M, R) systems. *Journal of Theoretical Biology*, 222, 261–272.
19. Luisi, P. L. (1993). Defining the transition to life: Self-replicating bounded structures and chemical autopoiesis. In W. Stein & F. J. Varela (Eds.), *Thinking about biology* (pp. 3–23). Vol. III of *SFI Studies in the Sciences of Complexity, Lecture Notes*. Redwood City, CA: Addison-Wesley.
20. Luisi, P. L., & Varela, F. (1989). Self replicating micelles: A minimal version of a chemical autopoietic system. *Origins of Life and Evolution of the Biosphere*, 19, 633–643.
21. Madina, D., Ono, N., & Ikegami, T. (2003). Cellular evolution in a 3D lattice artificial chemistry. In W. Banzhaf, T. Christaller, P. Dittrich, J. T. Kim, & J. Ziegler (Eds.), *Advances in artificial life* (pp. 59–68). 7th European Conference on Artificial Life, ECAL 2003, Dortmund, Germany, September 2003. Springer-Verlag.
22. Maturana, H. (1970). Neurophysiology of cognition. In P. Garvin (Ed.), *Cognition: A multiple view* (pp. 3–23). Washington, DC: Spartan Books.

23. Maturana, H. R., & Varela, F. J. (1980). *Autopoiesis and cognition: The Realization of the living*, Vol. 42 of *Boston Studies in the Philosophy of Science*. Dordrecht, The Netherlands: D. Reidel.
24. Maturana, H. R., & Varela, F. J. (1987). *The tree of knowledge: The biological roots of human understanding*. Boston: New Science Library (Shambhala).
25. McMullin, B. (1994). Artificial Darwinism: The very idea! In B. McMullin & N. Murphy (Eds.), *Autopoiesis & Perception* (pp. 71–94). Dublin: Dublin City University. Technical Report bmcm9401. Proceedings of the International Workshop held in DCU, August 1992. <http://www.eeng.dcu.ie/~alife/bmcm9401/mcmullin.pdf>.
26. McMullin, B. (1997)a. Code McMullin: The case of the independent test. *Santa Fe Institute Bulletin*, 12(2). <http://www.santafe.edu/sfi/publications/Bulletins/bulletin-summer97/code.html>.
27. McMullin, B. (1997)b. Computational autopoiesis: The original algorithm. Working Paper 97-01-001, Santa Fe Institute, Santa Fe, NM 87501, USA. <http://www.santafe.edu/sfi/publications/wpabstract/199701001>.
28. McMullin, B. (1997)c. SCL: An artificial chemistry in *Swarm*. Working Paper 97-01-002, Santa Fe Institute, Santa Fe, NM 87501, USA. <http://www.santafe.edu/sfi/publications/wpabstract/199701002>.
29. McMullin, B. (2000)a. John von Neumann and the evolutionary growth of complexity: Looking backwards, looking forwards. *Artificial Life*, 6(4), 347–361.
30. McMullin, B. (2000)b. Some remarks on autocatalysis and autopoiesis. *Annals of the New York Academy of Sciences*, 901, 163–174.
31. McMullin, B., & Groß (Chu), D. (2001). Towards the implementation of evolving autopoietic artificial agents. In J. Kelemen & P. Sosík (Eds.), *Advances in artificial life, Proceedings of the 6th European Conference on Artificial Life* (pp. 440–443). ECAL 2001, Prague, Czech Republic, September 2001. Springer-Verlag. <http://www.eeng.dcu.ie/~alife/bmcm-ecal-2001/>.
32. McMullin, B., & Murphy, N. (1994). Autopoiesis & perception. Technical Report bmcm9401, School of Electronic Engineering, Dublin City University, Dublin. Proceedings of the International Workshop held in DCU, August 1992. <http://www.eeng.dcu.ie/~alife/bmcm9401/>.
33. McMullin, B., & Varela, F. J. (1997). Rediscovering computational autopoiesis. In P. Husbands & I. Harvey (Eds.), *Proceedings of the Fourth European Conference on Artificial Life* (pp. 38–47). Cambridge, MA: MIT Press. <http://www.eeng.dcu.ie/~alife/bmcm-ecal97/>.
34. Mingers, J., & McMullin, B. (1997). JM's simple autopoiesis program. Program source in Pascal, for MS-DOS platform. [http://www.eeng.dcu.ie/~alife/jm\\_sap/](http://www.eeng.dcu.ie/~alife/jm_sap/).
35. Nomura, T. (2001). Formal description of autopoiesis based on the theory of category. In J. Kelemen & P. Sosík (Eds.), *Advances in artificial life, proceedings of the 6th European Conference on Artificial Life* (pp. 700–703). ECAL 2001, Prague, Czech Republic, September 2001. Springer-Verlag.
36. Ono, N., & Ikegami, T. (1999). Model of self-replicating cell capable of self-maintenance. In D. Floreano, J. D. Nicaud, & F. Mondada (Eds.), *Proceedings of the 5th European Conference on Artificial Life (ECAL'99)* (pp. 399–406). Lausanne, Switzerland: Springer-Verlag.
37. Ono, N., & Ikegami, T. (2000). Self-maintenance and self-reproduction in an abstract cell model. *Journal of Theoretical Biology*, 206, 243–253.
38. Ono, N., & Ikegami, T. (2001). Artificial chemistry: Computational studies on the emergence of self-reproducing units. In J. Kelemen & P. Sosík (Eds.), *Advances in artificial life, proceedings of the 6th European Conference on Artificial Life* (pp. 186–195). ECAL 2001, Prague, Czech Republic, September 2001. Springer-Verlag.
39. Ono, N., & Ikegami, T. (2002). Selection of catalysts through cellular reproduction. In R. Standish, M. A. Bedau, & H. A. Abbass (Eds.), *Proceedings of the 8th International Conference on Artificial Life (ALIFE 8)* (pp. 57–64). Sydney, Australia: MIT Press. <http://parallel.hpc.unsw.edu.au/complex/alife8/proceedings/sub2844.pdf>.

40. Rasmussen, S., Baas, N., Mayer, B., Nilsson, M., & Oleson, M. (2001). Ansatz for dynamical hierarchies. *Artificial Life*, 7(4), 367–373.
41. Rasmussen, S., Knudsen, C., Feldberg, R., & Hindsholm, M. (1990). The Coreworld: Emergence and evolution of cooperative structures in a computational chemistry. *Physica D*, 42, 111–134.
42. Ray, T. S. (1992). An approach to the synthesis of life. In C. G. Langton, C. Taylor, J. D. Farmer, & S. Rasmussen (Eds.), *Artificial life II* (pp. 371–408). Redwood City, CA: Addison-Wesley. Proceedings of the workshop on Artificial Life held February, 1990, in Sante Fe, New Mexico.
43. Rosen, R. (1991). *Life itself*. New York: Columbia University Press.
44. Varela, F. J. (1979). *Principles of biological autonomy*. New York: North-Holland.
45. Varela, F. J. (1996). The early days of autopoiesis: Heinz and Chile. *Systems Research*, 13(3), 407–416.
46. Varela, F. J., Maturana, H. R., & Uribe, R. (1974). Autopoiesis: The organization of living systems, its characterization and a model. *BioSystems*, 5, 187–196.
47. von Neumann, J. (1966). *Theory of self-reproducing automata*. Urbana, IL: University of Illinois Press. Edited and completed by Arthur W. Burks.
48. Zeleny, M. (1977). Self-organization of living systems: A formal model of autopoiesis. *International Journal of General Systems*, 4, 13–28.
49. Zeleny, M. (1978). APL-AUTOPOIESIS: Experiments in self-organization of complexity. In *Progress in cybernetics and systems research, Vol. III* (pp. 65–84). Washington, DC: Hemisphere.
50. Zeleny, M. (1980a). Autopoiesis: A paradigm lost. In M. Zeleny (Ed.), *Autopoiesis, dissipative structures and spontaneous social orders* (pp. 3–43). Boulder, CO: Westview Press. AAAS Selected Symposium 55.
51. Zeleny, M. (Ed.) (1980b). *Autopoiesis, dissipative structures and spontaneous social orders*. Boulder, CO: Westview Press. AAAS Selected Symposium 55.
52. Zeleny, M. (Ed.) (1981). *Autopoiesis: A theory of living organization*, Vol. 3 of North Holland Series in General Systems Research. New York: North Holland.
53. Zeleny, M., & Pierre, N. A. (1975). Simulation models of autopoietic systems. In *Summer Computer Simulation Conference, July 21–23 1975, San Francisco* (pp. 831–842).
54. Zeleny, M., & Pierre, N. A. (1976). Simulation of self-renewing systems. In E. Jantsch & C. H. Waddington (Eds.), *Evolution and consciousness: Human systems in transition* (pp. 150–165). Reading, MA: Addison-Wesley.