

# Nonhuman Primates Express Human Leukemia-Associated Antigens

By John M. Pesando and Ted A. Conrad

Serologic studies using four murine monoclonal antibodies specific for the common acute lymphoblastic leukemia antigen (CALLA) and five monoclonal antibodies specific for the gp24 surface antigen indicate that these leukemia-associated antigens are present on cells of comparable tissues in man and in four nonhuman primates. As in man, adherent cell populations obtained from skin, lung, and bone marrow of *Macaca fascicularis*, *M mulatta*, *M nemestrina*, and *Papio cynocephalus* react with these antibodies. Similarly, granulocytes from both man and these nonhu-

man primates bind CALLA- and gp24-specific antibodies. Radioimmune precipitation experiments confirm the identity of these antigens. Our studies suggest that nonhuman primates can be used to screen serologic reagents to leukemia-associated antigens for potential toxic effects on normal tissues prior to their use in man. Similarly, nonhuman primates could be employed to assess the possible role of antigen-positive stromal cells in the reconstitution of bone marrow following transplantation.

**M**ONOCLONAL ANTIBODIES to the common acute lymphoblastic leukemia antigen (CALLA) have been used alone<sup>1</sup> or with monoclonal antibodies to a 24-kD surface glycoprotein (gp24)<sup>2</sup> in efforts to rid marrow of malignant cells prior to autologous bone marrow transplantation for acute lymphoblastic leukemia (ALL). In addition, monoclonal antibodies to CALLA have also been used for passive serotherapy in this disease.<sup>3</sup> Since serologic and immune precipitation studies have demonstrated the presence of a CALLA-like molecule on adherent normal cell populations from human lung, skin, and bone marrow, on a small population of nonadherent cells in normal bone marrow, and on polymorphonuclear leukocytes,<sup>4,7</sup> the possible toxicity of CALLA-specific antibodies and conjugates on normal tissues must be evaluated. Similarly, both serologic and immune precipitation studies have demonstrated the presence of gp24 on normal cell populations from human lung and bone marrow<sup>8,9</sup> (J.M. Pesando, submitted for publication). Serologic studies have also identified gp24 on polymorphonuclear leukocytes.

The purpose of our present studies was to determine whether or not the CALLA and gp24 antigens are present on cells of corresponding normal tissues from nonhuman primates. Adherent cell populations from cultured specimens of lung, skin, and bone marrow from four nonhuman primates (*Macaca fascicularis*,

*M mulatta*, *M nemestrina*, and *Papio cynocephalus*) were tested for the expression of these antigens using a panel of four monoclonal antibodies specific for CALLA and five monoclonal antibodies specific for gp24. Polymorphonuclear leukocytes from these four species were also studied for expression of these two antigens both because of their ready availability and because they could be assayed immediately upon collection, without the need for prior growth in tissue culture.

## MATERIALS AND METHODS

### Antibodies

Four CALLA-specific monoclonal antibodies— $\alpha$ CALLA-1 (J5),  $\alpha$ CALLA-2 (J13),  $\alpha$ CALLA-3 (24.1), and  $\alpha$ CALLA-4—were employed in these studies. The first three of these antibodies have been described previously.<sup>5,6,10</sup>  $\alpha$ CALLA-4 was raised by immunizing BALB/c mice with leukemic cells from the CALLA-positive ALL cell line Laz 221. It identifies a 100,000-dalton surface glycoprotein on CALLA-positive leukemic cells and shows an identical reactivity profile with the other three antibodies against cells from a panel of 32 acute leukemias and 22 hematopoietic cell lines. Modulation of the expression of CALLA by J5 antibody ( $\alpha$ CALLA-1) on reactive cell populations inhibits binding of  $\alpha$ CALLA-4. The CALLA-specific 24.1 antibody ( $\alpha$ CALLA-3) was the gift of Dr Marcus Braun, Fred Hutchinson Cancer Research Center.

Five monoclonal antibodies specific for the gp24 antigen were employed— $\alpha$ gp24-1 (J2),  $\alpha$ gp 24-2 (J9),  $\alpha$ gp24-3,  $\alpha$ gp24-4, and  $\alpha$ gp24-5. One of these antibodies ( $\alpha$ gp24-1) has been described elsewhere,<sup>11</sup> while a detailed description of the remaining antibodies is the subject of another report (J.M. Pesando, submitted for publication). All of these antibodies identify a 24,000-dalton surface protein by radioimmune precipitation, and sequential immune precipitation studies confirm that they identify the same monomorphic antigen. They demonstrate identical patterns of reactivity against all human cell populations tested. All antibodies were obtained as ascites from tumor-bearing mice at a concentration of approximately 7 mg/mL and were ultracentrifuged and filtered (0.45  $\mu$ m Milllex, Millipore, Bedford, Mass) before use.

### Cell Populations

Sterile samples of primate lung, skin, and bone marrow obtained from animals sacrificed for experimental purposes at the Regional Primate Research Center at the University of Washington were cut

From the Divisions of Medical and Pediatric Oncology, Fred Hutchinson Cancer Research Center, Seattle.

Supported by a grant from the Cancer Research Institute, Inc, New York, and by National Institutes of Health grant No. CA15704, CA30924, and CA34206. J.M.P. is the recipient of a Junior Faculty Clinical Fellowship from the American Cancer Society.

Submitted March 26, 1984; accepted June 2, 1984.

Address reprint requests to Dr John M. Pesando, Fred Hutchinson Cancer Research Center, 1124 Columbia St, Seattle, WA 98104.

© 1984 by Grune & Stratton, Inc.  
0006-4971/84/6405-0021\$03.00/0

into small pieces and placed in RPMI medium with 20% fetal bovine serum (FBS) in a humidified atmosphere containing 7% CO<sub>2</sub> at 37 °C. Adherent monolayers were harvested by trypsinization for fluorescence-activated cell sorter (FACS) analysis after four weeks. Cell viability was greater than 95%. Most of these cells appeared either flat and angulated or spindle-shaped. A cryopreserved ampule of *M mulatta* cells was obtained through the American Type Culture Collection.

Granulocytes from peripheral blood samples from the aforementioned primates were isolated by Ficoll-Diatrizoate (LSM, Bionetics, Kensington, Md) density gradient centrifugation using standard procedures. Red cells were selectively removed from granulocyte-enriched pellets by hypotonic lysis in hemolytic buffer (0.155 mol/L ammonium chloride, 0.012 mol/L sodium bicarbonate, and 0.110 mol/L EDTA at pH 7.4). Wright-Giemsa staining confirmed that greater than 90% of the cells so obtained had the morphological appearance of granulocytes. Cell viability was greater than 95%, as measured by trypan blue exclusion.

Laz 221<sup>12</sup> is a CALLA- and gp24-positive human ALL cell line grown in RPMI with 10% FBS. These cells were used as positive controls in all immunofluorescence experiments.

### Immunofluorescence Analysis

Binding of monoclonal antibodies to target cell populations was determined using a standard indirect immunofluorescence assay<sup>13</sup> and monitored on the fluorescence-activated cell sorter (FACS-IV, Becton-Dickinson, Mountain View, Calif), with fluorescence displayed in the logarithmic mode. Prior to immunofluorescence analysis, both adherent cells and granulocytes were suspended in phosphate-buffered saline (PBS) containing 5% human AB serum to block Fc receptor sites. Monoclonal antibodies were used at a 1:1,000 dilution of ascites fluid (0.5 µg) to ensure antibody excess. Fluoresceinated goat antibodies to mouse IgG were purchased from

Meloy Laboratories (Springfield, Va). All procedures were performed at 4 °C to inhibit antibody-induced modulation of CALLA.

### Identification of Antigens

Cell surface radiolabeling with <sup>125</sup>I and immune precipitation of detergent-solubilized cell extracts were performed using standard methods, as described previously.<sup>14</sup> Antibodies bound to protein A coupled to Sepharose were used to extract specific antigen from radiolabeled cell extracts, followed by extensive washing with a detergent-containing buffer. Antigens were resolved by electrophoresis on a 7% to 15% gradient sodium dodecyl sulfate-polyacrylamide slab gel (SDS-PAGE) and visualized by autoradiography. A ratio of 1 mCi of <sup>125</sup>I to 50 × 10<sup>6</sup> hematopoietic cells and 20 × 10<sup>6</sup> adherent cells was used for radiolabeling.

### RESULTS

All CALLA-specific monoclonal antibodies tested reacted with adherent cell populations from all four species of primates. This reactivity is recorded in Table 1 and Fig 1. It is of note that not all target cell populations within each species were reactive. Adherent cells from lung tissue of both *P cynocephalus* and *M mulatta* were reactive, while those from *M nemestrina* and *M fascicularis* were not. Adherent cells of bone marrow from *M mulatta* but not *P cynocephalus* or *M nemestrina* were reactive, while those from *M fascicularis* were not available for testing. In contrast, adherent cells from skin of fetal and adult *M nemestrina*, *M fascicularis*, and *P cynocephalus* were all reactive, while those from *M mulatta* were not tested.


Table 1. Presence of Human Leukemia-Associated Antigens on Nonhematopoietic Cells of Primates\*

Source of Cells	Anti-CALLA Antibodies				Anti-gp24 Antibodies				
	CALLA-1 (J5)	CALLA-2 (J13)	CALLA-3 (24.1)	CALLA-4	gp24-1 (J2)	gp24-2 (J9)	gp24-3	gp24-4	gp24-5
<i>M nemestrina</i> (adult)									
Lung	—	—	—	—	+	+	+	+	+
Skin	+	+	+	+	+	+	+	+	+
Bone marrow	NT†	NT	NT	NT	NT	NT	NT	NT	NT
<i>M nemestrina</i> (151-day fetus)									
Lung	—	—	—	—	+	+	+	+	+
Skin	+	+	+	+	+	+	+	+	+
Bone marrow	—	—	—	—	+	+	+	+	+
<i>M fascicularis</i>									
Lung	—	—	—	—	+	+	+	+	+
Skin	+	+	+	+	+	+	+	+	+
Bone marrow	NT	NT	NT	NT	NT	NT	NT	NT	NT
<i>M mulatta</i>									
Lung	+	+	+	+	+	+	+	+	+
Skin	NT	NT	NT	NT	NT	NT	NT	NT	NT
Bone marrow	+	+	+	+	+	+	+	+	+
<i>P cynocephalus</i>									
Lung	+	+	+	+	+	+	+	+	W‡
Skin	+	+	+	+	+	+	+	+	+
Bone marrow	—	—	—	—	+	+	+	+	W

\*Binding of antibodies was detected using a standard indirect immunofluorescence assay, as described in the text. Background fluorescence was determined by incubating cells with a nonreactive antibody. Samples were scored as seropositive if the majority of cells stained more intensely with test than with control antibodies. Most samples were obtained from a single animal of a given species.

†NT, not tested.

‡W, weak reactivity.



**Fig 1.** Expression of human leukemia-associated CALLA and gp24 antigens on adherent cells from skin and on granulocytes of *M fascicularis*. Cell number is plotted against fluorescence intensity, and 4,000 cells were counted in each sample. (A and B) Binding of  $\alpha$ CALLA and  $\alpha$ gp24 antibodies to adherent cells of skin, respectively; (C and D) binding of these same antibodies to circulating blood granulocytes. Binding of a nonreactive control antibody is indicated by the thin dashed curve in each panel. In A and C, the curves for binding of antibodies  $\alpha$ CALLA-1, -2, and -4 are superimposable and are indicated by a single, thick solid line. That for  $\alpha$ CALLA-3 is indicated by a thin solid line. In B and D, binding of antibodies  $\alpha$ gp24-1, -2, and -3 is indicated by thick solid, thin solid, and thick dashed curves, respectively. Binding of antibodies  $\alpha$ gp24-4 and -5 is not shown.

Although cells having different morphologies were frequently observed in these adherent cell populations, the majority of cells in a specific culture had similar morphologies and demonstrated comparable reactivities with the antibodies being tested (Fig 1). Multiple cell populations were not readily demonstrable by FACS analysis. Our failure to demonstrate serologic reactivity on cells of comparable tissues from all four of these primates most likely reflects inherent differences in the phenotype of the predominant cell type growing in tissue culture rather than the absence of the antigen on all cells of that tissue for a specific primate. Since most samples were obtained from single animals, we cannot exclude the possibility of individual variation within species. We were unable to correlate the expression of CALLA with a specific cellular morphology in these studies. Similar heterogeneity in the expression of CALLA by cultures of bone marrow-derived adherent cells obtained from different normal human donors has also been observed (J.M. Pesando, unpublished observations, September 1983).

In contrast to the results with CALLA, adherent cells from all tissues tested from these four primates expressed the gp24 antigen as judged by their reactivity with a panel of five monoclonal antibodies specific for gp24 (Table 1 and Fig 1). The reduced binding of antibody  $\alpha$ gp24-5 to adherent cells from lung and bone marrow of *P cynocephalus* relative to that of the other four antibodies suggests that it may recognize a unique antigenic determinant. Simultaneous binding of  $\alpha$ gp24-5 and these other gp24-specific antibodies to antigen-positive human cell populations could not be demonstrated, suggesting that the ability of antibodies to bind to cells of different species is a more sensitive test for the identity of the epitopes that they recognize. Adverse steric interactions between bound immunoglobulin molecules are avoided in the latter system.

All CALLA-specific monoclonal antibodies tested reacted with polymorphonuclear leukocytes from man and all four of these primates (Table 2 and Fig 1). Once again, all cells in these samples reacted comparably with the different anti-CALLA antibodies. Polymorphonuclear leukocytes from cebus monkey (but not lemur) also reacted with these antibodies suggesting that CALLA first appeared in a common ancestor of Old World and New World monkeys. Similarly, monoclonal antibodies specific for the gp24 antigen bound to samples of polymorphonuclear leukocytes from all primates tested, though only weak binding to cells from some human donors was noted.  $\alpha$ Gp24-5 failed to show significant binding to granulocytes of *P cynocephalus*. Binding of  $\alpha$ gp24-5 to granulocytes from cebus monkey and lemur was also not detected, despite binding of the other four gp24-specific antibodies to these samples.

Radioimmune precipitation experiments were performed using  $\alpha$ CALLA and  $\alpha$ gp24 antibodies to exclude the possibility that identical antigenic determinants might be shared by otherwise dissimilar macromolecules (Figs 2 and 3). These studies confirm that similar molecular species are identified by these antibodies on the ALL cell line Laz 221, human


**Table 2.** Presence of Human Leukemia-Associated Antigens on Granulocytes From Primates\*

Source of Cells	Anti-CALLA Antibodies				Anti-gp24 Antibodies				
	CALLA-1 (J5)	CALLA-2 (J13)	CALLA-3 (24.1)	CALLA-4	gp24-1 (J2)	gp24-2 (J9)	gp24-3	gp24-4	gp24-5
<i>M nemestrina</i>	+	+	+	+	+	+	+	+	+
<i>M fascicularis</i>	+	+	+	+	+	+	+	+	+
<i>M mulatta</i>	+	+	+	+	NT†	NT	NT	NT	NT
<i>P cynocephalus</i>	+	+	+	+	+	+	W‡	+	-
Cebus monkey	+	+	+	+	+	+	+	+	-
Lemur	-	-	-	-	+	+	+	+	-

\*Binding of antibodies was detected as described in the text and in Table 1. For *M nemestrina*, *M fascicularis*, *M mulatta*, and *P cynocephalus*, granulocytes from at least two individuals were tested for expression of CALLA.

†NT, not tested.

‡W, weak binding.




**Fig 2.** Presence of CALLA and gp24 antigens on primate cells. Results obtained with the ALL cell line Laz 221 are shown in lanes A and F, with human granulocytes in lanes B and G, with granulocytes of *M fascicularis* in lanes C and H, and with adherent cells from skin of *M fascicularis* in lanes D and I. Cells were surface-labeled with  $^{125}\text{I}$  before immune precipitation of cell extracts with  $\alpha\text{CALLA-1}$  (lanes A through D) or  $\alpha\text{gp24-2}$  (lanes F through I). Molecular weight markers labeled with  $^{14}\text{C}$ -acetate from New England Nuclear (Boston) include myosin (200,000 daltons), phosphorylase B (97,000 daltons), albumin (69,000 daltons), ovalbumin (46,000 daltons), carbonic anhydrase (30,000 daltons), and lactoglobulin (18,000 daltons).

granulocytes, granulocytes from *M fascicularis*, and adherent cells from the skin of *M fascicularis*. Molecules identified by  $\alpha\text{CALLA-1}$  (J5 antibody) on Laz 221 cells and on granulocytes and adherent skin cells of *M fascicularis* have molecular weights of approximately 100,000 daltons. In contrast, that identified by  $\alpha\text{CALLA-1}$  on human granulocytes has a molecular weight of approximately 110,000 daltons, in agreement with the findings of Braun et al.<sup>6</sup> The basis for the difference in the molecular weight of the antigen detected by  $\alpha\text{CALLA-1}$  on granulocytes as compared to other human hematopoietic cells is currently under investigation. The  $\alpha\text{gp24-2}$  antibody identifies antigens having a molecular weight of approximately 24,000 daltons on all four cell types.

#### DISCUSSION

For several years, monoclonal antibodies have been used clinically in efforts to eradicate malignant cells and to manipulate the immune response for therapeutic purposes.<sup>1-3,15-18</sup> Monoclonal antibodies represent a diverse group of new pharmacologic reagents, but they have not been tested for toxicity in animals because of their apparent species specificity. Our serologic and immune precipitation studies indicate that the leukemia-associated CALLA and gp24 antigens are expressed by comparable normal cells of man and nonhuman primates. These observations suggest that nonhuman primates can be used to screen both conju-



**Fig 3.** Presence of CALLA on primate cells. Cells were surface labeled with  $^{125}\text{I}$ , and antigens detected by  $\alpha\text{CALLA-1}$  were isolated and visualized as described in the text. Samples are as follows: (A) ALL cell line Laz 221; (B) human granulocytes; (C) granulocytes of *M fascicularis*; and (D) adherent cells from skin of *M fascicularis*. Bands having molecular weights of approximately 100,000 daltons are seen in lanes A, C, and D, while once again, the antigen isolated from human granulocytes has a slightly higher molecular weight.

gated and unconjugated serologic reagents to leukemia-associated antigens for potential adverse side effects on normal tissues prior to their use in man. Similarly, the recent demonstration of cross-reactivity of monoclonal antibodies to human T cell and myeloid antigens with appropriate primate hematopoietic cells<sup>19-21</sup> suggests that primates may also serve as important models for evaluating the toxicities of serologic reagents to other human tumor-associated anti-

Gp24-specific monoclonal antibodies react with adherent cells obtained from every sample of primate and human bone marrow tested. CALLA-specific antibodies react with several but not all of these samples. These same adherent cells presumably include the hematopoietic stromal cells that are required for long-term culture of marrow cells in vitro.<sup>22,23</sup> Therefore, successful efforts to purge bone marrow of leukemic cells using gp24-specific monoclonal antibodies prior to autologous transplantation would most likely eliminate these adherent cells as well as leukemic ones. Whether or not the removal of such stromal cells from treated marrow would affect subsequent engraftment and hematopoiesis would depend on the efficacy of the cytolytic process employed and on whether or not the chemoradiotherapy preparative regimens also eliminated stromal cells in the patient. At least one report in the literature suggests that

stromal cells become progressively donor in origin with time after allogeneic bone marrow transplantation.<sup>24</sup> Our present studies suggest that nonhuman primates could be employed to assess the possible role of antigen-positive stromal cells in bone marrow reconstitution and engraftment.

Finally, it should be noted that despite the presence of a CALLA-like molecule on human granulocytes,

CALLA has not been demonstrated on normal or malignant myeloid blasts by ourselves or others, despite study of over 700 patients with acute nonlymphocytic leukemia (ANL)<sup>5,13,25</sup> (J.M. Pesando, unpublished observations, March 1984). Therefore, antibodies to this antigen retain their considerable operational utility as reagents with which to discriminate between ALL and ANL.

#### REFERENCES

- Ritz J, Sallan SE, Bast RC, Lipton JM, Clavell LA, Feeney M, Hercend J, Nathan DG, Schlossman SF: Autologous bone marrow transplantation in CALLA-positive acute lymphoblastic leukemia following in vitro treatment with J5 monoclonal antibody and complement. *Lancet* 2:60, 1982
- Ramsay N, LeBien T, Nesbit M, McGlave P, Weisdorf D, Hurd D, Kenyon P, Goldman A, Kim T, Kersey J: Autologous bone marrow transplantation (BMT) for acute lymphoblastic leukemia (ALL) following marrow treatment with BA-1, BA-2, BA-3 and rabbit complement (C'). *Blood* 62:228a, 1983 (abstr)
- Ritz J, Pesando JM, Sallan SE, Clavell LA, Notis-McConarty J, Rosenthal P, Schlossman SF: Serotherapy of acute lymphoblastic leukemia with monoclonal antibody. *Blood* 58:141, 1981
- Pesando JM, Ritz J, Lazarus H, Tomaselli KJ, Schlossman SF: Fate of a common acute lymphoblastic leukemia antigen during modulation by monoclonal antibody. *J Immunol* 126:540, 1981
- Pesando JM, Tomaselli KJ, Lazarus H, Schlossman SF: Distribution and modulation of a human leukemia-associated antigen (CALLA). *J Immunol* 131:2038, 1983
- Braun MP, Martin PJ, Ledbetter JA, Hansen JA: Granulocytes and cultured human fibroblasts express common acute lymphoblastic leukemia associated antigens. *Blood* 61:718, 1983
- Cossmann J, Neckers LM, Leonard WJ, Green WC: Polymorphonuclear neutrophils express the common acute lymphoblastic leukemia antigen. *J Exp Med* 157:1064, 1983
- Kersey JH, LeBien TW, Abramson CS, Newman R, Sutherland R, Greaves M: p24: A human leukemia-associated and lymphohematopoietic progenitor cell surface structure identified with monoclonal antibody. *J Exp Med* 153:726, 1981
- Jones NH, Borowitz NJ, Metzgar RS: Characterization and distribution of a 24,000 molecular weight antigen defined by a monoclonal antibody (DU-ALL-1) elicited to common acute lymphoblastic leukemia (cALL) cells. *Leuk Res* 6:449, 1982
- Ritz J, Pesando JM, Notis-McConarty J, Lazarus H, Schlossman SF: A monoclonal antibody to human acute lymphoblastic leukemia antigen. *Nature* 283:583, 1980
- Hercend T, Nadler LM, Pesando JM, Reinherz EL, Schlossman SF, Ritz J: Expression of a 26,000 dalton glycoprotein on activated human T cells. *Cell Immunol* 64:192, 1981
- Lazarus H, Barell EF, Krishan A, Livingston DM, Harris K, Schlossman SF: Characterization of a unique cell line (Laz 221) from a human acute lymphocytic ("null cell") leukemia. *Cancer Res* 38:1362, 1978
- Pesando JM, Ritz J, Lazarus H, Costello SB, Sallan SE, Schlossman SF: Leukemia-associated antigens in ALL. *Blood* 54:1240, 1979
- Pesando JM, Ritz J, Levine H, Terhorst C, Lazarus H, Schlossman SF: Human leukemia-associated antigen: Relation to a family of surface glycoproteins. *J Immunol* 124:2794, 1980
- Miller RA, Maloney DG, Warnke R, Levy R: Treatment of B-cell lymphoma with monoclonal anti-idiotypic antibody. *N Engl J Med* 306:517, 1982
- Dillman RO, Shawler DL, Sobol RE, Collins HA, Beauregard JC, Wormsley SB, Royston I: Murine monoclonal antibody therapy in two patients with chronic lymphocytic leukemia. *Blood* 59:1036, 1982
- Miller RA, Oseroff AR, Stratte PT, Levy R: Monoclonal antibody therapeutic trials in seven patients with T-cell lymphoma. *Blood* 62:988, 1983
- Bozdech MJ, Finlay JL, Trigg ME, Billing R, Hong R, Sugden W, Sondel PM: Monoclonal B-cell lymphoproliferative disorder following monoclonal antibody (CT2) T cell-depleted allogeneic bone marrow transplantation. *Blood* 62:218a, 1983 (abstr)
- Haynes BF, Dowell BL, Hensley LL, Gore I, Metzgar RS: Human T cell antigen expression by primate T cells. *Science* 215:298, 1982
- Letvin NL, King NW, Reinherz EL, Hunt RD, Lane H, Schlossman SF: T lymphocyte surface antigens in primates. *Eur J Immunol* 13:245, 1983
- Letvin NL, Todd RF, Palley LS, Schlossman SF, Griffin JD: Conservation of myeloid surface antigens on primate granulocytes. *Blood* 61:408, 1983
- Dexter TM, Allen TD, Lajtha LG: Conditions controlling the proliferation of hematopoietic stem cells in vitro. *J Cell Physiol* 91:335, 1977
- Dexter TM: Is the marrow stroma transplantable? *Nature* 298:222, 1982
- Keating A, Singer JW, Killen PD, Stricker GE, Salo AC, Sanders J, Thomas ED, Thorning D, Falikow PJ: Donor origin of the in vitro haematopoietic microenvironment after marrow transplantation in man. *Nature* 298:280, 1982
- Greaves MF, Hariri G, Newman RA, Sutherland DR, Ritter MA, Ritz J: Selective expression of the common acute lymphoblastic leukemia (gp100) antigen on immature lymphoid cells and their malignant counterparts. *Blood* 61:628, 1983