

## Nuclear Factor- $\kappa$ B in Development, Prevention, and Therapy of Cancer

Carter Van Waes

**Abstract** Nuclear factor- $\kappa$ B (NF- $\kappa$ B) is a signal transcription factor that has emerged as an important modulator of altered gene programs and malignant phenotype in development of cancer. Major carcinogens and oncogenic viruses induce NF- $\kappa$ B activation, and a variety of subsequent oncogenic events contribute to a progressive increase in constitutive NF- $\kappa$ B activation as an important common pathway in most forms of cancer. NF- $\kappa$ B target genes promote tumor cell proliferation, survival, migration, inflammation, and angiogenesis. Inhibition of NF- $\kappa$ B has been found to be an important mechanism of action of steroids, nonsteroidal anti-inflammatory drugs, and natural and synthetic compounds that show therapeutic and preventive activity. Newer agents targeting the proteasome, inhibitor- $\kappa$ B kinase, and other upstream kinases involved in NF- $\kappa$ B activation have shown anticancer activity in clinical or preclinical studies.

Nuclear factor- $\kappa$ B (NF- $\kappa$ B) was originally identified as a bacterial lipopolysaccharide-induced nuclear transcription factor that regulates immunoglobulin  $\kappa$  light chain expression in B lymphocytes (1). Subsequently, NF- $\kappa$ B family members have been shown to be expressed in all cells, where they regulate expression of multiple genes important in cell survival, host responses to injury and infection, and contribute to pathogenesis of various diseases, including cancer (1–4). Five NF- $\kappa$ B family members have been identified in mammalian cells, including NF- $\kappa$ B1 (p105/p50), NF- $\kappa$ B2 (p100/p52), REL A (p65), cREL, and RELB, which associate to form various heterodimeric and homodimeric combinations. These are initially expressed in inactive form in the cytoplasm, by virtue of self-folding with contiguous ankyrin repeat domains, or association with separate ankyrin-containing inhibitor- $\kappa$ Bs (I $\kappa$ B; Fig. 1). Signal-induced phosphorylation and ubiquitination of the 105 and p100 precursors and I $\kappa$ Bs results in proteasome-mediated processing of NF- $\kappa$ B1 (p50), NF- $\kappa$ B2 (p52), and I $\kappa$ Bs. Ankyrin/I $\kappa$ B degradation results in exposure of the nuclear localization sequence and DNA-binding sites for nuclear translocation and DNA binding of NF- $\kappa$ Bs (Fig. 1).

Signal regulation of I $\kappa$ B degradation and NF- $\kappa$ B transactivation involves a family of I $\kappa$ B kinases (IKK), other kinases, and the ubiquitin E3 ligase SCF/ $\beta$ TrCP, which play important roles in integrating a diversity of upstream signals (Fig. 1; reviewed in refs. 1–4). A current paradigm for signaling by

“classic” (canonical) and “alternative” (noncanonical) IKK/NF- $\kappa$ B pathways has evolved, based largely on studies of the functional effects of knockout of specific IKK and NF- $\kappa$ B/REL subunits on development, immunity, and the cancer phenotype (Fig. 1; reviewed in refs. 1, 2). The classic pathway, activated by tumor necrosis factor (TNF) and many other signals (Fig. 1A and B) through an IKK $\alpha$ /IKK $\beta$ /IKK $\gamma$  complex, has been reported to mediate IKK $\beta$ -induced phosphorylation- and proteasome-dependent degradation of I $\kappa$ Bs and activation of NF- $\kappa$ B1/REL A (p50/p65) and/or p50/cREL. Injury- or pathogen-induced activation of this pathway in normal cells has been shown to be important for cell survival and innate and adaptive immunity. An alternative pathway, activated by other TNF family members via the NF- $\kappa$ B inducing kinase, involves IKK $\alpha$ /IKK $\alpha$  homodimers, which activate p100/RelB for processing into NF- $\kappa$ B2 p52/RelB heterodimers (Fig. 1C). Knockout studies indicate that the alternative pathway components regulate survival of premature B lymphocytes and development of peripheral lymphoid tissues. Additional positive signal modifications of IKK and NF- $\kappa$ B components and interacting cofactors by CK2 or Akt, or interactions with negative regulators, such as the p14ARF/p53 pathway, have been shown to determine whether DNA-bound NF- $\kappa$ B results in gene activation or repression (Fig. 1) and will be reviewed below.

Many pathogens, growth factors, cytokines, and carcinogens that induce activation of the classic pathway and heterodimer NF- $\kappa$ B1/REL A (p50/p65) have been implicated in promotion and pathogenesis of cancer. Induced activation by cytotoxic agents, and constitutive activation of the classic pathway by oncogenic activation of upstream receptor and non-receptor tyrosine kinases, has been shown to promote therapeutic resistance and cancer cell survival (Fig. 1A and B; refs. 2, 3). These findings have resulted in interest in development of proteasome and  $\beta$ TrCP inhibitors for therapy, and more specific IKK $\beta$ -specific inhibitors, to target the classic pathway for prevention and therapy of cancer. However, recent evidence indicating that signal activation of both IKK $\alpha$  and IKK $\beta$ , and NF- $\kappa$ B1 and NF- $\kappa$ B2, may promote survival in different cancers,


**Author's Affiliation:** Head and Neck Surgery Branch, National Institute on Deafness and Other Communication Disorders and Center for Cancer Research, National Cancer Institute, Bethesda, Maryland

Received 9/6/06; revised 9/16/06; accepted 9/19/06.

**Grant support:** National Institute on Deafness and Other Communication Disorders intramural project Z01-DC-00016.

**Requests for reprints:** Carter Van Waes, Head and Neck Surgery Branch, National Institute on Deafness and Other Communication Disorders, CRC Room 4-2732, 10 Center Drive, Bethesda, MD 20892. Phone: 301-402-4216; Fax: 301-402-1140; E-mail: vanwaesc@nidcd.nih.gov.

©2007 American Association for Cancer Research.  
doi:10.1158/1078-0432.CCR-06-2221



**Fig. 1.** NF- $\kappa$ B activation in pathogenesis and therapy of cancer. *A*, NF- $\kappa$ B activation in cancer development is linked with chronic exposure to bacteria, certain viral products, chemical promoters, and carcinogens and reactive oxygen species, which cause repeated DNA damage. Induction also occurs in response to cytotoxic chemotherapy and ionizing radiation. *B*, classic NF- $\kappa$ B pathway activation occurs in response to these bacterial, viral, chemical or physical stimuli; by aberrant cytokine, integrin, and growth factor ligand/receptor activation (e.g., TNFR, IL-1R, integrin  $\alpha_6\beta_4$ , EGFR, HER-2/*neu*, and other serum factors); expression of activating fusion proteins following translocations (e.g., BCR-ABL and MALT1); or aberrant activation by intermediate kinases (e.g., PI3K, CK2, and AKT). Intermediate kinases convey signals to the I $\kappa$ B complex formed by IKK $\alpha$ , IKK $\beta$ , and IKK $\gamma$ , and IKK $\beta$  and CK2 phosphorylate I $\kappa$ B, marking it for ubiquitination by E3 ligase  $\beta$ TrCP (SCF) and proteasome degradation. I $\kappa$ B is processed to NF- $\kappa$ B1 (p50)/RELA or cREL heterodimers, which translocate to the nucleus and bind promoters of genes regulating proliferation, apoptosis, migration inflammation, angiogenesis, and innate immunity. *C*, alternative pathway. The alternative pathway may be activated by other TNF family members via the NF- $\kappa$ B inducing kinase and involves IKK $\alpha$ /IKK $\alpha$  homodimers, which activate NF- $\kappa$ B2/p100 for processing into p52/RelB heterodimers. The RelB/p52 heterodimer then translocates into the nucleus to bind the promoter of genes whose products are important for the malignant phenotype in some cancers and B-cell development and adaptive immunity. *D*, in certain leukemias, overexpression of BCL3 can activate NF- $\kappa$ B2. *E*, repression of gene activation can occur in the presence of intact p14ARF and p53, which favor replacement of CBP/p300 with histone deacetylases. Red highlighted inhibitors of NF- $\kappa$ B activation under clinical and preclinical investigation include proteasome,  $\beta$ TrCP and IKK antagonists, and inhibitors of receptor and intermediate kinases involved in activation. Abbreviations: NIK, NF- $\kappa$ B inducing kinase; IL-1R, interleukin 1 receptor; EGFR, epidermal growth factor receptor; TNFR, TNF receptor; TRAF, TNF receptor – associated factor; TAK, transforming growth factor- $\beta$  – activated kinase; FAK, focal adhesion kinase; CK2, casein kinase 2; PI3K, phosphatidylinositol 3-kinase; PDK, 3-phosphoinositide – dependent protein kinase.

and that NF- $\kappa$ B may inhibit or promote cancer development in different models, breaks down previously accepted paradigms, suggesting a more complex role for the IKKs and NF- $\kappa$ Bs as targets for cancer prevention and therapy.

### Links between NF- $\kappa$ B, Infection, Inflammation, and Carcinogens in Development of Cancer

Cumulative experimental and epidemiologic evidence indicates that pathogen-, carcinogen-, and inflammation-induced

activation of NF- $\kappa$ B plays direct or indirect roles in cellular promotion, transformation, and progression of experimental and human cancers (refs. 2, 4, 5; Fig. 1A and B). The first evidence implicating pathogen-induced NF- $\kappa$ B activation in development of cancer was identification of the REV-T viral oncogene that causes avian reticuloendothelial lymphomatosis, v-Rel (4, 6), which shares a Rel transactivation domain with the mammalian homologues NF- $\kappa$ B1, NF- $\kappa$ B2, RelA (p65), cRel, and RelB. Figure 1 summarizes a number of alterations and signal pathways that have been implicated in activation of NF- $\kappa$ B in human cancers (2–6). EBV gene *LMP-1* and human

T lymphocyte virus gene *Tax* encode proteins that activate NF- $\kappa$ B and commander gene programs that promote cell proliferation, survival, and inflammation, contributing to the pathogenesis of lymphomas, nasopharyngeal carcinomas, and adult T-cell leukemias. Human papillomavirus genes *E6* and *E7* that inactivate *p53* and *Rb* tumor suppressor genes have also been implicated in NF- $\kappa$ B activation and are associated with premalignant and malignant lesions of the cervix, oropharynx, and larynx, where human papillomavirus-associated neoplasms are prevalent. Hepatitis B and C are major causes of hepatocellular carcinoma; hepatitis B X protein and hepatitis C 5A and core proteins have been shown to activate NF- $\kappa$ B and are implicated in hepatocellular transformation. Recently, gastric *Helicobacter pylori* and colonic bacteria in patients with ulcerative colitis have also been implicated in NF- $\kappa$ B activation in epithelia and inflammatory leukocytes and promotion of gastric and colon carcinomas.

Major chemical and physical carcinogens implicated in initiation and/or promotion of human cancer can also activate NF- $\kappa$ B. Notably, nicotine and carcinogens in tobacco and betel nut (areca), which are linked to pathogenesis of head and neck and lung malignancies, induce AKT and NF- $\kappa$ B and promote cell proliferation, survival, and inflammation (4, 5, 7). Nicotine has been reported to directly activate these pathways via nicotinic receptors and Akt (7), whereas chemotherapy- and radiation-induced DNA damage has been reported to induce NF- $\kappa$ B activation via nuclear to cytoplasmic signaling mechanism involving sumoylation of the IKK complex (8). TNF,  $\gamma$  radiation, and certain chemotherapeutic drugs also induce NF- $\kappa$ B activation and several target antiapoptotic genes (*TRAF*, *IAP*, and *BCL-2* and *Bcl-XL*) that protect cells from therapeutic injury by these agents (4).

### Constitutive Activation of NF- $\kappa$ B in Cancer

There is now considerable evidence that sustained or constitutive activation of NF- $\kappa$ B is prevalent in cell lines and tumor tissue specimens and contributes to malignant progression and therapeutic resistance in most of the major forms of human cancer. NF- $\kappa$ B/Rels are constitutively activated in human lymphomas (9), carcinomas of the breast (10), prostate (11), lung (12), colon (13), pancreas (14), head and neck (15), esophagus (16), and cervix (17).

Although the mechanisms of activation are incompletely understood, aberrant activation of upstream tyrosine receptor and non-receptor kinases via IKKs or other kinases has been found to be the most common basis for NF- $\kappa$ B activation in epithelial and lymphoid malignancies (Fig. 1, highlighted in green). Autocrine or paracrine activation of NF- $\kappa$ B resulting from overexpression of epidermal growth factor (18), HER-2/*neu* (19), interleukin-1 (20), hepatocyte growth factor (21), and integrin family (22) ligands and receptors has been reported. Epidermal growth factor receptor and HER-2/*neu* signaling involving phosphatidylinositol 3-kinase (PI3K), IKK, and CK2 has been shown in breast cancer (23). Interleukin-1/interleukin-1R, transforming growth factor- $\alpha$ /epidermal growth factor receptor, PI3K, Akt, CK2, and IKK have been shown to mediate activation in head and neck squamous cell carcinomas (20, 24, 25). Hepatocyte growth factor/PI3K/Pak/IKK signaling is found in prostate carcinoma (21). Activation via CK2 and IKK is observed in colon carcinomas (26). PI3K/Akt and IKK

seem to be important in signal activation of NF- $\kappa$ B and cell survival in many cancers (27–30). The BCR-ABL fusion oncogene has been implicated in NF- $\kappa$ B activation, cell survival, and tumorigenesis in human leukemias (ref. 31; Fig. 1A). Activation by a translocation that produces a MALT-1 fusion protein has reported in diffuse large B-cell lymphomas (ref. 32; Fig. 1A). Constitutive activation of NF- $\kappa$ B p52:p52 due to overexpression and association with the transactivating family member BCL-3 has been detected in breast carcinomas and lymphomas (refs. 33, 34; Fig. 1D). Direct mutation or altered expression of NF- $\kappa$ B molecules has only rarely been found in human cancers and in Hodgkin's lymphomas, where mutations of I $\kappa$ B that favor activation have been identified (35, 36).

Loss of the inhibitory effects of certain tumor suppressor genes may also be important in function of NF- $\kappa$ B as a prosurvival factor (Fig. 1E, highlighted in *stippled red*). Evidence indicates that activation of functional ARF, ATR, and Chk2 mediates alternative phosphorylation of RelA (p65), and p53 competes for CBP/p300, resulting in repression of NF- $\kappa$ B prosurvival genes when the ARF-ATR-p53 pathway and proapoptotic target genes are activated (37). Inactivation of this proapoptotic pathway by hypermethylation of *p14ARF/p16INK4A* or mutation, or human papillomavirus inactivation of p53, are among the most common alterations in human cancers. Loss of phosphatase and tensin homologue deleted on chromosome 10 (PTEN) expression is associated with increased PI3K/Akt signal activation of NF- $\kappa$ B and c-Myc in prostate carcinoma cells (38).

NF- $\kappa$ B has been shown to regulate many of the genes differentially expressed and implicated in cell proliferation, survival, migration, and tumorigenesis and metastasis in cancer. NF- $\kappa$ B-related gene signatures have been identified and associated with malignant phenotype in squamous cell carcinomas (39–41), Hodgkin's and certain non-Hodgkin's lymphomas (42, 43), and inflammatory breast cancer (44). A number of key genes in cancer have been shown to be targets and transcriptionally regulated by NF- $\kappa$ B. Targets of NF- $\kappa$ B important in cell proliferation and survival include prominent oncogenes such as *cyclin D*, *Bcl-XL*, and *IAP* (45–47). Expression of key angiogenesis factors and adhesion molecules, such as GRO1, interleukin-8 and vascular endothelial growth factor, are directly or indirectly enhanced by NF- $\kappa$ B activation (48). Together, these genes contribute to the increase in proliferation, survival, inflammation, and angiogenesis that leads to rapid tumorigenesis and metastasis.

### Clinical-Translational Advances

Inhibition of NF- $\kappa$ B has been found to be an important mechanism of action of steroids, nonsteroidal anti-inflammatory drugs (NSAID), and natural and synthetic compounds that show therapeutic and preventive activity with acceptable safety profiles. Newer agents targeting the proteasome, IKK, and other upstream kinases involved in NF- $\kappa$ B activation have shown anticancer activity in clinical or preclinical studies.

**NSAIDs and cancer prevention.** NSAIDs, such as aspirin, sulindac, ibuprofen, and celecoxib, have been shown to inhibit NF- $\kappa$ B activation and arachidonic acid inflammatory pathways upstream and downstream of NF- $\kappa$ B (49–52). NSAIDs have shown inhibitory activity against cancer cells *in vitro* and/or *in vivo* models (53). NSAIDs have been shown to reduce

adenoma and colon cancer development in patients with inflammatory bowel disease and hereditary colon cancer, and long-term NSAID use has also been associated with a reduction in risk of colon, breast, and prostate cancer in population-based studies (54–57). Several natural compounds affecting NF- $\kappa$ B (e.g., curcumin, derived from tumeric, the spice in curry) have also shown promise as potential chemopreventive and therapeutic agents (58, 59). Further studies of the efficacy and NF- $\kappa$ B–dependent and NF- $\kappa$ B–independent mechanisms of action of synthetic and dietary NSAIDs are needed. In addition, potential benefits must be compared with possible risks of gastrointestinal or cardiovascular side effects observed with some of the synthetic agents.

**Corticosteroids and cytotoxic agents used for therapy.** The cytotoxic effects of corticosteroids in combination with other DNA-damaging agents led to the use of steroid-based regimens as a current mainstay of treatment of certain leukemias, lymphomas, and myelomas. Subsequently, corticosteroids were shown to mediate many of their anti-inflammatory and tumor cytotoxic effects through inhibition of NF- $\kappa$ B (60), and these lymphoid malignancies and supporting host responses were found to be exquisitely dependent on NF- $\kappa$ B–regulated survival or inflammatory mechanisms (61). Recently, *cis*-platinum, which has cytotoxic activity and is a standard radiation sensitizer in a wide range of epithelial malignancies, has also been shown to inhibit NF- $\kappa$ B activation and expression of antiapoptotic genes, in addition to its effects as a DNA-damaging agent (62). These agents have found a significant place in current cancer therapy, with supportive measures to manage the hematopoietic and immunosuppressive side effects expected with agents with broad effects on NF- $\kappa$ B, DNA replication, and other functions.

**Proteasome inhibitors.** A new class of therapeutic agents under development are proteasome inhibitors, which regulate degradation of I $\kappa$ B and inhibit NF- $\kappa$ B, as well as turnover of other cellular proteins (63). The proteasome inhibitor bortezomib (VELCADE, Millennium Pharmaceuticals, Cambridge, MA) has been shown to have significant preclinical and clinical activity and is now approved for treatment of patients with therapy-resistant multiple myeloma (63). Multiple myeloma cells, and the bone marrow stromal interactions on which they depend, exhibit sensitivity to bortezomib through NF- $\kappa$ B–dependent and NF- $\kappa$ B–independent mechanisms. Bortezomib has also recently been shown to have activity in other B cell–related malignancies, including mantle cell lymphoma and Waldenstrom macroglobulinemia (64, 65). Despite evidence for constitutive activation of NF- $\kappa$ B and prosurvival mechanisms in most solid cancers, limited or no activity has been observed with bortezomib monotherapy in Hodgkin's lymphoma, metastatic melanoma, or colon, breast, prostate, and renal cell carcinomas (66–71).

Evidence from preclinical and clinical studies suggest that combination with other cytotoxic therapies, or inhibitors of other prosurvival pathways, may enhance antitumor activity of bortezomib in some cancers. In preclinical and an ongoing phase I clinical study, we have found that bortezomib has NF- $\kappa$ B inhibitory, antiproliferative, proapoptotic, and radiation-sensitizing activity in head and neck squamous cell carcinoma (72). In combination with re-irradiation, partial clinical response and stabilization of disease was observed in a subset of patients. Dose-limiting hypotension and hyponatremia was observed with 6 weeks of twice weekly bortezomib (0.6 and 0.9

mg/M<sup>2</sup>/dose) but has been reduced with amendment of the schedule to include a mid-course treatment break and i.v. fluid support, and dose escalation is ongoing. A combination, including bortezomib, gemcitabine, and carboplatinum, has shown activity and prolonged survival in patients with non-small cell lung cancer (73). Combinations with platinum or taxanes have also shown clinical responses in patients with advanced malignancies in phase I studies (74–76). The possible potentiation of effects of combining proteasome and NF- $\kappa$ B inhibition together with DNA-damaging agents, or with agents targeting other pathways, is a subject of current interest. The role of proteasome inhibitors in combination with other standard or molecular targeted therapies in solid malignancies, and role of NF- $\kappa$ B, awaits determination in larger phase II studies.

**IKK inhibitors.** IKK $\beta$  antagonists that more specifically inhibit I $\kappa$ B and the NF- $\kappa$ B classic pathway initially implicated in cancers have been the subject of intensive development and preclinical studies (2, 77). This is based largely on evidence that IKK $\beta$  and p65 knockout sensitize cells to apoptosis and cytotoxic therapies, and anticipation that fewer side effects may be encountered with more specific agents. The IKK $\beta$ -selective antagonist PS-1145 or ML120b derivatives (Millennium Pharmaceuticals) have shown antiproliferative, cytotoxic, and antitumor activity in preclinical studies with multiple myeloma cells, diffuse large B-cell lymphoma, chronic myelogenous leukemia, and prostate carcinoma cells (78–80). Another IKK $\beta$  antagonist, BMS-345541 (Bristol-Myers Squibb, Princeton, NJ), has shown apoptotic and antitumor activity in melanoma xenograft models (81). These agents exhibit IC<sub>50</sub>s for their molecular targets in the submicromolar range and for inhibition of sensitive cells in the 5 to 30 micromolar range (78, 81). To date, no clinical trials with IKK inhibitors have been reported.

It is not clear that targeting IKK $\beta$  and the classic pathway according to the current paradigm will provide sufficient blockade of NF- $\kappa$ B signal pathways. Our finding that IKK $\beta$  inhibitors only partially inhibit NF- $\kappa$ B activation and are not cytotoxic in head and neck cancers led us to the discovery that IKK $\alpha$  and IKK $\beta$  as well as protein kinase CK2 (formerly casein kinase 2) contribute to activation of NF- $\kappa$ B (25). Several other studies indicate IKK $\alpha$  may contribute to activation of both NF- $\kappa$ B1 and NF- $\kappa$ B2 and survival of cancer cells (82–84), suggesting that alternative and classic NF- $\kappa$ B pathway activation may promote cell survival in other cancers. These findings suggest there is a need to re-examine the paradigm upon which IKK $\beta$  drug development has been based, and to identify inhibitors of both IKK $\alpha$  and IKK $\beta$ , CK2, other kinases, or ubiquitin ligases that specifically regulate NF- $\kappa$ B transactivation (Fig. 1).

### NF- $\kappa$ B as Proapoptotic or Antiapoptotic Signal in Cancer?

Differing results between human cancers reviewed previously and among several experimental animal models indicate that the mechanism or stage of carcinogenesis, affected cell subpopulation, or other molecular changes may determine whether NF- $\kappa$ B acts as a repressor or promoter of malignancy. Some of these findings have raised questions about whether targeting NF- $\kappa$ B for prevention could have deleterious rather than beneficial effects. These experimental animal studies

examined effects of tissue targeted alterations in expression of I $\kappa$ B, IKK $\beta$ , or NF- $\kappa$ B on cancer development. Consistent with a role of IKK/NF- $\kappa$ B in promotion and pathogenesis of cancer, inhibition of NF- $\kappa$ B by targeted IKK $\beta$  knockout in colonic epithelia exposed to bacteria in an inflammatory bowel disease model repressed colon carcinogenesis (85). Pilarsky et al. found that inhibition of NF- $\kappa$ B by targeted over-expression of I $\kappa$ B signal mutant or treatment with anti-TNF antibodies was associated with a decrease in NF- $\kappa$ B activation and tumorigenesis in mdr-2 knockout mice that develop inflammation-associated hepatocellular carcinomas (86). By contrast, targeted disruption of the classic pathway by IKK $\beta$  knockout in hepatocytes during exposure to carcinogen DEN increased tumorigenesis (87). TNF and c-Jun NH<sub>2</sub>-terminal kinase 1 pathway activations were implicated in necrosis-induced inflammation and promotion of the increase in carcinogenesis in this model (87, 88).

Apparent differences in role of NF- $\kappa$ B have also been observed in human squamous cell carcinoma and among some murine models. Constitutive activation of NF- $\kappa$ B has been found with progression of squamous cell carcinoma derived from skin of BALB/c and  $\alpha$ -catenin knockout mice (89, 90) and in squamous dysplasias and carcinomas of the cervix and the head and neck (15, 17, 91). Inhibition of NF- $\kappa$ B by I $\kappa$ B repressor, steroids, or proteasome inhibitors promotes apoptosis and inhibits tumorigenesis of these cancers *in vitro* and *in vivo* (40, 72, 91, 92). By contrast, in transgenic mouse models targeting squamous epithelia, selective expression of an I $\kappa$ B repressor of NF- $\kappa$ B during carcinogen exposure or with expression of *ras* oncogene resulted in increased carcinogenesis of squamous cell carcinoma of the skin (93, 94). With inhibition of NF- $\kappa$ B in that setting, increased TNF, c-Jun NH<sub>2</sub>-terminal kinase, and activator protein pathway activation were shown to contribute to the increase in carcinogenesis (95). These findings suggest there may be greater complexity and differences in the mechanism of activation of NF- $\kappa$ B during carcinogenesis in human and murine cancers than are thus far reflected in some of these targeted murine models.

Several molecular alterations that occur with cancer development have been identified that could help explain the apparent differences in function of NF- $\kappa$ B in these different contexts. First, most knockout models to date do not account for the multiple positive signal mechanisms activated during cancer development that can contribute to constitutive activation of the classic or alternative NF- $\kappa$ B pathways. As reviewed above, multiple growth factor and cytokine receptors, PI3K, Akt, CK2, IKK $\alpha$  and IKK $\beta$ , other signal kinases, and cofactors may be coactivated in cancer and contribute to transactivation of NF- $\kappa$ B1 and/or NF- $\kappa$ B2. In addition, most cancers exhibit inactivation or loss of expression of tumor suppressors that

have been shown to negatively regulate NF- $\kappa$ B activation. Perkins et al. have shown that in embryonic fibroblasts or malignant cells deficient in functional p14ARF, p53, and proapoptotic gene expression, NF- $\kappa$ B acts as an activator of prosurvival genes, such as *cyclin D1* and *Bcl-XL* (37, 96). However, re-expression and/or activation of functional ARF, ATR, Chk2, and p53 tumor suppressor pathway components can alter NF- $\kappa$ B to become a functional repressor. ARF, ATR, and Chk2 were shown to mediate phosphorylation of Thr<sup>505</sup> of RelA (p65), and p53 was shown to compete for the coactivator CBP/p300, resulting in association with histone deacetylases and inactivation of NF- $\kappa$ B and prosurvival genes. Consistent with this model, others have found that NF- $\kappa$ B overexpression in immortalized keratinocyte cell lines with mutant p53 exhibits increased malignant potential and tumorigenesis (97). Mice with targeted deletion of  $\alpha$ -catenin, another tumor suppressor commonly inactivated in human cancer, also exhibit increased NF- $\kappa$ B activation, inflammation, proliferation, and malignant transformation (90). Together, these observations may help account for the apparent disparities in role of NF- $\kappa$ B in different experimental knockout models and human SCC above because the *p14ARF/p16INK4a* locus encoding p14ARF and p53, and  $\alpha$ -catenin frequently undergo inactivation or mutation in human and murine cancers in which constitutive NF- $\kappa$ B activation has been shown.

## Conclusions

The alterations in multiple signal pathways shown to positively and negatively regulate NF- $\kappa$ B activation are consistent with its frequent activation and important functional role in cancer. The redundancy in alterations that may lead to activation can also explain in part why drugs targeting receptors and kinases further upstream are effective primarily in cancers in which limited alterations and a dominant role of these molecules is established. Agents with broad activities that include inhibition of NF- $\kappa$ B, such as NSAIDs, corticosteroids, and proteasome inhibitors, have already shown evidence of efficacy and safety in prevention or therapy of certain cancers. The important role of CK2, IKKs, ubiquitin ligase  $\beta$ TrCP, and proteasome in integrating these signals and the final steps of activation have made these important molecules for study as potential targets for prevention and therapy.

## Acknowledgments

We thank the many contributors in the Van Waes lab, colleagues in the NF- $\kappa$ B field and their forbearance for oversimplifications or errors I may have made in reviewing the field, and Keith Brown, James Battey, and Patricia Steeg for the review and comments.

## References

- Hayden MS, Ghosh S. Signaling to NF- $\kappa$ B. *Genes Dev* 2004;18:2195–224.
- Luo JL, Kamata H, Karin M. IKK/NF- $\kappa$ B signaling: balancing life and death—a new approach to cancer therapy. *J Clin Invest* 2005;115:2625–32.
- Karin M, Ben-Neriah Y. Phosphorylation meets ubiquitination: the control of NF- $\kappa$ B activity. *Annu Rev Immunol* 2000;18:621–63.
- Baldwin AS. Control of oncogenesis and cancer therapy resistance by the transcription factor NF- $\kappa$ B. *J Clin Invest* 2001;107:241–6.
- Aggarwal BB, Shishodia S, Sandur SK, Pandey MK, Sethi G. Inflammation and cancer: how hot is the link? *Biochem Pharmacol* 2006;72:1605–21. Epub 2006 Aug 4.
- Hiscott J, Kwon H, Genin P. Hostile takeovers: viral appropriation of the NF- $\kappa$ B pathway. *J Clin Invest* 2001;107:143–51.
- Tsurutani J, Castillo SS, Brognard J, et al. Tobacco components stimulate Akt-dependent proliferation and NF- $\kappa$ B-dependent survival in lung cancer cells. *Carcinogenesis* 2005;26:1182–95.
- Huang TT, Wuerzberger-Davis SM, Wu ZH, Miyamoto S. Sequential modification of NEMO/IKK $\gamma$  by SUMO-1 and ubiquitin mediates NF- $\kappa$ B activation by genotoxic stress. *Cell* 2003;115:565–76.
- Bargou RC, Emmerich F, Krappmann D, et al. Consti-

- tive nuclear factor- $\kappa$ B-RelA activation is required for proliferation and survival of Hodgkin's disease tumor cells. *J Clin Invest* 1997;100:2961–9.
10. Sovak MA, Bellas RE, Kim DW, et al. Aberrant nuclear factor- $\kappa$ B/Rel expression and the pathogenesis of breast cancer. *J Clin Invest* 1997;100:2952–60.
  11. Suh J, Payvandi F, Edelstein LC, et al. Mechanisms of constitutive NF- $\kappa$ B activation in human prostate cancer cells. *Prostate* 2002;52:183–200.
  12. Mukhopadhyay T, Roth JA, Maxwell SA. Altered expression of the p50 subunit of the NF- $\kappa$ B transcription factor complex in non-small cell lung carcinoma. *Oncogene* 1995;11:999–1003.
  13. Kojima M, Morisaki T, Sasaki N, et al. Increased nuclear factor- $\kappa$ B activation in human colorectal carcinoma and its correlation with tumor progression. *Anticancer Res* 2004;24:675–81.
  14. Wang W, Abbruzzese JL, Evans DB, Larry L, Cleary KR, Chiao PJ. The nuclear factor- $\kappa$ B RelA transcription factor is constitutively activated in human pancreatic adenocarcinoma cells. *Clin Cancer Res* 1999;5:119–27.
  15. Ondrey FG, Dong G, Sunwoo J, et al. Constitutive activation of transcription factors NF- $\kappa$ B, AP-1, and NF-IL6 in human head and neck squamous cell carcinoma cell lines that express pro-inflammatory and pro-angiogenic cytokines. *Mol Carcinog* 1999;26:119–29.
  16. Abdel-Latif MM, O'Riordan J, Windle HJ, et al. NF- $\kappa$ B activation in esophageal adenocarcinoma: relationship to Barrett's metaplasia, survival, and response to neoadjuvant chemoradiotherapy. *Ann Surg* 2004;239:491–500.
  17. Nair A, Venkatraman M, Maliekal TT, Nair B, Karunagaran D. NF- $\kappa$ B is constitutively activated in high-grade squamous intraepithelial lesions and squamous cell carcinomas of the human uterine cervix. *Oncogene* 2003;22:50–8.
  18. Biswas DK, Cruz AP, Gansberger E, Pardee AB. Epidermal growth factor-induced nuclear factor  $\kappa$ B activation: a major pathway of cell-cycle progression in estrogen-receptor negative breast cancer cells. *Proc Natl Acad Sci U S A* 2000;97:8542–7.
  19. Pianetti S, Arsura M, Romieu-Mourez R, Coffey RJ, Sonenshein GE. Her-2/neu overexpression induces NF- $\kappa$ B via a PI3-kinase/Akt pathway involving calpain-mediated degradation of I $\kappa$ B- $\alpha$  that can be inhibited by the tumor suppressor PTEN. *Oncogene* 2001;20:1287–99.
  20. Wolf JS, Chen Z, Dong G, et al. IL (interleukin)-1 $\alpha$  promotes nuclear factor- $\kappa$ B and AP-1-induced IL-8 expression, cell survival, and proliferation in head and neck squamous cell carcinomas. *Clin Cancer Res* 2001;7:1812–20.
  21. Fan S, Gao M, Meng Q, Laterra JJ, Symons MH, Coniglio S. Role of NF- $\kappa$ B signaling in hepatocyte growth factor/scatter factor-mediated cell protection. *Oncogene* 2005;24:1749–66.
  22. Nikolopoulos SN, Blaikie P, Yoshioka T, Guo W, Giancotti FG. Integrin  $\beta$ 4 signaling promotes tumor angiogenesis. *Cancer Cell* 2004;6:471–83.
  23. Romieu-Mourez R, Landesman-Bollag E, Seldin DC, Traish AM, Mercurio F, Sonenshein GE. Roles of IKK kinases and protein kinase CK2 in activation of nuclear factor- $\kappa$ B in breast cancer. *Cancer Res* 2001;61:3810–8.
  24. Bancroft CC, Chen Z, Yeh J, Sunwoo JB, Yeh NT, Jackson S. Effects of pharmacologic antagonists of epidermal growth factor receptor, PI3K and MEK signal kinases on NF- $\kappa$ B and AP-1 activation and IL-8 and VEGF expression in human head and neck squamous cell carcinoma lines. *Int J Cancer* 2002;99:538–48.
  25. Yu M, Yeh J, Van Waes C. Protein kinase casein kinase 2 mediates inhibitor- $\kappa$ B kinase and aberrant nuclear factor- $\kappa$ B activation by serum factor(s) in head and neck squamous carcinoma cells. *Cancer Res* 2006;66:6722–31.
  26. Farah M, Parhar K, Moussavi M, Eivemark S, Salh B. 5,6-Dichloro-ribifuranosylbenzimidazole- and apigenin-induced sensitization of colon cancer cells to TNF- $\alpha$ -mediated apoptosis. *Am J Physiol Gastrointest Liver Physiol* 2003;285:G919–28.
  27. Tamatani T, Azuma M, Aota K, Yamashita T, Bando T, Sato M. Enhanced I $\kappa$ B kinase activity is responsible for the augmented activity of NF- $\kappa$ B in human head and neck carcinoma cells. *Cancer Lett* 2001;171:165–72.
  28. Gasparian AV, Yao YJ, Kowalczyk D, Lyakh LA, Karseladze A, Slaga TJ. The role of IKK in constitutive activation of NF- $\kappa$ B transcription factor in prostate carcinoma cells. *J Cell Sci* 2002;115:141–51.
  29. Madrid LV, Wang CY, Guttridge DC, Schottelius AJ, Baldwin AS, Jr., Mayo MW. Akt suppresses apoptosis by stimulating the transactivation potential of the RelA/p65 subunit of NF- $\kappa$ B. *Mol Cell Biol* 2000;20:1626–38.
  30. Madrid LV, Mayo MW, Reuther JY, Baldwin AS, Jr. Akt stimulates the transactivation potential of the RelA/p65 Subunit of NF- $\kappa$ B through utilization of the I $\kappa$ B kinase and activation of the mitogen-activated protein kinase p38. *J Biol Chem* 2001;276:18934–40.
  31. Reuther JY, Reuther GW, Cortez D, Pendergast AM, Baldwin AS, Jr. A requirement for NF- $\kappa$ B activation in Bcr-Abl-mediated transformation. *Genes Dev* 1998;12:968–81.
  32. Ho L, Davis RE, Conne B, et al. MALT1 and the API2–1 fusion act between CD40 and IKK and confer NF- $\kappa$ B-dependent proliferative advantage and resistance against FAS-induced cell death in B cells. *Blood* 2005;105:2891–9.
  33. Westerheide SD, Mayo MW, Anest V, Hanson JL, Baldwin AS, Jr. The putative oncoprotein Bcl-3 induces cyclin D1 to stimulate G(1) transition. *Mol Cell Biol* 2001;21:8428–36.
  34. Mathas S, Johrens K, Joos S, et al. Elevated NF- $\kappa$ B p50 complex formation and Bcl-3 expression in classical Hodgkin, anaplastic large-cell, and other peripheral T-cell lymphomas. *Blood* 2005;106:4287–93.
  35. Wood KM, Roff M, Hay RT. Defective I $\kappa$ B $\alpha$  in Hodgkin cell lines with constitutively active NF- $\kappa$ B. *Oncogene* 1998;16:2131–9.
  36. Emmerich F, Meiser M, Hummel M, et al. Overexpression of I $\kappa$ B  $\alpha$  without inhibition of NF- $\kappa$ B activity and mutations in the I $\kappa$ B  $\alpha$  gene in Reed-Sternberg cells. *Blood* 1999;94:3129–34.
  37. Campbell KJ, Perkins ND. Regulation of NF- $\kappa$ B function. *Biochem Soc Symp* 2006;73:165–80.
  38. Asano T, Yao Y, Zhu J, Li D, Abbruzzese JL, Reddy SA. The PI 3-kinase/Akt signaling pathway is activated due to aberrant Pten expression and targets transcription factors NF- $\kappa$ B and c-Myc in pancreatic cancer cells. *Oncogene* 2004;23:8571–80. Erratum in: *Oncogene* 2005;24:4320.
  39. Dong G, Loukinova E, Chen Z, et al. Molecular profiling of transformed and metastatic murine squamous carcinoma cells by differential display and cDNA microarray reveals altered expression of multiple genes related to growth, apoptosis, angiogenesis, and the NF- $\kappa$ B signal pathway. *Cancer Res* 2001;61:4797–808.
  40. Loercher A, Lee TL, Ricker JL, et al. Nuclear factor- $\kappa$ B is an important modulator of the altered gene expression profile and malignant phenotype in squamous cell carcinoma. *Cancer Res* 2004;64:6511–23. Erratum in: *Cancer Res* 2004 Nov 1;64:8130–2.
  41. Chung CH, Parker JS, Ely K, et al. Gene Expression profiles identify epithelial-to-mesenchymal transition and activation of nuclear factor- $\kappa$ B signaling as characteristics of a high-risk head and neck squamous cell carcinoma. *Cancer Res* 2006;66:8210–8.
  42. Hinz M, Lemke P, Anagnostopoulos I, et al. Nuclear factor  $\kappa$ B-dependent gene expression profiling of Hodgkin's disease tumor cells, pathogenetic significance, and link to constitutive signal transducer and activator of transcription 5a activity. *J Exp Med* 2002;196:605–17.
  43. Staudt LM. Gene expression profiling of lymphoid malignancies. *Annu Rev Med* 2002;53:303–18.
  44. Van Laere SJ, Van der Auwera I, Van den Eynden GG, et al. Nuclear factor- $\kappa$ B signature of inflammatory breast cancer by cDNA microarray validated by quantitative real-time reverse transcription-PCR, immunohistochemistry, and nuclear factor- $\kappa$ B DNA-binding. *Clin Cancer Res* 2006;12:3249–56.
  45. Wang CY, Mayo MW, Korneluk RG, Goeddel DV, Baldwin AS, Jr. NF- $\kappa$ B antiapoptosis: induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation. *Science* 1998;281:1680–3.
  46. Guttridge DC, Albanese C, Reuther JY, Pestell RG, Baldwin AS, Jr. NF- $\kappa$ B controls cell growth and differentiation through transcriptional regulation of cyclin D1. *Mol Cell Biol* 1999;19:5785–99.
  47. Chen C, Edelstein LC, Gelinas C. The Rel/NF- $\kappa$ B family directly activates expression of the apoptosis inhibitor Bcl-x(L). *Mol Cell Biol* 2000;20:2687–95.
  48. Richmond A. NF- $\kappa$ B, chemokine gene transcription and tumour growth. *Nat Rev Immunol* 2002;2:664–74.
  49. Yin MJ, Yamamoto Y, Gaynor RB. The anti-inflammatory agents aspirin and salicylate inhibit the activity of I $\kappa$ B kinase- $\beta$ . *Nature* 1998;396:77–80.
  50. Yamamoto Y, Yin MJ, Lin KM, Gaynor RB. Sulindac inhibits activation of the NF- $\kappa$ B pathway. *J Biol Chem* 1999;274:27307–14.
  51. Palayoor ST, Youmell MY, Calderwood SK, Coleman CN, Price BD. Constitutive activation of I $\kappa$ B kinase  $\alpha$  and NF- $\kappa$ B in prostate cancer cells is inhibited by ibuprofen. *Oncogene* 1999;18:7389–94.
  52. Takada Y, Bhardwaj A, Potdar P, Aggarwal BB. Non-steroidal anti-inflammatory agents differ in their ability to suppress NF- $\kappa$ B activation, inhibition of expression of cyclooxygenase-2 and cyclin D1, and abrogation of tumor cell proliferation. *Oncogene* 2004;23:9247–58.
  53. Narayanan BA, Narayanan NK, Pittman B, Reddy BS. Regression of mouse prostatic intraepithelial neoplasia by nonsteroidal anti-inflammatory drugs in the transgenic adenocarcinoma mouse prostate model. *Clin Cancer Res* 2004;10:7727–37.
  54. Sangha S, Yao M, Wolfe MM. Non-steroidal anti-inflammatory drugs and colorectal cancer prevention. *Postgrad Med J* 2005;81:223–7.
  55. Hoffmeister M, Chang-Claude J, Brenner H. Do older adults using NSAIDs have a reduced risk of colorectal cancer? *Drugs Aging* 2006;23:513–23.
  56. Blair CK, Sweeney C, Anderson KE, Folsom AR. NSAID use and survival after breast cancer diagnosis in post-menopausal women. *Breast Cancer Res Treat* 2007;101:191–7. Epub 2006 Jun 6.
  57. Pruthi RS, Derksen JE, Moore D, et al. Phase II trial of celecoxib in prostate-specific antigen recurrent prostate cancer after definitive radiation therapy or radical prostatectomy. *Clin Cancer Res* 2006;12:2172–7.
  58. Aggarwal BB, Shishodia S. Molecular targets of dietary agents for prevention and therapy of cancer. *Biochem Pharmacol* 2006;71:1397–421.
  59. Aggarwal S, Takada Y, Singh S, Myers JN, Aggarwal BB. Inhibition of growth and survival of human head and neck squamous cell carcinoma cells by curcumin via modulation of nuclear factor- $\kappa$ B signaling. *Int J Cancer* 2004;111:679–92. Erratum in: *Int J Cancer* 2004;112:1086.
  60. Nakanishi C, Toi M. Nuclear factor- $\kappa$ B inhibitors as sensitizers to anticancer drugs. *Nat Rev Cancer* 2005;5:297–309.
  61. Hideshima T, Chauhan D, Richardson P, et al. NF- $\kappa$ B as a therapeutic target in multiple myeloma. *J Biol Chem* 2002;277:16639–47.
  62. Campbell KJ, Witty JM, Rocha S, Perkins ND. Cisplatin mimics ARF tumor suppressor regulation of RelA (p65) nuclear factor- $\kappa$ B transactivation. *Cancer Res* 2006;66:929–35.
  63. Richardson PG, Mitsiades C, Hideshima T, Anderson

- KC. Bortezomib: proteasome inhibition as an effective anticancer therapy. *Annu Rev Med* 2006;57:33–47.
64. O'Connor OA, Wright J, Moskowitz C, et al. Phase II clinical experience with the novel proteasome inhibitor bortezomib in patients with indolent non-Hodgkin's lymphoma and mantle cell lymphoma. *J Clin Oncol* 2005;23:676–84.
65. Chen CI, Kouroukis T, White D, et al. Bortezomib is active in Waldenstrom's macroglobulinemia (WM): results of a National Cancer Institute of Canada (NCIC) phase II study in previously untreated WM. *J Clin Oncol Suppl* 2006;24:7543.
66. Papandreou CN, Daliani DD, Nix D, et al. Phase I trial of the proteasome inhibitor bortezomib in patients with advanced solid tumors with observations in androgen-independent prostate cancer. *J Clin Oncol* 2004;22:2108–21.
67. Davis NB, Taber DA, Ansari RH, et al. Phase II trial of PS-341 in patients with renal cell cancer: a University of Chicago phase II consortium study. *J Clin Oncol* 2004;22:115–9.
68. Kondagunta GV, Drucker B, Schwartz L, et al. Phase II trial of bortezomib for patients with advanced renal cell carcinoma. *J Clin Oncol* 2004;22:3720–5.
69. Markovic SN, Geyer SM, Dawkins F, et al. A phase II study of bortezomib in the treatment of metastatic malignant melanoma. *Cancer* 2005;103:2584–9.
70. Mackay H, Hedley D, Major P, et al. A phase II trial with pharmacodynamic endpoints of the proteasome inhibitor bortezomib in patients with metastatic colorectal cancer. *Clin Cancer Res* 2005;11:5526–33.
71. Yang CH, Gonzalez-Angulo AM, Reuben JM, et al. Bortezomib (VELCADE) in metastatic breast cancer: pharmacodynamics, biological effects, and prediction of clinical benefits. *Ann Oncol* 2006;17:813–7.
72. Van Waes C, Chang AA, Lebowitz PF, et al. Inhibition of nuclear factor-kappaB and target genes during combined therapy with proteasome inhibitor bortezomib and reirradiation in patients with recurrent head-and-neck squamous cell carcinoma. *Int J Radiat Oncol Biol Phys* 2005;63:1400–12.
73. Lara PN, Gumerlock PH, Crowley J, Gandara DR. Bortezomib and gemcitabine/carboplatin results in encouraging survival in advanced non-small cell lung cancer: results of a phase II Southwest oncology group trial (SO339). *J Clin Oncol Suppl* 2006;24:368s. Abstract 7017.
74. Aghajanian C, Dizon DS, Sabbatini P, Raizer JJ, Dupont J, Spriggs DR. Phase I trial of bortezomib and carboplatin in recurrent ovarian or primary peritoneal cancer. *J Clin Oncol* 2005;23:5943–9.
75. Ma C, Mandrekar SJ, Alberts SR, et al. A phase I and pharmacologic study of sequences of the proteasome inhibitor, bortezomib (PS-341, Velcade™), in combination with paclitaxel and carboplatin in patients with advanced malignancies. *Cancer Chemother Pharmacol* 2007;59:207–15. Epub 2006 Jun 9.
76. Messersmith WA, Baker SD, Lassiter L, et al. Phase I trial of bortezomib in combination with docetaxel in patients with advanced solid tumors. *Clin Cancer Res* 2006;12:1270–5.
77. Karin M, Yamamoto Y, Wang QM. The IKK NF-kappa B system: a treasure trove for drug development. *Nat Rev Drug Discov* 2004;3:17–26.
78. Lam LT, Davis RE, Pierce J, et al. Small molecule inhibitors of I kappa B kinase are selectively toxic for subgroups of diffuse large B-cell lymphoma defined by gene expression profiling. *Clin Cancer Res* 2005;11:28–40.
79. Cillonì D, Messa F, Arruga F, et al. The NF-kappaB pathway blockade by the IKK inhibitor PS1145 can overcome imatinib resistance. *Leukemia* 2006;20:61–7.
80. Yemelyanov A, Gasparian A, Lindholm P, et al. Effects of IKK inhibitor PS1145 on NF-kappaB function, proliferation, apoptosis and invasion activity in prostate carcinoma cells. *Oncogene* 2006;25:387–98.
81. Yang J, Amiri KI, Burke JR, Schmid JA, Richmond A. BMS-345541 targets inhibitor of kappaB kinase and induces apoptosis in melanoma: involvement of nuclear factor kappaB and mitochondria pathways. *Clin Cancer Res* 2006;12:950–60.
82. Anest V, Cogswell PC, Baldwin AS, Jr. I kappa B kinase alpha and p65/RelA contribute to optimal epidermal growth factor-induced c-fos gene expression independent of I kappa Balpha degradation. *J Biol Chem* 2004;279:31183–9.
83. Bernal-Mizrachi L, Lovly CM, Ratner L. The role of NF- $\kappa$ B-1 and NF- $\kappa$ B-2-mediated resistance to apoptosis in lymphomas. *Proc Natl Acad Sci U S A* 2006;103:9220–5.
84. Matta H, Chaudhary PM. Activation of alternative NF-kappa B pathway by human herpes virus 8-encoded Fas-associated death domain-like IL-1 beta-converting enzyme inhibitory protein (vFLIP). *Proc Natl Acad Sci U S A* 2004;101:9399–404.
85. Luo JL, Maeda S, Hsu LC, Yagita H, Karin M. Inhibition of NF-kappaB in cancer cells converts inflammation-induced tumor growth mediated by TNFalpha to TRAIL-mediated tumor regression. *Cancer Cell* 2004;6:297–305.
86. Pikarsky E, Porat RM, Stein I, et al. NF-kappaB functions as a tumour promoter in inflammation-associated cancer. *Nature* 2004;431:461–6.
87. Maeda S, Kamata H, Luo JL, Leffert H, Karin M. IKKbeta couples hepatocyte death to cytokine-driven compensatory proliferation that promotes chemical hepatocarcinogenesis. *Cell* 2005;121:977–90.
88. Sakurai T, Maeda S, Chang L, Karin M. Loss of hepatic NF-kappaB activity enhances chemical hepatocarcinogenesis through sustained c-Jun N-terminal kinase 1 activation. *Proc Natl Acad Sci U S A* 2006;103:10544–51.85.
89. Dong G, Chen Z, Kato T, Van Waes C. The host environment promotes the constitutive activation of nuclear factor-kappaB and proinflammatory cytokine expression during metastatic tumor progression of murine squamous cell carcinoma. *Cancer Res* 1999;59:3495–504.
90. Kobiela A, Fuchs E. Links between alpha-catenin, NF-kappaB, and squamous cell carcinoma in skin. *Proc Natl Acad Sci U S A* 2006;103:2322–7.
91. Duffey DC, Chen Z, Dong G, et al. Expression of a dominant-negative mutant inhibitor-kappaBalpha of nuclear factor-kappaB in human head and neck squamous cell carcinoma inhibits survival, proinflammatory cytokine expression, and tumor growth *in vivo*. *Cancer Res* 1999;59:3468–74.
92. Sunwoo JB, Chen Z, Dong G, et al. Novel proteasome inhibitor PS-341 inhibits activation of nuclear factor-kappa B, cell survival, tumor growth, and angiogenesis in squamous cell carcinoma. *Clin Cancer Res* 2001;7:1419–28.
93. van Hogerlinden M, Rozell BL, Ahrlund-Richter L, Toftgard R. Squamous cell carcinomas and increased apoptosis in skin with inhibited Rel/nuclear factor-kappaB signaling. *Cancer Res* 1999;59:3299–303.
94. Dajee M, Lazarov M, Zhang JY, et al. NF-kappaB blockade and oncogenic Ras trigger invasive human epidermal neoplasia. *Nature* 2003;421:639–43.
95. Zhang JY, Green CL, Tao S, Khavari PA. NF-kappaB RelA opposes epidermal proliferation driven by TNFR1 and JNK. *Genes Dev* 2004;18:17–22.
96. Perkins ND, Gilmore TD. Good cop, bad cop: the different faces of NF-kappaB. *Cell Death Differ* 2006;13:759–72.
97. Ren Q, Kari C, Quadros MR, et al. Malignant transformation of immortalized HaCaT keratinocytes through deregulated nuclear factor kappaB signaling. *Cancer Res* 2006;66:5209–15.