

# Fish Oil Inhibits Human Lung Carcinoma Cell Growth by Suppressing Integrin-Linked Kinase

ShouWei Han,<sup>1</sup> XiaoJuan Sun,<sup>1</sup> Jeffrey D. Ritzenthaler,<sup>1</sup> and Jesse Roman<sup>1,2</sup>

<sup>1</sup>Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, Emory University School of Medicine; <sup>2</sup>Atlanta Veterans Affairs Medical Center, Atlanta, Georgia

## Abstract

We previously showed that synthetic peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) ligands inhibit non-small cell lung carcinoma (NSCLC) cell growth through multiple signaling pathways. Here, we show that dietary compounds, such as fish oil (which contains certain kinds of fatty acids like  $\omega$ 3 and  $\omega$ 6 polyunsaturated fatty acids), also inhibit NSCLC cell growth by affecting PPAR $\gamma$  and by inhibiting the expression of integrin-linked kinase (ILK). Exogenous expression of ILK overcame, whereas silencing ILK enhanced the inhibitory effect of fish oil on cell growth. The inhibitor of p38 mitogen-activated protein kinase, SB239023, abrogated the inhibitory effect of fish oil on ILK expression, whereas the inhibitor of extracellular signal-regulated kinase, PD98059, had no effect. Transient transfection experiments showed that fish oil reduced ILK promoter activity, and this effect was abolished by AP-2 $\alpha$  small interfering RNA and SB239023 and by deletion of a specific portion of the *ILK* gene promoter. Western blot analysis and gel mobility shift assay showed that fish oil significantly induced AP-2 $\alpha$  protein expression and AP-2 DNA-binding activity in the *ILK* gene promoter and that this was dependent on PPAR $\gamma$  activation. Blockade of AP-2 $\alpha$  abrogated the effect of fish oil on ILK expression and on cell growth, whereas exogenous expression of AP-2 $\alpha$  enhanced cell growth in the setting of fish oil exposure. Taken together, these findings show that fish oil inhibits ILK expression through activation of PPAR $\gamma$ -mediated and p38 mitogen-activated protein kinase-mediated induction of AP-2 $\alpha$ . In turn, this leads to inhibition of NSCLC cell proliferation. This study unveils a novel mechanism by which fish oil inhibits human lung cancer cell growth. (*Mol Cancer Res* 2009;7(1):108–17)

## Introduction

Lung cancer is the leading cause of cancer death in the world with an estimated 213,000 new cases and 160,000 deaths in 2007 in the United States (1). Despite marked improvements in understanding the molecular biology of cancer and in advances in diagnostics, surgery, and chemotherapy, the 5-year poor survival rate (<15%) has not changed substantially (1, 2). New approaches to the treatment and prevention of lung cancer are clearly indicated but depend greatly on a better understanding of the molecular mechanisms that control oncogenesis and tumor growth in lung. Fish oils are enriched with polyunsaturated fatty acids (PUFA) of the  $\omega$ 3 family and are believed to have protective effects against cancer (3). Several investigators have shown that diets enriched with fish oil can reduce the growth rates of implanted tumors *in vivo* (4–6). However, the mechanisms responsible for the anticancer effects of fish oil remain incompletely elucidated.

Both *n*-6 PUFAs and *n*-3 PUFAs modulate peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) and decrease cell growth in human lung cancer cells (7). PPAR $\gamma$  is a member of the ligand-inducible nuclear transcription factors that heterodimerize with retinoid X receptors and bind to peroxisome proliferator response elements located in the promoter region of PPAR target genes (8). These lipid-sensitive receptors can be activated in a variable isotype-specific manner by natural/dietary ligands, including long-chain PUFAs, which are found in fish oil (e.g., *n*-3 PUFA and *n*-6 PUFA), various eicosanoids (e.g., 15-d-PGJ2), lipid hydroperoxides [e.g., 9(*S*)-HODE and 13(*S*)-HODE], and linoleic acid (9, 10). The efficacy of these compounds as anticancer agents has been examined in a variety of cancers, including colon, breast, and prostate, and they have been found to inhibit cancer cell growth *in vitro* and *in vivo* (11).

In view of the above, it has been suggested that the anticancer properties of fish oil are dependent on activation of PPAR $\gamma$ ; however, the downstream events involved in this process remain unclear. One of the potential targets for PPAR $\gamma$  ligands is integrin-linked kinase (ILK), which links cell adhesion receptors, integrins, and growth factors to the actin cytoskeleton and to a range of signaling pathways that are implicated in the regulation of anchorage-dependent cell growth/survival, cell cycle progression, invasion and migration, and tumor angiogenesis (12). Moreover, overexpression of ILK results in oncogenic transformation and progression to invasive and metastatic phenotypes (13, 14). Thus, we explored the effects of fish oil on ILK expression. This work revealed that fish oil inhibits non-small cell lung carcinoma (NSCLC) proliferation by suppressing ILK expression through activation of PPAR $\gamma$ . This results in the activation of p38

Received 8/15/08; revised 10/3/08; accepted 10/30/08.

**Grant support:** American Thoracic Society/LUNGevity Foundation Partnership grant LC-06-004, Emory University Research Fund 2-55016, and NIH/National Cancer Institute grant CA123104 (S. Han) and Department of Veterans Affairs Merit Review Grant and NIH/National Cancer Institute grant CA116812 (J. Roman).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Requests for reprints:** ShouWei Han, Division of Pulmonary, Allergy and Critical Care Medicine, Emory University School of Medicine, Whitehead Bioresearch Building, 615 Michael Street, Suite 205-M, Atlanta, GA 30322. Phone: 404-712-2661; Fax: 404-712-2151. E-mail: shan2@emory.edu

Copyright © 2009 American Association for Cancer Research.  
doi:10.1158/1541-7786.MCR-08-0384

mitogen-activated protein kinase (p38 MAPK) and induction of AP-2 $\alpha$ , which in turn inhibits *ILK* gene expression. To our knowledge, this is the first report linking fish oil to ILK expression.

## Results

### *Fish Oil Decreases NSCLC Cell Proliferation, Which Is Associated with Inhibition of ILK Protein Expression, through PPAR $\gamma$ Activation*

We began by examining the effects of fish oil on NSCLC cell proliferation. As shown in Fig. 1A, fish oil resulted in a significant decrease in cell proliferation of two NSCLC cell lines (H1838 and H1792) compared with the control cells (no fish oil) with maximal effects noted at 15  $\mu$ g/mL. Note that two human bronchial epithelial cell lines (BEAS-2B and 16-HBE) showed no inhibition in response to fish oil (Fig. 1A).

The ILK signal pathway has been implicated in the regulation of anchorage-dependent cell growth/survival, cell cycle progression, invasion and migration, and tumor angiogenesis (15, 16). We found that fish oil inhibited ILK protein levels in a dose-dependent manner in H1838 NSCLC cells (Fig. 1B). Similar results were also observed in several other NSCLC cell lines (Fig. 1C), whereas fish oil had little effect on ILK protein expression in BEAS-2B, 16-HBE, and NIH3T3 fibroblast cells (Fig. 1D).

Because fish oil can activate nuclear receptors such as PPAR $\gamma$ , we examined if inhibition of ILK expression by fish oil was mediated by PPAR $\gamma$ . To this end, we tested the cells with GW9662, a specific PPAR $\gamma$  antagonist. As shown in Fig. 1E, the effects of fish oil on ILK expression were eliminated in the presence of GW9662, and this was confirmed by PPAR $\gamma$  small interfering RNA (siRNA; Fig. 1F). Note that fish oil induced PPAR $\gamma$  protein expression (Fig. 1F); this effect was due to up-regulation of PPAR $\gamma$  gene transcription because fish oil stimulated PPAR $\gamma$  gene promoter activity in separate experiments (data not shown). Similar results were also observed with another PPAR $\gamma$  natural activator, linoleic acid (Fig. 1E).

### *Overexpression of ILK Overcomes, whereas Silencing of ILK Enhances the Inhibitory Effect of Fish Oil on Cell Growth*

Having established that fish oil may act via affecting expression of ILK, we tested the role of ILK further. Specifically, we tested if overexpression of ILK blocked the effect of fish oil on cell growth. We transiently transfected cells with ILK kinase-inactive (ILK-S343A) and ILK kinase-hyperactive (ILK-S343D) cDNAs. Exogenous high ILK protein expression was confirmed by Western blot with anti-ILK antibody in ILK-S343A and ILK-S343D treatment groups (Fig. 2A, *top*). Cell viability assays showed that ILK kinase-hyperactive (ILK-S343D) cDNA opposed the inhibitory effect of fish oil on cell growth (Fig. 2A, *bottom*), whereas the inactive ILK slightly enhanced the effect of fish oil. Knockdown of ILK with siRNA reduced ILK protein production (Fig. 2B, *top*) and enhanced the inhibitory effect of fish oil on cell growth as determined by CellTiter-Glo Luminescent Cell Viability Assay (Fig. 2B, *bottom*).

### *The Inhibitor of p38 MAPK, SB239023, Abrogates the Effect of Fish Oil on ILK, whereas the Inhibitor of Extracellular Signal-Regulated Kinase, PD98059, Has No Effect*

We and others have shown that PPAR $\gamma$  ligands activate kinase signaling related to p38 MAPK and extracellular signal-regulated kinase 1/2 (ERK1/2) in several cell systems, including lung cancer (17-19). Therefore, we tested whether regulation of ILK expression by fish oil was related to p38 MAPK and ERK activation. We found that fish oil increased the phosphorylation of p38 MAPK (Fig. 3A) and ERK (Fig. 3B). However, the inhibitors of p38 MAPK, SB239023 and SB203580 (data not shown), blocked the effect of fish oil on ILK protein expression (Fig. 3C) and cell growth (data not shown). The ERK1/2 inhibitor PD98059 had no effect (Fig. 3D). This indicated that activation of p38 MAPK, but not that of ERK, was involved in the regulation of ILK by fish oil.

### *Fish Oil Reduces ILK Promoter Activity and This Is Abolished by the p38 MAPK Inhibitor*

We next examined whether the effects of fish oil on ILK expression occurred at the transcriptional level. As shown in Fig. 4A, the ILK promoter contains multiple transcription factor binding sites, including AP-2, nuclear factor- $\kappa$ B (NF- $\kappa$ B), and Sp1. We found that H1838 cells transfected with a wild-type ILK promoter luciferase reporter construct showed decreased promoter activity when exposed to fish oil, linoleic acid, and rosiglitazone, a synthetic PPAR $\gamma$  ligand, whereas little effect was observed in cells exposed to WY14643, a PPAR $\alpha$  ligand (Fig. 4B). The inhibitory effect of fish oil on ILK promoter activity was abolished by the inhibitor of p38 MAPK, SB239023 (Fig. 4C).

### *Fish Oil Induces AP-2 DNA-Binding Activity in the ILK Promoter through Activation of PPAR $\gamma$*


Having established the importance of ILK, we turned our attention to transcription factors affecting its expression. Transcription factors AP-2 and NF- $\kappa$ B have been inversely correlated with tumor cell growth. PPAR $\gamma$  ligand inhibition of cancer cell proliferation has been associated with inactivation of NF- $\kappa$ B and activation of AP-2 in several studies (20, 21). We found that H1838 cells treated with fish oil (10  $\mu$ g/mL) for 24 hours show increased AP-2 nuclear protein binding activity compared with the solvent controls (Fig. 5A). In contrast, fish oil slightly decreased Sp1 and NF- $\kappa$ B DNA-binding activity (Fig. 5B and C). The specific bands for AP-2, Sp1, and NF- $\kappa$ B were attenuated by a 100-fold molar excess of unlabeled respective oligonucleotides but were not inhibited by a mutated unlabeled AP-2, Sp1, or NF- $\kappa$ B oligonucleotides (mAP-2, mSp1, and mNF- $\kappa$ B). Consistent with the role of PPAR $\gamma$ , we found that GW9662 blocked the stimulatory effect of fish oil on AP-2 DNA-binding activity (Fig. 5D). Moreover, consistent with the stimulatory effect of fish oil on PPAR $\gamma$  protein expression (Fig. 1F), we showed that fish oil enhanced protein complex formation between AP-2 $\alpha$  and PPAR $\gamma$  (Fig. 5E).

### *Blockade of AP-2 $\alpha$ Abrogates the Effect of Fish Oil on ILK Expression and on Cell Growth*

To further test the role of AP-2 in mediating the effect of fish oil on ILK expression, we examined the effect of fish oil on

AP-2 family protein levels. We found that fish oil increased the expression of AP-2 $\alpha$  subunits but had no effect on AP-2 $\gamma$  and AP-2 $\beta$  (Fig. 6A). The stimulatory effect of fish oil on AP-2 $\alpha$  protein expression was blocked by GW9662

(Fig. 6B) and by the inhibitor of p38 MAPK (Fig. 6C). In addition, the inhibitory effect of fish oil on ILK promoter activity was abolished in cells with silencing of AP-2 $\alpha$  (Fig. 6D) and with a deletion ILK promoter construct most



**FIGURE 1.** Fish oil decreases NSCLC cell growth and ILK protein expression through PPAR $\gamma$ . **A.** NSCLC cells (H1838 and H1792) and normal bronchial epithelial cells (BEAS-2B and 16-HBE) were plated in different concentrations of fish oil–enriched medium for up to 48 h. Afterwards, viable cells were detected using CellTiter-Glo Luminescent Cell Viability Assay kit according to the protocol of the manufacturer. Actin was used as internal control for loading purposes. Con, untreated control cells. **B.** Cellular protein was isolated from H1838 cells that were cultured with increased concentrations of fish oil as indicated for 24 h followed by Western blot analysis with antibodies against ILK protein. Actin served as internal control for normalization purposes. **C.** Cellular protein was isolated from several NSCLC cell lines (H1838, A549, H522, and H1792) that were cultured with fish oil (10  $\mu$ g/mL) for up to 24 h followed by Western blot analysis with antibodies against ILK protein. **D.** Cellular protein was isolated from normal epithelial cells (BEAS-2B and 16-HBE) and NIH3T3 cells that were cultured with fish oil (10  $\mu$ g/mL) for up to 24 h followed by Western blot analysis with antibodies against ILK protein. **E.** Cellular protein was isolated from H1838 cells treated with GW9662 (20  $\mu$ mol/L) for 2 h before exposure of the cells to linoleic acid (10  $\mu$ mol/L) or fish oil (10  $\mu$ g/mL) for an additional 24 h and then subjected to Western blot analysis with antibodies against ILK. **F.** Cellular protein was isolated from H1838 cells transfected with control or PPAR $\gamma$  siRNA (100 nmol/L each) for 30 h before exposure of the cells to fish oil–enriched (10  $\mu$ g/mL) medium for an additional 24 h. Afterwards, Western blot analysis was done using polyclonal antibodies against PPAR $\gamma$  and ILK. Actin served as internal control for normalization purposes.

likely lacking AP-2 sites (Fig. 6E). This suggests that inhibition of ILK expression by fish oil occurs at the transcriptional level through activation of p38 MAPK and induction of AP-2 $\alpha$  in NSCLC cells.


Next, we assessed the role of AP-2 $\alpha$  in mediating the inhibitory effect of fish oil on lung cancer cell growth. We showed that cells transfected with AP-2 $\alpha$  siRNA diminished the inhibitory effects of fish oil on cell growth as determined by CellTiter-Glo Luminescent Cell Viability Assay (Fig. 6F, *bottom*). The control siRNA had no effect. Note that AP-2 $\alpha$  siRNA blocked protein production of AP-2 $\alpha$  (Fig. 6F, *top*). On the contrary, cells transfected with AP-2 expression vector [SP(RSV)AP-2] showed enhanced inhibition of the effect of fish oil on cell proliferation (Fig. 6G, *bottom*). Note that cells transfected with [SP(RSV)AP-2] showed increased AP-2 protein levels, whereas the control vector had no effect (Fig. 6G, *top*).

## Discussion


The expression of PPAR $\gamma$  and the effects of dietary PPAR $\gamma$  ligands, such as fish oil, on cell growth have been extensively studied in many carcinoma cell types, including lung cancer (3, 7). However, the exact mechanisms mediating the effects of dietary PPAR $\gamma$  ligands on cell growth inhibition are not fully understood. Fish oil, which is an activator of PPAR $\gamma$ , contains PUFAs of the  $\omega$ 3 and  $\omega$ 6 family and is believed to have anticancer properties (4-6). In this study, we show that fish oil inhibits human lung carcinoma cell growth through suppression of *ILK* gene expression, suggesting that ILK represents a target for PPAR $\gamma$  ligands. Interestingly, fish oil had little effect on the growth of normal cells and this seems to correlate with the relative lower PPAR $\gamma$  protein levels found in normal cells.

The concentrations of fish oil used in this study are similar or even lower than those reported by others showing inhibition of cell growth and invasion (22, 23). One study found that fish oil at a dose range between 1 and 100  $\mu$ g/mL significantly inhibited growth of human hepatoma cells (22). The maximal dose of fish oil tolerated by patients seems to be  $\sim$ 6 g/d (24). Therefore, the concentrations of fish oil used here are much lower than the 1.2 mg/mL found in human blood in those studies.

Alterations in integrin function and ILK expression enhance the aggressive capabilities of several tumor cells (25). Furthermore, ILK overexpression has been correlated with tumor progression (13, 26). ILK is a ubiquitously expressed serine-threonine protein kinase capable of interacting with the cytoplasmic domain of the integrin  $\beta$ 1 and plays a critical role in integrin-dependent cell adhesion, spreading, and cell shape change (12). ILK has been implicated in the regulation of cell growth/survival, cell cycle progression, invasion, and tumor angiogenesis (15, 16). Overexpression of ILK results in oncogenic transformation and progression to invasive and metastatic phenotypes (13, 14). ILK and the activated form of phosphatidylinositol 3-kinase downstream signal Akt are mutually associated with poor prognosis in NSCLC, and the simultaneous overexpression of these proteins is an independent prognostic factor (27). Thus, ILK-pAkt signaling may provide a novel prognostic marker and therapeutic target for NSCLC (27). We found that knockdown of ILK resulted in further inhibition of cell growth, whereas exogenous expression of ILK blocked the inhibitory effect of fish oil on cell growth. Interestingly, silencing of ILK was not as effective as fish oil in inhibiting tumor cell growth. This is probably due to the fact that other kinases, such as Akt and ERK, tumor promoters, such as prostaglandin E $_2$ /cyclooxygenase-2, and



**FIGURE 2.** The role of ILK in mediating the inhibitory effect of fish oil on cell growth. **A.** H1838 cells were transfected with the inactive (ILK-S343A) and hyperactive ILK cDNA (ILK-S343D) using the Oligofectamine reagent according to the manufacturer's instructions. After 24 h of incubation, cells were treated with or without fish oil (15  $\mu$ g/mL) for an additional 24 h. Afterwards, viable cells were detected using CellTiter-Glo Luminescent Cell Viability Assay kit according to the protocol of the manufacturer. Western blot results for ILK protein. Actin was used as internal control for loading purposes. **B.** H1838 cells were transfected with control or ILK siRNA (100 nmol/L) for 40 h followed by exposing the cells to fish oil (15  $\mu$ g/mL) for an additional 24 h. Afterwards, viable cells were detected using CellTiter-Glo Luminescent Cell Viability Assay kit according to the protocol of the manufacturer. Western blot results for ILK protein. Actin was used as internal control for loading purposes.



**FIGURE 3.** The role of p38 MAPK and ERK signaling in ILK protein expression in the setting of fish oil treatment. **A.** Cellular protein was isolated from H1838 cells treated with fish oil in the indicated time period. Afterwards, Western blot analysis was done with antibodies against phospho-p38 MAPK and total p38 MAPK. Actin was used as internal control for loading purposes. **B.** Cellular protein was isolated from H1838 cells treated with fish oil in the indicated time period. Afterwards, Western blot analysis was done with antibodies against phospho-ERK1/2 and total ERK1/2. Actin was used as internal control for loading purposes. **C.** Cellular protein was isolated from H1838 cells treated with SB239063 (10 µmol/L) for 2 h before exposure of the cells to fish oil-enriched medium (10 µg/mL) for an additional 24 h and then subjected to Western blot analysis for ILK protein. **D.** Cellular protein was isolated from H1838 cells treated with PD98059 (25 µmol/L) for 2 h before exposure of the cells to fish oil-enriched medium (10 µg/mL) for an additional 24 h and then subjected to Western blot analysis for ILK protein. Actin was used as internal control for loading purposes.

extracellular matrixes, such as fibronectin, may also drive NSCLC cell growth (28, 29). These effects may be unrelated to ILK but affected by fish oil.

Consistent with other reports (7, 30), we found that the effect of fish oil on the expression of ILK was dependent on activation of PPAR $\gamma$ . In other work, eicosapentaenoic acid (EPA), a component of fish oil, induced a PPAR response element reporter assay and stimulated cell growth in colon cancer cells, which was inhibited by GW9662, a PPAR $\gamma$  antagonist. In contrast, overexpression of PPAR $\gamma$  enhanced the effect of EPA (30). However, PPAR $\gamma$ -independent mechanisms have also been reported (31). These discrepancies may be due to the different cells studied and the concentrations used for these reagents.

ILK has been linked to p38 MAPK and ERK signaling pathways in many cell systems (25, 32). We showed that inhibition of p38 MAPK, but not ERK, blocked the inhibitory effect of fish oil on ILK. Consistent with this, another study showed that ILK played an important role in interleukin-1 $\alpha$ -induced adhesion and invasion of pancreatic cancer cells through p38 MAPK signaling (25). Thus, our results suggest that ILK is downstream of p38 MAPK.

Having shown the important role of ILK, we investigated whether down-regulation of ILK by fish oil reflected inhibition of transactivation of the ILK gene. Our results show that fish oil decreased ILK promoter activity through activation of p38 MAPK. Several transcription factor binding sites within regions of the ILK promoter have been characterized, including

regulatory elements for AP-2, NF- $\kappa$ B, and Sp1 (33). As such, we evaluated the possibility that these sites might play a role in inhibition of ILK expression by fish oil. The effect of fish oil on ILK promoter activity was blocked when the upstream portion of the ILK gene promoter containing AP-2 sites was deleted and in cells transfected with AP-2 $\alpha$  siRNA, suggesting that AP-2 $\alpha$  plays a critical role in mediating the effect of fish oil on ILK gene expression. Furthermore, increased AP-2 $\alpha$  expression and AP-2 $\alpha$  DNA-binding activity were responsible for the inhibitory effect of fish oil on ILK expression. On the contrary, silencing AP-2 $\alpha$  blocked the inhibitory effect of fish oil on NSCLC cell proliferation, further indicating the tumor-suppressive property of this transcription factor. Similarly, another study showed that targeting constitutive expression of AP-2 $\alpha$  in colon cancer cells with siRNA resulted in an increase in their invasive potential, down-regulation of E-cadherin, and increased expression of matrix metalloproteinase-9 (34). The AP-2 transcription factor is involved in the regulation of cell proliferation, differentiation, apoptosis, and carcinogenesis. AP-2 $\alpha$  functions as a tumor suppressor in several cancers (35). Overexpression of AP-2 $\alpha$  significantly decreased cell proliferation and invasion concomitant to the up-regulation of p27 expression in pancreatic cancer cells (36). In addition, transient expression of AP-2 $\alpha$  resulted in the induction of PTEN transcription in AP-2 $\alpha$ -negative colon cancer cells (37).

Taken together, our results show that fish oil inhibits ILK expression through PPAR $\gamma$  signaling that results in activation of p38 MAPK and induction of AP-2 $\alpha$  protein expression


followed by AP-2 DNA-binding activity. In turn, the inhibition of ILK expression results in inhibition of NSCLC cell proliferation. This study unveils a novel mechanism by which fish oil inhibits human lung cancer cell growth.

## Materials and Methods


### Culture, Chemicals, and Fish Oil Treatment

The human NSCLC cell lines (H522, H1792, H1838, and A549), normal bronchial epithelial cell lines (BEAS-2B and 16-HBE), and NIH3T3 cells were obtained from the American Type Culture Collection and routinely grown in RPMI 1640 supplemented with 10% heat-inactivated fetal bovine serum, HEPES buffer, 50 IU/mL penicillin/streptomycin, and 1  $\mu$ g amphotericin (complete medium) as previously described (38). Fish oil was obtained from Sigma. As described in detail elsewhere (39), the fish oil was obtained from Mendaden fish and is typically composed of the following fatty acids: 14:0 myristic acid (6–9%), 16:0 palmitic acid (15–20%), 16:1 palmitoleic acid (9–14%), 18:0 stearic acid (3–4%), 18:1 oleic acid (5–12%), 18:2 linoleic acid (<3%), 18:3 linolenic acid (<3%), 18:4 octadecatetraenoic acid (2–4%), 20:4 arachidonic acid (<3%), 20:5 EPA (10–15%), and 22:6 docosahexaenoic acid (8–15%). These sum components added up to 80% (the

remaining 20% represents other unidentified fatty acids). Note that EPA and docosahexaenoic acid  $\omega$ 3 fatty acids, known to act as PPAR $\gamma$  activators, are the major components for fish oil. The fish oil was emulsified with 5% (w/w) egg phosphatidylcholine (Sigma) and 0.03% (w/w) butylated hydroxytoluene (as antioxidant) in PBS at a final oil concentration of 15 mg/mL. For treatment, cells were plated in regular growth medium for 24 h, after which the medium was replaced with fish oil-enriched medium at indicated concentrations of fish oil. The control group received egg phosphatidylcholine alone at an equal volume. The oil-enriched medium was changed daily to decrease the reactions of fatty acid oxidation on cells. Polyclonal antibodies specific for ILK, p38 MAPK, phospho-p38 MAPK (Thr<sup>180</sup>/Tyr<sup>182</sup>), PD98059, and SB239023 were purchased from Cell Signaling. GW9662, rosiglitazone, and WY14643 were purchased from Cayman Chemical Co. Polyclonal antibodies against AP-2 $\alpha$ , AP-2 $\beta$ , and AP-2 $\gamma$  were purchased from Santa Cruz Biotechnology, Inc. The inactive (ILK-S343A) and superactive ILK cDNAs (ILK-S343D) in pUSEamp vectors under the control of the cytomegalovirus promoter were purchased from Upstate USA, Inc. Linoleic acid, antibodies against PPAR $\gamma$ , and other chemicals were purchased from Sigma-Aldrich unless otherwise indicated.



**FIGURE 4.** Fish oil decreases ILK promoter activity in human lung carcinoma cells. **A.** The human ILK wild-type and deletion promoter constructs schematics are presented. These regions contain several transcription factor binding sites, including AP-2, Sp1, and NF- $\kappa$ B. **B.** H1838 cells ( $1 \times 10^5$ ) were cotransfected with a wild-type human ILK promoter reporter constructs (shown in **A**) ligated to a luciferase reporter gene and an internal control pRL-TK Synthetic *Renilla* Luciferase Reporter Vector as described in Materials and Methods for 24 h using the Oligofectamine reagent according to the manufacturer's instructions. After 24 h of incubation, cells were treated with vehicle control (Con), rosiglitazone (10  $\mu$ mol/L), fish oil (10  $\mu$ g/mL), linoleic acid (10  $\mu$ mol/L), and WY14643 (15  $\mu$ mol/L) for an additional 24 h. **C.** H1838 cells ( $1 \times 10^5$ ) were cotransfected with a wild-type human ILK promoter reporter construct (shown in **A**) ligated to luciferase reporter gene and an internal control pRL-TK Synthetic *Renilla* Luciferase Reporter Vector as described in Materials and Methods for 24 h using the Oligofectamine reagent according to the manufacturer's instructions. After 24 h of incubation, cells were treated with SB239023 (10  $\mu$ mol/L) for 2 h before exposure of the cells to fish oil (10  $\mu$ g/mL) for an additional 24 h. The ratio of firefly luciferase to *Renilla* luciferase activity was quantified as described in Materials and Methods. Columns, mean of at least three independent experiments for each condition; bars, SD. \*, significant increase of activity compared with controls; \*\*, significance of combination treatment compared with fish oil alone ( $P < 0.05$ ).



**FIGURE 5.** Effect of fish oil on AP-2, Sp1, and NF- $\kappa$ B binding activities. **A.** Oligonucleotides containing the AP-2 (**A**), Sp1 (**B**), and NF- $\kappa$ B (**C**) sites were end labeled with [ $\gamma$ - $^{32}$ P]ATP and incubated with nuclear extracts (5  $\mu$ g) from H1838 cells treated with fish oil (10  $\mu$ g/mL) for an additional 24 h. For competition assays, a molar excess ( $\times 100$ ) of consensus Sp1 (*Cold Sp1*) or NF- $\kappa$ B (*Cold NF- $\kappa$ B*) or AP-2 (*Cold AP-2*) oligonucleotide was added to the binding reaction. Oligonucleotides containing a mutated Sp1 (*Mut Sp1*) or NF- $\kappa$ B (*Mut NF- $\kappa$ B*) or AP-2 (*Mut AP-2*) site that were end labeled with [ $\gamma$ - $^{32}$ P]ATP were used to confirm the binding specificity. **D.** Oligonucleotides containing the AP-2 site were end labeled with [ $\gamma$ - $^{32}$ P]ATP and incubated with nuclear extracts (5  $\mu$ g) from H1838 cells treated with GW9662 (20  $\mu$ M) for 2 h before exposure of the cells to fish oil (10  $\mu$ g/mL) for an additional 24 h. For competition assays, a molar excess ( $\times 100$ ) of consensus AP-2 (*Cold AP-2*) oligonucleotide was added to the binding reaction. Oligonucleotides containing a mutated AP-2 (*Mut AP-2*) site that was end labeled with [ $\gamma$ - $^{32}$ P]ATP were used to confirm the binding specificity. **E.** H1838 cells were treated with fish oil (10  $\mu$ g/mL) for 24 h and cell lysates were immunoprecipitated with either IgG or anti-PPAR $\gamma$ . Immunoprecipitates were separated by SDS-PAGE in 10% Tris-glycine gels and transferred onto nitrocellulose, and Western blotting (WB) was carried out with either anti-PPAR $\gamma$  or anti-AP-2 $\alpha$  antibody.

#### Cell Viability Assay

NSCLC cells ( $10^5$  per well) were cultured with increasing concentrations of fish oil-enriched medium for up to 48 h or were transfected with control or AP-2 $\alpha$  siRNA for 30 h before exposure of the cells to fish oil-enriched medium for an additional 48 h in 96-well plates. In a separate experiment, cells were transfected with inactive (ILK-S343A) or superactive ILK (ILK-S343D) cDNA or with control or AP-2 expression reporter construct SP(RSV)AP-2 (12100; purchased from Addgene, Inc.; ref. 40) using the Oligofectamine reagent (Invitrogen) according to the manufacturer's instructions. After 24 h of incubation, cells were treated with or without fish oil-enriched medium for 48 h. Afterwards, the number of viable cells in culture was determined using the CellTiter-Glo Luminescent Cell Viability Assay kit, which is based on quantitation of the ATP present according to the manufacturer's instructions (Promega).

#### Western Blot Analysis

The procedure was done as previously described (38). Protein concentrations were determined by the Bio-Rad

protein assay. Equal amounts of protein from whole-cell lysates were solubilized in  $2\times$  SDS sample buffer and separated on 8% to 10% SDS-polyacrylamide gels. Blots were incubated with antibodies against PPAR $\gamma$ , ILK, p38 MAPK, and their phosphorylated forms (1:1,000), as well as AP-2 $\alpha$ , AP-2 $\beta$ , and AP-2 $\delta$  (1:500). The blots were washed and followed by incubation with a secondary goat antibody raised against rabbit IgG conjugated to horseradish peroxidase (1:2,000; Cell Signaling). The blots were washed, transferred to freshly made enhanced chemiluminescence solution (Amersham) for 1 min, and exposed to X-ray film. In controls, the primary antibodies were omitted or replaced with a control rabbit IgG.


#### Treatment with AP-2 $\alpha$ siRNA

The AP-2 $\alpha$  siRNA and control nonspecific siRNA oligonucleotides were purchased from Santa Cruz Biotechnology, Inc. For the transfection procedure, cells were grown to 60% confluence, and AP-2 $\alpha$  and control siRNAs were transfected using the Oligofectamine reagent according to the manufacturer's instructions. Briefly, Oligofectamine reagent was

incubated with serum-free medium for 10 min. Subsequently, a mixture of respective siRNA was added. After incubation for 15 min at room temperature, the mixture was diluted with medium and added to each well. The final concentration of siRNAs in each well was 100 nmol/L. After culturing for 30 h, cells were washed and resuspended in fish oil-enriched culture medium for an additional 24 or 48 h for luciferase assays or cell growth assays.

#### Immunoprecipitation Assays

Protein lysates were prepared from cells treated with fish oil for 24 h by extraction in modified radioimmunoprecipitation assay buffer [50 mmol/L Tris (pH 7.4), 0.5% NP40, 0.25% sodium deoxycholate, 125 mmol/L NaCl, 1 mmol/L EDTA, 1 mmol/L EGTA, 1 mmol/L  $\text{Na}_3\text{VO}_4$ , protease inhibitor cocktail] on ice for 1 h. Cells were sonicated for 10 s, lysates were centrifuged at  $12,000 \times g$  for 15 min at  $4^\circ\text{C}$ , and the



**FIGURE 6.** The role of transcription factor AP-2 $\alpha$  in inhibition of ILK expression by fish oil. **A.** Cellular proteins were isolated from H1838 cells treated with fish oil (10  $\mu\text{g}/\text{mL}$ ) for the indicated time period. Afterwards, Western blot analyses were done using polyclonal antibodies against AP-2 $\alpha$ , AP-2 $\beta$ , and AP-2 $\gamma$ . **B.** H1838 cells were treated with GW9662 (20  $\mu\text{mol}/\text{L}$ ) for 2 h before exposure of the cells to fish oil (10  $\mu\text{g}/\text{mL}$ ) for an additional 24 h followed by Western blot analysis for AP-2 $\alpha$  protein. **C.** H1838 cells were treated with SB239023 (10  $\mu\text{mol}/\text{L}$ ) for 2 h before exposure of the cells to fish oil (10  $\mu\text{g}/\text{mL}$ ) for an additional 24 h followed by Western blot analysis for AP-2 $\alpha$  protein. **D.** H1838 cells were transfected with control or AP-2 $\alpha$  siRNA (100 nmol/L) together with a wild-type ILK promoter constructs (-267/+463 bp) for 30 h followed by exposing the cells to fish oil (10  $\mu\text{g}/\text{mL}$ ) for an additional 24 h. **E.** H1838 cells were transfected with a truncated human ILK promoter reporter construct (shown in Fig. 4A) ligated to luciferase reporter gene and an internal control phRL-TK Synthetic *Renilla* Luciferase Reporter Vector as described in Materials and Methods for 24 h using the Oligofectamine reagent according to the manufacturer's instructions. After 24 h of incubation, cells were treated with vehicle control (Con) and fish oil (10  $\mu\text{g}/\text{mL}$ ) for an additional 24 h. **F.** H1838 cells were transfected with control or AP-2 $\alpha$  siRNA (100 nmol/L) for 30 h before exposure of the cells to fish oil (10  $\mu\text{g}/\text{mL}$ ) for an additional 48 h. Afterwards, viable cells were detected using CellTiter-Glo Luminescent Cell Viability Assay kit according to the protocol of the manufacturer. Actin was used as internal control for loading purposes. Western blot result for AP-2 $\alpha$  protein production. **G.** H1838 cells were transfected with control and AP-2 expression reporter constructs [SP(RSV)AP-2] and an internal control phRL-TK Synthetic *Renilla* Luciferase Reporter Vector as described in Materials and Methods for 24 h before exposure of the cells to fish oil (10  $\mu\text{g}/\text{mL}$ ) for an additional 48 h. Western blot results for AP-2 $\alpha$  protein. The ratio of firefly luciferase activity to *Renilla* luciferase activity was quantified as described in Materials and Methods. Columns, mean of at least four independent experiments for each condition; bars, SD. \*, significant increase of activity compared with controls; \*\*, significance of combination treatment compared with the fish oil alone ( $P < 0.05$ ).



supernatant was removed for immunoprecipitation. Samples containing 200 µg proteins were precleared for 30 min with 30 µL of protein A/G Plus-agarose (Santa Cruz Biotechnology) and incubated for 1 h at 4°C with the appropriate antibodies (anti-PPAR $\gamma$ ) or normal IgG preabsorbed to protein A/G Plus-agarose. Immune complexes were collected after incubation overnight at 4°C and washed once with lysis buffer and three times with PBS. Protein was eluted by boiling in 25 µL 2 $\times$  SDS sample buffer [125 mmol/L Tris (pH 6.8), 10% 2-mercaptoethanol, 4% SDS, 20% glycerol], and eluted proteins were analyzed by SDS-PAGE and Western blotting using the appropriate antibodies described above.

#### Transient Transfection Assays

The wild-type human ILK (pILK-Pr) and deletion ILK promoter constructs (dILK-pr) ligated to the luciferase reporter gene were a gift from Drs. Michalik and Desvergne (Center for Integrative Genomics, University of Lausanne, Lausanne, Switzerland) and have been reported previously (41). The ILK promoter construct contains ~730 bp of the 5' flanking region of the human *ILK* gene connected to the pGL3 Basic Luciferase reporter vector (Promega). Briefly, NSCLC cells were seeded at a density of  $5 \times 10^5$  per well in six-well dishes and grown to 50% to 60% confluence. For each well, 2 µg of the above ILK plasmid DNA constructs, with or without 0.2 µg of the internal control phRL-TK Synthetic *Renilla* Luciferase Reporter Vector, were cotransfected into the cells using the Oligofectamine reagent according to the manufacturer's instructions as described in our earlier work (42). After 24 h of incubation, cells were treated with or without rosiglitazone, linoleic acid, WY14643, or fish oil for 24 h or with SB239023 for 2 h before exposure of the cells to fresh oil-enriched medium for an additional 24 h. The preparation of cell extracts and measurement of luciferase activities were carried out using the Dual-Luciferase Reporter kit according to recommendations by the manufacturer (Promega). The assays for firefly luciferase activity and *Renilla* luciferase activity were done sequentially in a Labsystems Luminoskan Ascent luminometer equipped with dual injectors. Changes in firefly luciferase activity were calculated and plotted after normalization with changes in *Renilla* luciferase activity within the same sample.

#### Electrophoretic Mobility Shift Assays

Electrophoretic mobility shift assay experiments were done as described before (39). The oligonucleotides used as probes were wild-type AP-2 (5'-TCCTCCCCGCTCCGC-3') and mutant AP-2 (5'-TCCTCttGCCTCCGC-3'), wild-type Sp1 (5'-GGCCCCACGGGCGGG-3') and mutant Sp1 (5'-GGCCCCACGGtGGG-3'), and wild-type NF- $\kappa$ B (5'-ACGGGAGTCCCCG-3') and mutant NF- $\kappa$ B (5'-ACGtAGTCCCCG-3'), which were based on the ILK promoter sequences (33) and consensus AP-2 binding motifs (5'-GATCGAACTGACCGCCCCGGCCCGT-3'). The complementary oligonucleotides were annealed and purified following the manufacturer's protocol. The AP-2, Sp1, and NF- $\kappa$ B oligonucleotides were end labeled with [ $\gamma$ -<sup>32</sup>P]ATP using T4 polynucleotide kinase as recommended by the manufacturer. Nuclear proteins (5 µg) were first incubated under binding

conditions [10 mmol/L HEPES, Tris-HCl (pH 7.9), 50 mmol/L KCl, 0.1 mmol/L EDTA, 1 mmol/L DTT, 12% (v/v) glycerol, and 2 µg poly(deoxyinosinic-deoxycytidylic acid)] for 10 min, and then [ $\gamma$ -<sup>32</sup>P]ATP probe was added for another 20 min at room temperature in a final volume of 20 µL. For cold competition, a 100-fold excess of the respective unlabeled consensus oligonucleotides was incubated for 15 min before adding the probe. The same amount of mutated oligonucleotides mixed with the probe was used as another control. All of these were done in the same binding conditions as described before. Afterwards, the protein-DNA complexes were electrophoresed on a native 4.5% polyacrylamide gel at 150 V using 1 $\times$  Tris-glycine buffer. Each gel was then dried and subjected to autoradiography at -80°C.

#### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

#### Acknowledgments

We thank Drs. Michalik and Desvergne for providing the ILK promoter constructs and Dr. Shi-Yong Sun (Winship Cancer Institute, Emory University) for providing H1792 and H522 cells.

#### References

- Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007;57:43–66.
- Collins LG, Haines C, Perkel R, Enck RE. Lung cancer: diagnosis and management. *Am Fam Physician* 2007;75:56–63.
- Reddy BS, Patlolla JM, Simi B, Wang SH, Rao CV. Prevention of colon cancer by low doses of celecoxib, a cyclooxygenase inhibitor, administered in diet rich in  $\omega$ -3 polyunsaturated fatty acids. *Cancer Res* 2005;65:8022–7.
- Nowak J, Weylandt KH, Habel P, et al. Colitis-associated colon tumorigenesis is suppressed in transgenic mice rich in endogenous *n*-3 fatty acids. *Carcinogenesis* 2007;28:1991–5.
- Hardman WE. Dietary canola oil suppressed growth of implanted MDA-MB 231 human breast tumors in nude mice. *Nutr Cancer* 2007;57:177–83.
- Sauer LA, Blask DE, Dauchy RT. Dietary factors and growth and metabolism in experimental tumors. *J Nutr Biochem* 2007;18:637–49.
- Trombetta A, Maggiora M, Martinasso G, Cotogni P, Canuto RA, Muzio G. Arachidonic and docosahexaenoic acids reduce the growth of A549 human lung-tumor cells increasing lipid peroxidation and PPARs. *Chem Biol Interact* 2007; 165:239–50.
- Kliwer SA, Willson TM. The nuclear receptor PPAR $\gamma$ —bigger than fat. *Curr Opin Genet Dev* 1998;8:576–81.
- Chene G, Dubourdeau M, Balard P, et al. *n*-3 and *n*-6 polyunsaturated fatty acids induce the expression of COX-2 via PPAR $\gamma$  activation in human keratinocyte HaCaT cells. *Biochim Biophys Acta* 2007;1771:576–89.
- Schopfer FJ, Lin Y, Baker PR, et al. Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor  $\gamma$  ligand. *Proc Natl Acad Sci U S A* 2005;102:2340–5.
- Grommes C, Landreth GE, Heneka MT. Antineoplastic effects of peroxisome proliferator-activated receptor  $\gamma$  agonists. *Lancet Oncol* 2004;5:419–29.
- Hannigan G, Troussard AA, Dedhar S. Integrin-linked kinase: a cancer therapeutic target unique among its ILK. *Nat Rev Cancer* 2005;5:51–63.
- Liu J, Costello PC, Pham NA, et al. Integrin-linked kinase inhibitor KP-392 demonstrates clinical benefits in an orthotopic human non-small cell lung cancer model. *J Thorac Oncol* 2006;1:771–9.
- Cordes N. Overexpression of hyperactive integrin-linked kinase leads to increased cellular radiosensitivity. *Cancer Res* 2004;64:5683–92.
- Hannigan GE, Leung-Hagsteejn C, Fitz-Gibbon L, et al. Regulation of cell adhesion and anchorage-dependent growth by a new  $\beta$ 1-integrin-linked protein kinase. *Nature* 1996;379:91–6.
- Koul D, Shen R, Bergh S, et al. Targeting integrin-linked kinase inhibits Akt signaling pathways and decreases tumor progression of human glioblastoma. *Mol Cancer Ther* 2005;4:1681–8.
- Yano M, Matsumura T, Senokuchi T, et al. Statins activate peroxisome

- proliferator-activated receptor  $\gamma$  through extracellular signal-regulated kinase 1/2 and p38 mitogen-activated protein kinase-dependent cyclooxygenase-2 expression in macrophages. *Circ Res* 2007;100:1442–51.
18. Li M, Lee TW, Mok TS, Warner TD, Yim AP, Chen GG. Activation of peroxisome proliferator-activated receptor- $\gamma$  by troglitazone (TGZ) inhibits human lung cell growth. *J Cell Biochem* 2005;96:760–74.
  19. Han S, Zheng Y, Roman J. Rosiglitazone, an agonist of PPAR $\gamma$ , inhibits non-small cell carcinoma cell proliferation in part through activation of tumor sclerosis complex-2. *PPAR Res* 2007;2007:29632.
  20. Costa V, Foti D, Paonessa F, et al. The insulin receptor: a new anticancer target for peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) and thiazolidinedione-PPAR $\gamma$  agonists. *Endocr Relat Cancer* 2008;15:325–35.
  21. Patel KM, Wright KL, Whittaker P, Chakravarty P, Watson ML, Ward SG. Differential modulation of COX-2 expression in A549 airway epithelial cells by structurally distinct PPAR $\gamma$  agonists: evidence for disparate functional effects which are independent of NF- $\kappa$ B and PPAR $\gamma$ . *Cell Signal* 2005;17:1098–110.
  22. Pandey MK, Pant AB, Das M. *In vitro* cytotoxicity of polycyclic aromatic hydrocarbon residues arising through repeated fish fried oil in human hepatoma Hep G2 cell line. *Toxicol In Vitro* 2006;20:308–16.
  23. Istfan NW, Chen ZY, Rex S. Fish oil slows S phase progression and may cause upstream shift of DHFR replication origin ori- $\beta$  in CHO cells. *Am J Physiol Cell Physiol* 2002;283:C1009–24.
  24. Cerchietti LC, Navigante AH, Castro MA. Effects of eicosapentaenoic and docosahexaenoic *n*-3 fatty acids from fish oil and preferential Cox-2 inhibition on systemic syndromes in patients with advanced lung cancer. *Nutr Cancer* 2007;59:14–20.
  25. Sawai H, Okada Y, Funahashi H, et al. Integrin-linked kinase activity is associated with interleukin-1 $\alpha$ -induced progressive behavior of pancreatic cancer and poor patient survival. *Oncogene* 2006;25:3237–46.
  26. Imanishi Y, Hu B, Jarzynka MJ, et al. Angiopoietin-2 stimulates breast cancer metastasis through the  $\alpha$ 5 $\beta$ 1 integrin-mediated pathway. *Cancer Res* 2007;67:4254–63.
  27. Okamura M, Yamaji S, Nagashima Y, et al. Prognostic value of integrin  $\beta$ 1-ILK-pAkt signaling pathway in non-small cell lung cancer. *Hum Pathol* 2007;38:1081–91.
  28. Han S, Ritzenthaler JD, Sitaraman SV, Roman J. Fibronectin increases matrix metalloproteinase 9 expression through activation of c-Fos via extracellular-regulated kinase and phosphatidylinositol 3-kinase pathways in human lung carcinoma cells. *J Biol Chem* 2006;281:29614–24.
  29. Marks JL, Gong Y, Chitale D, et al. Novel MEK1 mutation identified by mutational analysis of epidermal growth factor receptor signaling pathway genes in lung adenocarcinoma. *Cancer Res* 2008;68:5524–8.
  30. Allred CD, Talbert DR, Southard RC, Wang X, Kilgore MW. PPAR $\gamma$ 1 as a molecular target of eicosapentaenoic acid in human colon cancer (HT-29) cells. *J Nutr* 2008;138:250–6.
  31. Yee LD, Young DC, Rosol TJ, Vanbuskirk AM, Clinton SK. Dietary (*n*-3) polyunsaturated fatty acids inhibit HER-2/neu-induced breast cancer in mice independently of the PPAR $\gamma$  ligand rosiglitazone. *J Nutr* 2005;135:983–8.
  32. Zhang Y, Ikegami T, Honda A, et al. Involvement of integrin-linked kinase in carbon tetrachloride-induced hepatic fibrosis in rats. *Hepatology* 2006;44:612–22.
  33. Melchior C, Kreis S, Janji B, Kieffer N. Promoter characterization and genomic organization of the gene encoding integrin-linked kinase 1. *Biochim Biophys Acta* 2002;1575:117–22.
  34. Schwartz B, Melnikova VO, Tellez C, et al. Loss of AP-2 $\alpha$  results in deregulation of E-cadherin and MMP-9 and an increase in tumorigenicity of colon cancer cells *in vivo*. *Oncogene* 2007;26:4049–58.
  35. Pellikainen JM, Kosma VM. Activator protein-2 in carcinogenesis with a special reference to breast cancer—a mini review. *Int J Cancer* 2007;120:2061–7.
  36. Fauquette V, Aubert S, Groux-Degroote S, et al. Transcription factor AP-2 $\alpha$  represses both the mucin MUC4 expression and pancreatic cancer cell proliferation. *Carcinogenesis* 2007;28:2305–12.
  37. Choi HJ, Chung TW, Kim SJ, et al. The AP-2 $\alpha$  transcription factor is required for the ganglioside GM3-stimulated transcriptional regulation of a PTEN gene. *Glycobiology* 2008;18:395–407.
  38. Han S, Ritzenthaler JD, Wingerd B, Roman J. Activation of peroxisome proliferator-activated receptor  $\beta/\delta$  (PPAR $\beta/\delta$ ) increases the expression of prostaglandin E2 receptor subtype EP4. The roles of phosphatidylinositol 3-kinase and CCAAT/enhancer-binding protein  $\beta$ . *J Biol Chem* 2005;280:33240–9.
  39. Fox PL, DiCorleto PE. Fish oils inhibit endothelial cell production of platelet-derived growth factor-like protein. *Science* 1988;241:453–6.
  40. Williams T, Tjian R. Analysis of the DNA-binding and activation properties of the human transcription factor AP-2. *Genes Dev* 1991;5:670–82.
  41. Di-Poi N, Tan NS, Michalik L, Wahli W, Desvergne B. Antipoptotic role of PPAR $\beta$  in keratinocytes via transcriptional control of the Akt1 signaling pathway. *Mol Cell* 2002;10:721–33.
  42. Han S, Sidell N, Fisher PB, Roman J. Up-regulation of p21 gene expression by peroxisome proliferator-activated receptor  $\gamma$  in human lung carcinoma cells. *Clin Cancer Res* 2004;10:1911–9.