

Consequences of a Family History of Type 1 and Type 2 Diabetes on the Phenotype of Patients With Type 2 Diabetes

HAIYAN LI, BM
BO ISOMAA, MD
M.-R. TASKINEN, MD, PHD

LEIF GROOP, MD, PHD
TIINAMAJA TUOMI, MD, PHD

OBJECTIVE — To investigate the impact of a family history of type 1 and type 2 diabetes on the phenotype of patients with type 2 diabetes.

RESEARCH DESIGN AND METHODS — In a population-based study, we compared the phenotype in 3 groups of patients with type 2 diabetes. The first group had no family history of diabetes (FH⁻, n = 148); the second group had a family history of type 2 diabetes only (FH^{+ TYPE2}, n = 1,211); and the third group had a family history of both type 1 and type 2 diabetes (FH^{+ MIXED}, n = 240). Furthermore, we studied the frequency of GAD antibodies (GADabs), HLA-DQB1 risk genotypes, and the presence of coronary heart disease (CHD) according to family history in unrelated patients with type 2 diabetes from 787 families (148 FH⁻, 546 FH^{+ TYPE2}, and 93 FH^{+ MIXED}).

RESULTS — Patients with no family history of diabetes were older at the onset of the disease, had a better β -cell function (P = 0.004), and had higher HDL cholesterol concentrations (P = 0.006) than patients with a family history of diabetes. Patients with a family history of only type 2 diabetes had higher BMI and fasting C-peptide concentrations (P = 0.031) but lower frequency of GADab (11 vs. 23%, P = 0.001) and DQB1 risk genotypes (37 vs. 54%, P = 0.003) compared with patients with a family history of both type 1 and type 2 diabetes. In addition, hypertension (P = 0.05) and CHD (P = 0.031) were more common in FH^{+ TYPE2} than in FH^{+ MIXED} patients. In patients <60 years old, a family history of type 1 diabetes was associated with a reduced risk of CHD independent of age, hypertension, and HDL cholesterol concentrations. The results were similar when the GADab⁺ patients were excluded from the analysis.

CONCLUSIONS — A family history of both type 1 and type 2 diabetes had a profound influence on the phenotype of patients with type 2 diabetes, which suggests a genetic interaction between type 1 and type 2 diabetes.

Diabetes Care 23:589–594, 2000

Both type 1 and type 2 diabetes have a strong genetic component, and both types of diabetes cluster in the same families (1–12). Type 2 diabetes is a heterogeneous disease characterized by insulin resistance and impaired β -cell function. Studies in nondiabetic offspring of patients with type 2 diabetes have shown that defects in both insulin action and insulin secretion seem to be inherited (13,14).

From the Diabetes and Endocrine Research Laboratory (H.L., L.G., T.T.), Department of Endocrinology, Lund University, Malmö, Sweden; Jakobstad Hospital (B.I.), Jakobstad; and the Department of Internal Medicine (M.-R.T., T.T.), Helsinki University Central Hospital, Helsinki, Finland.

Address correspondence and reprint requests to Tiinamajja Tuomi, MD, PhD, Wallenberg Laboratory, Department of Endocrinology, Lund University, S-20502 Malmö, Sweden. E-mail: tiinamajja.tuomi@endo.mas.lu.se. Received for publication 25 August 1999 and accepted in revised form 10 January 2000.

Abbreviations: AER, albumin excretion rate; apoB, apolipoprotein B; CHD, coronary heart disease; CV, coefficient of variation; dBp, diastolic blood pressure; FH⁺, family history of diabetes; FH⁻, no family history of diabetes; FH^{+ MIXED}, family history of both type 1 and type 2 diabetes; FH^{+ TYPE2}, family history of type 2 diabetes only; FS, fasting serum; GADab, GAD antibody; MODY, maturity-onset diabetes of the young; OGTT, oral glucose tolerance test; OHA, oral hypoglycemic agents; RR, relative risk; sBP, systolic blood pressure.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

However, in these studies, only a family history of type 2 diabetes has been considered without taking into account the possible effect of a family history of type 1 diabetes. Furthermore, it is not known whether familial patients with type 2 diabetes are phenotypically similar to sporadic patients with type 2 diabetes without a family history of diabetes.

To address these questions, we compared the clinical phenotype in patients with type 2 diabetes who were divided into the following 3 groups: 1) those without a family history of diabetes (FH⁻), 2) those with a family history of type 2 diabetes only (FH^{+ TYPE2}), and 3) those with a family history of both type 1 and type 2 diabetes (FH^{+ MIXED}). In addition, we compared the type 1 diabetes-associated HLA-DQB1 genotype frequency in the 3 groups of patients with type 2 diabetes.

The data emphasize the importance of considering the genetic background of diabetes when studying the phenotype of patients with type 2 diabetes.

RESEARCH DESIGN AND METHODS

Study subjects

The Botnia Study is a population-based study aiming at the identification of genes that increase susceptibility to type 2 diabetes (13). Since 1990, patients with type 2 diabetes and their available family members have been recruited from a total of 1,377 families. The average number of family members was 8, and the average number of diabetic family members was 2.6. Families with maturity-onset diabetes of the young (MODY) were excluded after DNA analysis (15). All available family members were subjected to an oral glucose tolerance test (OGTT). Type 2 diabetes was diagnosed according to the revised World Health Organization criteria (16). Diagnosis of type 1 diabetes was based on initiation of insulin treatment within 6 months after diagnosis and/or a fasting C-peptide concentration <0.2 nmol/l. Patients who did not fulfill these criteria and did not have MODY mutations were considered to

have type 2 diabetes. The protocols were approved by the local ethics committees.

The study included the first consecutive 1,599 patients with type 2 diabetes from 787 families, most of them originating from the western coast of Finland (Botnia). The families comprised 5,670 subjects altogether. The presence of diabetes in the relative was either confirmed with measurements of fasting blood glucose or an OGTT (>90% of subjects) or based on the use of oral hypoglycemic agents (OHA) or insulin. The OGTT was performed for 91% of the nondiabetic subjects. Patients were divided into the following 2 groups according to family history of type 2 diabetes in their first- or second-degree relatives: 1) patients without a family history of diabetes (FH⁻, *n* = 148) and 2) patients with a family history of type 2 diabetes (FH⁺). The FH⁺ group was further subdivided into those with a family history of type 2 diabetes only (FH⁺_{TYPE2}, *n* = 1,211) and those with a family history of both type 1 and type 2 diabetes (FH⁺_{MIXED}, *n* = 240). Patients with a third-degree family history of type 1 diabetes (e.g., cousins) were included in the FH⁺_{MIXED} group, but 76% of the patients in that group had either first- or second-degree relatives with type 1 diabetes. The frequency of HLA-DQB1 genotypes (02/0302, 0302/X, 02/X, and 0602(3)/X) associated with an increased or decreased risk of type 1 diabetes, GAD antibodies (GADabs), insulin treatment, hypertension, or coronary heart disease (CHD) were compared between the unrelated patients with FH⁻ (*n* = 148), FH⁺_{TYPE2} (*n* = 546), and FH⁺_{MIXED} (*n* = 93) diabetes randomly selected from the 787 families (1 patient per family).

A standardized health questionnaire covering the subjects' medical history, including current and previous medication, was completed by specially trained nurses; the information included hypertension, cardiovascular diseases, myocardial infarction, stroke, smoking habits, alcohol consumption, and physical activity, in addition to a family history of diabetes. Combined with the local hospital records, myocardial infarction and stroke (including both ischemic and hemorrhagic) were defined as events requiring hospitalization; CHD was defined as the use of oral nitroglycerine or typical chest pain or a history of previous myocardial infarction. This information on the CHD was validated against electrocardiogram changes compatible with ischemic heart disease (17). Hypertension was defined as systolic blood pressure (SBP) of

≥160 mmHg and/or a diastolic blood pressure (DBP) of ≥90 mmHg or the use of anti-hypertensive drugs.

Microalbuminuria was defined as urinary albumin excretion rate (AER) of 30–300 mg/24 h or 20–200 μg/min in overnight urine collections and macroalbuminuria as AER >300 mg/24 h or >200 μg/min in 2 of 3 urine collections. This information was available for 63% of the subjects.

The subjects were classified as smokers if they were currently smoking or were known to have smoked within 1 year before the interview, or as nonsmokers if they had never smoked at any time or had not smoked for >1 year before the interview. This information was available for 54% of the subjects.

Metabolic measurements

An OGTT was performed for all subjects aged >15 years with fasting blood glucose <10 mmol/l and not treated with insulin. After a 12-h overnight fast, the subjects ingested 75 g of glucose in a volume of 300 ml. Samples for measurements of blood glucose and serum insulin were drawn at -10, 0, 30, 60, and 120 min, and incremental glucose and insulin areas under the curve were calculated. The 30-min incremental insulin area was used to evaluate the early insulin response. Blood glucose was measured with a hexokinase method (Boehringer Mannheim, Mannheim, Germany). Serum insulin concentrations were measured by radioimmunoassay (Pharmacia, Uppsala, Sweden) with an interassay coefficient of variation (CV) of 5%. Fasting serum (FS)-C-peptide concentrations were measured by a radioimmunoassay with an interassay CV of 9% (Human C-peptide RIA Kit; Linco, St. Charles, MO). HbA_{1c} concentration was measured by a high-performance liquid chromatography with a reference value of 5–7%.

The plasma concentrations of cholesterol and triglycerides and separate lipoprotein fractions were determined by enzymatic methods using commercial kits (Boehringer Mannheim). The ratio of total cholesterol to apolipoprotein B (apoB) concentrations was calculated (after converting apoB to mmol/l) and used as a measure of LDL particle size. Urinary AER was measured by immunoturbidimetry from 3 overnight urine collections. GADab were measured by a radio-immunoprecipitation method using ³⁵S-labeled recombinant human GAD₆₅ produced by in vitro transcription/translation (18).

HLA DQB1 genotypes

HLA DQB1 genotyping was performed by polymerase chain reaction for the second exon followed by dot-blot hybridization with sequence-specific oligonucleotide probes labeled with digoxigenin (DIG Oligonucleotide 3'-End Labelling Kit; Boehringer Mannheim) (18). Three DQB1 probes were used to distinguish DQB1 alleles 0201 or 0202 (02), 0302, and 0602 or 0603 [0602(3)] alleles. The genotypes were presented as 02/0302, 0302/X, 02/X, 0602(3)/X, 0302/0602(3), 02/0602(3), and X/X, whereby X could mean either a homozygous allele or any allele other than 02, 0302, or 0602(3). The 02/0302, 0302/X, or 02/X genotypes were considered DQB1-risk genotypes (19).

Statistical analysis

Data are presented as means ± SD or medians (interquartile ranges). The statistical analysis was performed with either Biomedical Data Processing (BMDP, Los Angeles, CA) or Number Cruncher Statistical Systems (NCSS, Kaysville, UT) statistical software. The group frequencies were compared using the χ² or Fisher's exact tests. Differences between group means were tested with analysis of variance or covariance with age, BMI, and waist-to-hip ratio as covariates, or using Mann-Whitney or the Kruskal-Wallis tests where appropriate. Logarithmic transformation was performed to variables with skewed distribution. A multiple logistic regression analysis was carried out with CHD as the dependent variable and age, FS-C-peptide, HDL cholesterol concentrations, and presence of hypertension, DQB1 risk genotype, family history of type 1 diabetes, or family history of type 2 diabetes as independent variables. Serum cholesterol concentration, micro- or macroalbuminuria, or smoking history did not differ between patients with and without CHD and were not included in the regression analysis. Hypertension, DQB1 risk genotypes, and family history of type 1 or type 2 diabetes were treated as binary variables in the analysis (presence = 1, absence = 0).

RESULTS — The FH⁻ patients presented with diabetes at a later age than the FH⁺_{TYPE2} patients (*P* < 0.0001, Table 1). Also, the FH⁻ patients had a lower BMI than the FH⁺_{TYPE2} patients (27.4 ± 4.1 vs. 29.1 ± 4.9 kg/m², *P* < 0.0001). β-Cell function, as judged from FS-C-peptide concentration (0.64 ± 0.38 vs. 0.67 ± 0.35 nmol/l, *P* = 0.020) and incremental early insulin secretion during OGTT

Table 1—Clinical characteristics of patients with type 2 diabetes without family history of diabetes (FH⁻), with family history of type 2 diabetes only (FH⁺_{TYPE2}), or with family history of both type 1 and type 2 diabetes (FH⁺_{MIXED})

	FH ⁻	P	FH ⁺ _{TYPE2}	P	FH ⁺ _{MIXED}
n (M/F)	148 (62/86)		1,211 (539/672)		240 (140/136)
Age (years)	71.3 ± 11.6	<0.0001	64.8 ± 11.1		64.4 ± 12.8
Age at onset (years)	62.9 ± 12.1	<0.0001	55.7 ± 12.2		54.2 ± 13.7
Duration (years)	7.2 (7.0)		8.5 (11.2)		9.1 (11.3)
BMI (kg/m ²)	27.4 ± 4.1	<0.0001	29.1 ± 4.9	0.005	28.1 ± 4.5
Waist-to-hip ratio	0.91 ± 0.08	<0.0001	0.94 ± 0.08		0.93 ± 0.08
Fasting blood glucose (mmol/l)	8.7 ± 3.1		8.7 ± 2.8		8.5 ± 3.1
HbA _{1c} (%)	7.9 ± 1.8		7.6 ± 1.7		7.7 ± 1.8
Fasting insulin (mU/l)	16.6 ± 14.2		18.1 ± 20.4		18.6 ± 23.1
FS-C-peptide (nmol/l)	0.67 ± 0.35	0.020*	0.64 ± 0.38	0.031*	0.54 ± 0.35
30-Min incremental insulin area (mU/l)	412 (485)	0.024*	289 (402)		283 (355)
dBp (mmHg)	81.5 ± 10.0		81.8 ± 10.8		82.2 ± 10.3
sBP (mmHg)	148.7 ± 19.7		144.6 ± 20.7		144.5 ± 18.7
Cholesterol (mmol/l)	5.8 ± 1.2		5.8 ± 1.2		5.8 ± 1.2
Triglycerides (mmol/l)	1.9 ± 1.4		2.0 ± 1.4		1.9 ± 1.5
HDL cholesterol (mmol/l)	1.25 ± 0.30	0.0003	1.16 ± 0.32		1.18 ± 0.32
ApoA1 (mg/dl)	133.1 ± 23.5		133.1 ± 22.1		133.4 ± 23.5
ApoA2 (mg/dl)	34.1 ± 6.3		35.3 ± 6.2		34.3 ± 5.8
ApoB (mg/dl)	97.2 ± 21.3	0.007	103.3 ± 25.2		101.1 ± 24.4
Micro-/macroalbuminuria† (%)	15/1		15/3		12/5
Smoking history (%)	26	0.020	36		28
Treatment: insulin/OHA/diet (%)	17/34/49	<0.001 (df2)	25/41/35	<0.001 (df2)	36/33/30

Data are n, means ± SD, medians (interquartile ranges), or %. *Adjusted for age, BMI, and waist-to-hip ratio. †Microalbuminuria or macroalbuminuria was defined as AER >20 µg/min or >200 µg/min, respectively.

(median [interquartile range]: 289 [402] vs. 412 [485] mU/l, $P = 0.024$) was reduced in FH⁺_{TYPE2} patients compared with FH⁻ patients; the lowest concentrations were seen in patients with a family history of both type 1 and type 2 diabetes (FH⁺_{MIXED}: 283 [355] mU/l; $P = 0.002$ vs. FH⁻ patients) (Table 1). HDL cholesterol concentrations were lower in FH⁺_{TYPE2} patients than in FH⁻ patients (1.16 ± 0.32 vs. 1.25 ± 0.30 mmol/l, $P = 0.0003$) and no difference was seen between FH⁺_{TYPE2} and FH⁺_{MIXED}. ApoB concentrations were higher (Table 1) and the total cholesterol/apoB ratio (used as a measure of LDL particle size) was lower in FH⁺_{TYPE2} patients than in FH⁻ patients ($2,044 \pm 444$ vs. $2,146 \pm 442$, $P = 0.001$).

GADabs were more common in patients with a family history of both type 1 and type 2 diabetes (23%) than in those with a family history of only type 2 diabetes (11%) or without a family history of diabetes (9%) ($P < 0.001$). Also, the frequency of DQB1 risk genotypes was higher in FH⁺_{MIXED} patients (54%) than in FH⁺_{TYPE2} patients (37%, $P = 0.003$) or FH⁻ patients (36%, $P < 0.008$) (Table 2). To exclude the possibility that the phenotypic differences were a result of patients with a latent auto-

immune form of type 1 diabetes, we repeated the analysis after exclusion of the GADab⁺ patients; this analysis yielded virtually identical results (Table 3).

The frequency of hypertension (47 vs. 58%, $P = 0.05$) and CHD (19 vs. 30%, $P = 0.031$) was lower in FH⁺_{MIXED} patients than in FH⁺_{TYPE2} patients. There was no

difference in frequency of micro- or macroalbuminuria or smoking history between the FH⁺_{MIXED} and the FH⁺_{TYPE2} patients (Table 1). Compared with the risk in FH⁻ patients, the relative risk (RR) of CHD was decreased in FH⁺_{MIXED} patients (RR = 0.57, $P = 0.053$) and increased in FH⁺_{TYPE2} patients (RR = 1.53, $P = 0.052$). In a mul-

Table 2—Frequency of HLA-DQB1 genotypes, GADab, mode of treatment, albuminuria, smoking, hypertension, and CHD in 787 unrelated patients with type 2 diabetes randomly selected from the families (1 per family)

	Family history of diabetes				
	FH ⁻ (%)	P	FH ⁺ _{TYPE2} (%)	P	FH ⁺ _{MIXED} (%)
HLA DQB1 genotype					
Risk	36	NS	37	0.003	54
Protective	35	0.064	26	0.068	16
GADab ⁺	9	NS	11	0.001	23
Insulin/OHA/diet	17/34/49	—	23/42/35	<0.0001	44/31/24
Albuminuria (micro-/macro-)	16 (15/1)	NS	17 (15/2)	NS	18 (9/9)
Smoking	26	NS	33	NS	32
Hypertension	59	NS	58	0.050	47
CHD	29	NS	30	0.031	19

Data are %. The risk genotypes include 02/0302 or 0302/X or 02/X, and the protective genotypes 0602(3)/X (X denotes either a homozygous allele or any allele other than 02, 0302, or 0602[3]). $n = 462$ (129 FH⁻, 241 FH⁺_{TYPE2}, and 92 FH⁺_{MIXED} patients).

Family history of type 1 and type 2 diabetes

Table 3—GADab⁻ patients with type 2 diabetes divided into those without family history of diabetes (FH⁻), with family history of type 2 diabetes only (FH⁺_{TYPE2}), or with family history of both type 1 and type 2 diabetes (FH⁺_{MIXED})

	FH ⁻	P	FH ⁺ _{TYPE2}	P	FH ⁺ _{MIXED}
n (M/F)	134 (53/81)		1,114 (505/609)		198 (85/113)
Age (years)	71.0 ± 11.3	<0.0001	64.9 ± 11.0		65.5 ± 12.6
Age at onset (years)	62.7 ± 11.8	<0.0001	55.9 ± 12.0		55.3 ± 13.6
Duration (years)	7.3 (6.2)		8.4 (11.0)		9.1 (12.1)
BMI (kg/m ²)	24.4 ± 4.0	0.0001	29.2 ± 4.9	0.038	28.4 ± 4.2
Waist-to-hip ratio	0.90 ± 0.08	<0.0001	0.93 ± 0.08		0.93 ± 0.08
Fasting blood glucose (mmol/l)	8.6 ± 2.9		8.6 ± 2.7		8.3 ± 2.8
HbA _{1c} (%)	7.8 ± 1.8		7.5 ± 1.6		7.6 ± 1.8
Fasting insulin (mU/l)	15.6 ± 12.2		17.7 ± 18.1		17.9 ± 19.9
FS-C-peptide (nmol/l)	0.66 ± 0.35		0.66 ± 0.38		0.61 ± 0.35
30-Min incremental insulin area (mU/l)	460 (484)	0.009*	288 (395)		324 (341)
dBp (mmHg)	82.2 ± 10.2		81.9 ± 10.7		82.5 ± 10.3
sBP (mmHg)	149.4 ± 19.8		146.8 ± 20.6		145.5 ± 18.9
Cholesterol (mmol/l)	5.8 ± 1.2		5.8 ± 1.1		5.9 ± 1.2
Triglyceride (mmol/l)	1.9 ± 1.4		2.1 ± 1.6		2.0 ± 1.3
HDL cholesterol (mmol/l)	1.24 ± 0.28	0.0003	1.15 ± 0.30		1.15 ± 0.29
ApoB (mg/dl)	97.9 ± 21.3	0.018	103.9 ± 25.3		103.4 ± 24.8
Micro-/macroalbuminuria† (%)	14/1		15/3		13/6
Smoking (n [%])	33 (26)	0.019	197 (36)	0.041	25 (26)
Insulin/OHA/diet (%)	16/38/48	0.02 (df 2)	22/41/36	0.02 (df 2)	32/34/34
Hypertension (n [%])	83 (62)		632 (57)		105 (53)
CHD (n [%])	40 (30)		349 (32)	0.027	47 (24)

Data are n, n (%), means ± SD, or medians (interquartile ranges). *Adjusted for age, BMI, and waist-to-hip ratio. †Microalbuminuria or macroalbuminuria was defined as AER >20 µg/min or >200 µg/min, respectively.

multiple logistic regression analysis, we tested whether age, FS-C-peptide and HDL cholesterol concentrations, hypertension, DQB1 risk genotype, and family history of type 1 and/or type 2 diabetes were associated with CHD (Table 4). Age, hyperten-

sion, and HDL cholesterol concentrations were independent risk factors for CHD in patients with type 2 diabetes. In addition, in patients <60 years of age, a family history of type 1 diabetes was associated with a reduced risk of CHD (P = 0.033). The

inclusion of cholesterol, microalbuminuria, or smoking history in the regression analysis did not change the results (data not shown).

CONCLUSIONS — The study emphasizes the importance of not only a family history of diabetes but also the type of diabetes as determinants of the phenotype of patients with type 2 diabetes. Without knowledge about the underlying genetic defects, familial type 2 diabetes cannot be distinguished as a different disease entity from sporadic type 2 diabetes. However, the data clearly demonstrate that familial type 2 diabetes is associated with a more severe phenotype (i.e., earlier age at onset, more severe β-cell dysfunction, and more features of the metabolic syndrome) compared with sporadic patients with type 2 diabetes. Furthermore, among the patients with a family history of type 2 diabetes, the introduction of type 1 diabetes in the family history markedly “changed the picture.” Type 2 patients from the mixed type 1/type 2 diabetes families had an increased frequency of type 1 diabetes-associated HLA-DQB1 genotypes and GADabs but lower frequency of hypertension and cardiovas-

Table 4—Multiple regression analysis of factors associated with coronary heart disease in patients with type 2 diabetes

Variables	Regression coefficient	SE	P
All patients (n = 301)			
Age (years)	0.0466	0.0135	0.0005
Hypertension	0.9594	0.2955	0.0011
HDL cholesterol	-1.6199	0.4987	0.0011
Age <60 years (n = 82)			
Hypertension	1.8418	0.8422	0.0287
HDL cholesterol	-5.7977	2.1783	0.0077
Family history of type 1 diabetes	-2.5522	1.2034	0.0339
Age >60 years (n = 219)			
Age (years)	0.0338	0.0204	0.0969
Hypertension	0.8562	0.3246	0.0083
HDL cholesterol	-1.2366	0.5200	0.0174

The following variables were included in the analysis: age, FS-C-peptide concentration, HDL cholesterol concentrations, presence of hypertension, DQB1 risk genotype, family history of type 1 diabetes, and family history of type 2 diabetes.

cular disease. They also required insulin therapy more often than patients with a family history of type 2 diabetes only.

The difference in age at onset between familial and sporadic patients with type 2 diabetes suggests that if the genetic load for type 2 diabetes is low, the manifestation of type 2 diabetes is delayed. Only a few studies have reported on the influence of a family history of type 2 diabetes on the phenotype of patients with type 2 diabetes (20–22). However, either the studies have included a relatively small number of patients with type 2 diabetes or the patients have had a mild form of diabetes (22). On the other hand, several studies have examined the effect of a family history of diabetes on nondiabetic first-degree relatives (13,14,22,23). These individuals have in general displayed early features of the metabolic syndrome, such as abdominal obesity, elevated blood pressure, and dyslipidemia consistent with the findings in the diabetic probands in the present study (13). The study also demonstrates that family history of type 1 diabetes changes the phenotype of type 2 diabetic patients toward a more type 1-like phenotype. However, this phenotype with late onset and high BMI differs from that in classical type 1 diabetes. Although the patients with type 2 diabetes from the mixed type 1/type 2 families had an increased frequency of type 1 diabetes-associated allele, HLA-DQB1*0302, the genotype conferring the highest risk for classical type 1 diabetes, 0201/0302, was rare among them (data not shown). This is compatible with data in type 1 diabetes that 0201/0302 genotype would predispose toward a more severe autoimmune destruction of the β -cells than the 0302/X genotype (24,25). We do not know whether the patients with type 2 diabetes show signs of insulinitis, but the increased frequency of GADabs, observed also in other populations (26), points at the coexistence of pancreatic autoimmunity. Of note, the present study demonstrates that although the frequency of GADab is higher in the mixed families than in the other type 2 families, the phenotypic differences associated with the type 1 diabetes family history are not restricted to the GADab⁺ patients. The 0302 allele is in linkage disequilibrium with the serologically determined HLA-antigen DR4. An increased frequency of DR4 or DQB1*0302 has been shown in type 2 diabetic patients (27–29), particularly in type 2 patients with impaired β -cell function and increased need for insulin

therapy (18,27). A higher than expected transmission of DR4 from patients with type 2 diabetes to offspring with type 1 diabetes has also been shown (6).

The patients with type 2 diabetes from the mixed type 1/type 2 families had less cardiovascular disease than the common type 2 diabetic patients. There is other information to support this finding. An increased prevalence of type 1 diabetes-associated HLA haplotypes and HLA-DR4 has been reported in elderly Finnish men with type 2 diabetes (29), suggesting that the HLA-DR4 antigen may be associated with a longer survival in patients with type 2 diabetes. In support of this, HLA-DR4 was increased in patients with type 2 diabetes who survived during a 10-year follow-up period (30).

In conclusion, the study demonstrates a profound influence of a family history of type 1 and type 2 diabetes on the phenotype and the outcome of patients with type 2 diabetes. The finding also points at a genetic interaction between type 1 and type 2 diabetes and the importance of a shared genetic background.

Acknowledgments — This study was financially supported by the Sigrid Juselius Foundation, the Pahlsson Foundation, the Medical Faculty of Lund University, the Malmö University Hospital, the Swedish Medical Doctors Association, the Crafoord Foundation, the Ernhold Lundström Foundation, and the Finnish Medical Society (B.I.). The Botnia study is supported by grants from the Sigrid Juselius Foundation, JDF Wallenberg, EC (BM4-CT95-0662), Swedish Medical Research Foundation, Academy of Finland, Finnish Diabetes Research Society, Swedish Diabetes Association, and Novo Nordisk Foundation.

Anita Nilsson and Britt Bruveris-Svenburg are acknowledged for technical assistance. The Botnia Research Group is acknowledged for recruiting and clinically studying the subjects.

References

- Gottlieb MS: Diabetes in offspring and siblings of juvenile- and maturity-onset-type diabetics. *J Chronic Dis* 33:331–339, 1980
- Chern MM, Anderson VE, Barbosa J: Empirical risk for insulin-dependent diabetes (IDD) in sibs: further definition of genetic heterogeneity. *Diabetes* 31:1115–1118, 1982
- Wagener DK, Sacks JM, LaPorte RE, MacGregor JM: The Pittsburgh study of insulin-dependent diabetes mellitus: risk for diabetes among relatives of IDDM. *Diabetes* 31:136–144, 1982
- Dahlquist G, Blom L, Tuvemo T, Nystrom L, Sandstrom A, Wall S: The Swedish childhood diabetes study: results from a nine year case register and a one year case-referent study indicating that type 1 (insulin-dependent) diabetes mellitus is associated with both type 2 (non-insulin-dependent) diabetes mellitus and autoimmune disorders. *Diabetologia* 32:2–6, 1989
- Quatraro A, Consoli G, Magno M, Caretta F, Ceriello A, Giugliano D: Analysis of diabetic family connection in subjects with insulin-dependent diabetes mellitus (IDDM). *Diabete Metab* 16:449–452, 1990
- Rich SS, Panter SS, Goetz FC, Hedlund B, Barbosa J: Shared genetic susceptibility of type 1 (insulin-dependent) and type 2 (non-insulin-dependent) diabetes mellitus: contributions of HLA and haptoglobin. *Diabetologia* 34:350–355, 1991
- Pozzilli P, Visalli N, Signore A, Andreani D: Clinical remission in patients with IDDM and family history of NIDDM (Letter; Comment). *Lancet* 337:1165, 1991
- Teupe B, Bergis K: Epidemiological evidence for “double diabetes” (Letter; Comment). *Lancet* 337:361–362, 1991
- Quatraro A, Consoli G, Ceriello A, Giugliano D: Heterogeneity of diabetes mellitus (Letter). *Diabetologia* 30:678–679, 1987
- Landin-Olsson M, Karlsson FA, Lernmark A, Sundkvist G, and the Diabetes Incidence Study in Sweden Group: Islet cell and thyrogastroic antibodies in 633 consecutive 15- to 34-yr-old patients in the Diabetes Incidence Study in Sweden. *Diabetes* 41:1022–1027, 1992
- Carel JC, Boitard C, Bougneres PF: Decreased insulin response to glucose in islet cell antibody-negative siblings of type 1 diabetic children. *J Clin Invest* 92:509–513, 1993
- Li H, Lindholm E, Groop L, Tuomi T: High prevalence of families with both type 1 and type 2 diabetes (Abstract). *Diabetologia* 41:A70, 1998
- Groop L, Forsblom C, Lehtovirta M, Tuomi T, Karanko S, Nissen M, Ehrnstrom BO, Forsen B, Isomaa B, Snickars B, Taskinen MR: Metabolic consequences of a family history of NIDDM (the Botnia study): evidence for sex-specific parental effects. *Diabetes* 45:1585–1593, 1996
- Vauhkonen I, Niskanen L, Vanninen E, Kainulainen S, Uusitupa M, Laakso M: Defects in insulin secretion and insulin action in non-insulin-dependent diabetes mellitus are inherited: metabolic studies on offspring of diabetic probands. *J Clin Invest* 101:86–96, 1998
- Lehto M, Tuomi T, Mahtani MM, Widen E, Forsblom C, Sarelin L, Gullstrom M, Isomaa B, Lehtovirta M, Hyrkkö A, Kaminen T, Orho M, Manley S, Turner RC, Brettnin T, Kirby A, Thomas J, Duyk G, Lander E, Taskinen MR, Groop L: Characterization of

- the MODY3 phenotype: early-onset diabetes caused by an insulin secretion defect. *J Clin Invest* 99:582–591, 1997
16. Alberti KG, Zimmet PZ: Definition, diagnosis and classification of diabetes mellitus and its complications. I. Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation (Comments). *Diabet Med* 15:539–553, 1998
 17. Isomaa B, Tuomi T, Almgren P, Groop LC, Henricsson M, Sarelin L, Taskinen MR: Chronic complications in patients with slowly progressing autoimmune type 1 Diabetes (LADA). *Diabetes Care* 22:1347–1353, 1999
 18. Tuomi T, Carlsson A, Li H, Isomaa B, Mietinen A, Nilsson A, Nissen M, Ehrnstrom BO, Forsen B, Snickars B, Lahti K, Forsblom C, Saloranta C, Taskinen MR, Groop LC: Clinical and genetic characteristics of type 2 diabetes with and without GAD antibodies. *Diabetes* 48:150–157, 1999
 19. Veijola R, Reijonen H, Vahasalo P, Sabbah E, Kulmala P, Ilonen J, Akerblom HK, Knip M: HLA-DQB1-defined genetic susceptibility, beta cell autoimmunity, and metabolic characteristics in familial and nonfamilial insulin-dependent diabetes mellitus: Childhood Diabetes in Finland (DiMe) Study Group. *J Clin Invest* 98: 2489–2495, 1996
 20. Leonetti DL, Fujimoto WY, Wahl PW: Early-life background and the development of non-insulin-dependent diabetes mellitus. *Am J Phys Anthropol* 79:345–355, 1989
 21. Boer JM, Feskens EJ, Kromhout D: Characteristics of non-insulin-dependent diabetes mellitus in elderly men: effect modification by family history. *Int J Epidemiol* 25:394–402, 1996
 22. Grill V, Persson G, Carlsson S, Norman A, Alvarsson M, Ostensson CG, Svanstrom L, Efendic S: Family history of diabetes in middle-aged Swedish men is a gender unrelated factor which associates with insulinopenia in newly diagnosed diabetic subjects. *Diabetologia* 42:15–23, 1999
 23. Ishikawa M, Pruneda ML, Adams-Huet B, Raskin P: Obesity-independent hyperinsulinemia in nondiabetic first-degree relatives of individuals with type 2 diabetes. *Diabetes* 47:788–792, 1998
 24. Deschamps I, Boitard C, Hors J, Busson M, Marcelli Barge A, Mogenet A, Robert JJ: Life table analysis of the risk of type 1 (insulin-dependent) diabetes mellitus in siblings according to islet cell antibodies and HLA markers: an 8-year prospective study. *Diabetologia* 35:951–957, 1992
 25. Schenker M, Hummel M, Ferber K, Walter M, Ziegler AG: Early expression and high prevalence of islet autoantibodies for DR3/4 heterozygous and DR4/4 homozygous offspring of parents with type 1 diabetes: the German BABYDIAB Study. *Diabetologia* 42: 671–677, 1999
 26. Turner R, Stratton I, Horton V, Manley S, Zimmet P, Mackay IR, Shattock M, Bottazzo GF, Holman R: UKPDS: autoantibodies to islet-cell cytoplasm and glutamic acid decarboxylase for prediction of insulin requirement in type 2 diabetes. *Lancet* 350: 1288–1293, 1997
 27. Groop L, Groop PH, Koskimies S: Relationship between B-cell function and HLA antigens in patients with type 2 (non-insulin-dependent) diabetes. *Diabetologia* 29:757–760, 1986
 28. Rich SS, French LR, Sprafka JM, Clements JP, Goetz FC: HLA-associated susceptibility to type 2 (non-insulin-dependent) diabetes mellitus: the Wadena City Health Study. *Diabetologia* 36:234–238, 1993
 29. Tuomilehto-Wolf E, Tuomilehto J, Hitman GA, Nissinen A, Stengard J, Pekkanen J, Kivinen P, Kaarsalo E, Karvonen MJ: Genetic susceptibility to non-insulin dependent diabetes mellitus and glucose intolerance are located in HLA region (Comments). *Br Med J* 307:155–159, 1993
 30. Forsblom CM, Sane T, Groop PH, Totterman KJ, Kallio M, Saloranta C, Laasonen L, Summanen P, Lepantalo M, Laatikainen L, Matikainen E, Teppo AM, Koskimies S, Groop L: Risk factors for mortality in Type II (non-insulin-dependent) diabetes: evidence of a role for neuropathy and a protective effect of HLA-DR4. *Diabetologia* 41: 1253–1262, 1998